

Top 100 SAS Variant-Disease Hits

Ranked by unmet-need index, L2G score, and SAS global MAF. Generated from
ot_sas_gi_vep115_plof_deleterious_missense_disease_gene_variant_map.india_exclusive.score_ge_0.15.parquet.

Table of Contents

1. 1. PLCE1 - thoracic aortic aneurysm L2G 0.851; unmet 5.73
2. 2. ELN - thoracic aortic aneurysm L2G 0.801; unmet 5.73
3. 3. ULK4 - thoracic aortic aneurysm L2G 0.774; unmet 5.73
4. 4. ABO - pancreatic adenocarcinoma L2G 0.836; unmet 5.35
5. 5. HNF1B - squamous cell lung carcinoma L2G 0.865; unmet 5.31
6. 6. CYP2A6 - squamous cell lung carcinoma L2G 0.238; unmet 5.31
7. 7. ERC1 - squamous cell lung carcinoma L2G 0.187; unmet 5.31
8. 8. ABO - Ischemic stroke L2G 0.961; unmet 5.27
9. 9. HDLBP - Ischemic stroke L2G 0.763; unmet 5.27
10. 10. CUX2 - Ischemic stroke L2G 0.717; unmet 5.27
11. 11. SPPL2A - Ischemic stroke L2G 0.45; unmet 5.27
12. 12. ADH5 - Ischemic stroke L2G 0.334; unmet 5.27
13. 13. MMP3 - Ischemic stroke L2G 0.277; unmet 5.27
14. 14. ICA1L - Ischemic stroke L2G 0.231; unmet 5.27
15. 15. PLCE1 - esophageal squamous cell carcinoma L2G 0.294; unmet 5.25
16. 16. WDR72 - chronic kidney disease L2G 0.963; unmet 5.24
17. 17. SHROOM3 - chronic kidney disease L2G 0.885; unmet 5.24
18. 18. PDILT - chronic kidney disease L2G 0.884; unmet 5.24
19. 19. TCF7L2 - chronic kidney disease L2G 0.884; unmet 5.24
20. 20. BCAS3 - chronic kidney disease L2G 0.639; unmet 5.24
21. 21. C6orf118 - chronic kidney disease L2G 0.525; unmet 5.24
22. 22. PCSK9 - chronic kidney disease L2G 0.458; unmet 5.24
23. 23. IVNS1ABP - chronic kidney disease L2G 0.452; unmet 5.24
24. 24. CYP1A2 - chronic kidney disease L2G 0.402; unmet 5.24
25. 25. DAB2 - chronic kidney disease L2G 0.318; unmet 5.24
26. 26. DHRS11 - chronic kidney disease L2G 0.269; unmet 5.24
27. 27. OR52N2 - chronic kidney disease L2G 0.218; unmet 5.24
28. 28. GAB2 - chronic kidney disease L2G 0.197; unmet 5.24
29. 29. PSEN1 - intracranial hemorrhage L2G 0.751; unmet 5.2
30. 30. EXOC4 - intracranial hemorrhage L2G 0.711; unmet 5.2
31. 31. ANKS1B - intracranial hemorrhage L2G 0.621; unmet 5.2
32. 32. WWOX - intracranial hemorrhage L2G 0.611; unmet 5.2
33. 33. ACACA - intracranial hemorrhage L2G 0.594; unmet 5.2
34. 34. PSPH - intracranial hemorrhage L2G 0.504; unmet 5.2
35. 35. TMEM130 - intracranial hemorrhage L2G 0.395; unmet 5.2

36. 36. RTKL1 - glioblastoma multiforme L2G 0.451; unmet 5.07
37. 37. HNF1B - ovarian carcinoma L2G 0.865; unmet 5.07
38. 38. ABO - ovarian carcinoma L2G 0.799; unmet 5.07
39. 39. HOXD3 - ovarian carcinoma L2G 0.739; unmet 5.07
40. 40. POU5F1B - ovarian carcinoma L2G 0.54; unmet 5.07
41. 41. ATP6V0A1 - ovarian carcinoma L2G 0.357; unmet 5.07
42. 42. SPPL2C - ovarian carcinoma L2G 0.234; unmet 5.07
43. 43. MAST4 - bullous pemphigoid L2G 0.871; unmet 5.07
44. 44. PARD3 - cardiac arrest L2G 0.877; unmet 5.03
45. 45. IGSF5 - cardiac arrest L2G 0.758; unmet 5.03
46. 46. THSD7B - cardiac arrest L2G 0.752; unmet 5.03
47. 47. PREX2 - cardiac arrest L2G 0.726; unmet 5.03
48. 48. FMN2 - cardiac arrest L2G 0.69; unmet 5.03
49. 49. ADCY5 - cardiac arrest L2G 0.641; unmet 5.03
50. 50. SPDYE21 - cardiac arrest L2G 0.539; unmet 5.03
51. 51. OR10AD1 - cardiac arrest L2G 0.226; unmet 5.03
52. 52. LMTK3 - cardiac arrest L2G 0.214; unmet 5.03
53. 53. TP63 - lung adenocarcinoma L2G 0.906; unmet 5
54. 54. SECISBP2L - lung adenocarcinoma L2G 0.884; unmet 5
55. 55. HNF1B - lung adenocarcinoma L2G 0.865; unmet 5
56. 56. FOXP4 - lung adenocarcinoma L2G 0.575; unmet 5
57. 57. TCF7L2 - lung adenocarcinoma L2G 0.435; unmet 5
58. 58. RTKL1 - lung adenocarcinoma L2G 0.277; unmet 5
59. 59. CYP2A6 - lung adenocarcinoma L2G 0.238; unmet 5
60. 60. GPC5 - gastric cancer L2G 0.869; unmet 5
61. 61. POU5F1B - gastric cancer L2G 0.828; unmet 5
62. 62. HNF1B - gastric cancer L2G 0.71; unmet 5
63. 63. STPG2 - gastric cancer L2G 0.342; unmet 5
64. 64. NR2E3 - gastric cancer L2G 0.252; unmet 5
65. 65. L1TD1 - neuroblastoma L2G 0.931; unmet 4.96
66. 66. KANSL1L - neuroblastoma L2G 0.692; unmet 4.96
67. 67. WRN - pneumococcal pneumonia L2G 0.585; unmet 4.94
68. 68. ASB5 - pneumococcal pneumonia L2G 0.153; unmet 4.94
69. 69. ACKR1 - microscopic polyangiitis L2G 0.889; unmet 4.91
70. 70. ADH1B - oropharynx cancer L2G 0.965; unmet 4.86
71. 71. INPP5D - oropharynx cancer L2G 0.745; unmet 4.86
72. 72. ANKS1A - oropharynx cancer L2G 0.612; unmet 4.86
73. 73. CAPN13 - oropharynx cancer L2G 0.502; unmet 4.86
74. 74. SULT1C4 - oropharynx cancer L2G 0.353; unmet 4.86
75. 75. GPR27 - oropharynx cancer L2G 0.197; unmet 4.86
76. 76. RTL1 - oropharynx cancer L2G 0.168; unmet 4.86
77. 77. DPY19L4 - polyarteritis nodosa L2G 0.73; unmet 4.85

78. 78. NEDD1 - polyarteritis nodosa L2G 0.504; unmet 4.85
79. 79. GCFC2 - polyarteritis nodosa L2G 0.501; unmet 4.85
80. 80. SEMA3C - polyarteritis nodosa L2G 0.48; unmet 4.85
81. 81. C7orf33 - polyarteritis nodosa L2G 0.392; unmet 4.85
82. 82. ELFN1 - polyarteritis nodosa L2G 0.222; unmet 4.85
83. 83. TYW1B - polyarteritis nodosa L2G 0.164; unmet 4.85
84. 84. BLK - esophageal adenocarcinoma L2G 0.831; unmet 4.84
85. 85. NNT - esophageal adenocarcinoma L2G 0.325; unmet 4.84
86. 86. UST - amyotrophic lateral sclerosis L2G 0.771; unmet 4.81
87. 87. DUSP16 - amyotrophic lateral sclerosis L2G 0.751; unmet 4.81
88. 88. ANKS1B - amyotrophic lateral sclerosis L2G 0.75; unmet 4.81
89. 89. ACOXL - amyotrophic lateral sclerosis L2G 0.674; unmet 4.81
90. 90. G2E3 - amyotrophic lateral sclerosis L2G 0.643; unmet 4.81
91. 91. CRYZ - amyotrophic lateral sclerosis L2G 0.608; unmet 4.81
92. 92. NFIA - amyotrophic lateral sclerosis L2G 0.52; unmet 4.81
93. 93. WDR72 - amyotrophic lateral sclerosis L2G 0.459; unmet 4.81
94. 94. NDST4 - amyotrophic lateral sclerosis L2G 0.451; unmet 4.81
95. 95. CCNA1 - amyotrophic lateral sclerosis L2G 0.367; unmet 4.81
96. 96. KIAA0232 - amyotrophic lateral sclerosis L2G 0.311; unmet 4.81
97. 97. MTNR1B - amyotrophic lateral sclerosis L2G 0.305; unmet 4.81
98. 98. GGT5 - amyotrophic lateral sclerosis L2G 0.169; unmet 4.81
99. 99. POU5F1B - hepatocellular carcinoma L2G 0.828; unmet 4.79
100. 100. HNF1B - hepatocellular carcinoma L2G 0.71; unmet 4.79

1. PLCE1 - thoracic aortic aneurysm

ENSG00000138193 | EFO_0004282

L2G score
0.851

Unmet need index
5.73

SAS global MAF
0.3109

Genome India global MAF
0.3439

Variants
2

Interpretations
Likely consequential

Disease Description

An aneurysm formed in the wall of the proximal portion of the descending aorta proceeding from the arch of the aorta.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.851 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
10_94032003_G_C	missense_variant	Likely consequential	0.001762	0.001842	0	0	21.2	0.6314		0.24	0.2131	0.3109	0.3439
10_94171330_G_T	missense_variant	Likely consequential	0.3097	0.3427	0.1691	0.1296	23.8	0.3728	0.5887	0.086	0.01078	0.3109	0.3439

2. ELN - thoracic aortic aneurysm

ENSG00000049540 | EFO_0004282

L2G score 0.801	Unmet need index 5.73	SAS global MAF 0.0003826
Genome India global MAF 0.002047	Variants 1	Interpretations Likely consequential

Disease Description

An aneurysm formed in the wall of the proximal portion of the descending aorta proceeding from the arch of the aorta.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.801 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_74042640_T_C	missense_variant	Likely consequential	0.0003826	0.002047	0.0002662	0.00732	20.7	0.834		0.125	0.1057	0.0003826	0.002047

3. ULK4 - thoracic aortic aneurysm

ENSG00000168038 | EFO_0004282

L2G score 0.774	Unmet need index 5.73	SAS global MAF 0.007181
Genome India global MAF 0	Variants 2	Interpretations not_missense

Disease Description

An aneurysm formed in the wall of the proximal portion of the descending aorta proceeding from the arch of the aorta.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.774 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
3_41819492_CT_C	frameshift_variant	not_missense	0.00245		0.0002843	0.001709	28.5					0.007181	0
3_41819492_C_CT	frameshift_variant	not_missense	0.004742		0.001492	0.003278	33					0.007181	0

4. ABO - pancreatic adenocarcinoma

ENSG00000175164 | EFO_1000044

L2G score 0.836	Unmet need index 5.35	SAS global MAF 0.04705
Genome India global MAF 0.03973	Variants 2	Interpretations not_missense

Disease Description

A carcinoma that arises from glandular epithelial cells of the pancreas

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.836 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
9_133256189_C_T	stop_gained	not_missense	0.0329	0.02446	0.01796	0.01162	51					0.04705	0.03973
9_133256234_G_T_G	frameshift_variant	not_missense	0.01463	0.01566	0.002961	0.0003027	17.65					0.04705	0.03973

5. HNF1B - squamous cell lung carcinoma

ENSG00000275410 | EFO_0000708

L2G score 0.865	Unmet need index 5.31	SAS global MAF 0.0169
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

A carcinoma arising from squamous bronchial epithelial cells. It may be keratinizing or non-keratinizing. Keratinizing squamous cell carcinoma is characterized by the presence of keratinization, pearl formation, and/or intercellular bridges. Non-keratinizing squamous cell carcinoma is characterized by the absence of keratinization, pearl formation, and intercellular bridges. Cigarette smoking and arsenic exposure are strongly associated with squamous cell lung carcinoma.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.865 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37710582_GTCACCAT_G	frameshift_variant	not_missense	0.0169		0.0023	0.02255	35					0.0169	0

6. CYP2A6 - squamous cell lung carcinoma

ENSG00000255974 | EFO_0000708

L2G score 0.238	Unmet need index 5.31	SAS global MAF 0.01779
Genome India global MAF 0.00764	Variants 3	Interpretations Likely consequential; not_missense

Disease Description

A carcinoma arising from squamous bronchial epithelial cells. It may be keratinizing or non-keratinizing. Keratinizing squamous cell carcinoma is characterized by the presence of keratinization, pearl formation, and/or intercellular bridges. Non-keratinizing squamous cell carcinoma is characterized by the absence of keratinization, pearl formation, and intercellular bridges. Cigarette smoking and arsenic exposure are strongly associated with squamous cell lung carcinoma.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.238 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
19_40843845_C_A	missense_variant	Likely consequential	0.001052		0.000346	0.001268	23.1	0.6058		0.147	0.1383	0.01779	0.00764
19_40843881_G_T	stop_gained	not_missense	0.009121		0.01919	0.003588	33					0.01779	0.00764
19_40848304_C_A	missense_variant	Likely consequential	0.007708	0.00764	0	0	29	0.7887		0.312	0.2805	0.01779	0.00764

7. ERC1 - squamous cell lung carcinoma

ENSG00000082805 | EFO_0000708

L2G score 0.187	Unmet need index 5.31	SAS global MAF 0.001693
Genome India global MAF 0	Variants 1	Interpretations Likely consequential

Disease Description

A carcinoma arising from squamous bronchial epithelial cells. It may be keratinizing or non-keratinizing. Keratinizing squamous cell carcinoma is characterized by the presence of keratinization, pearl formation, and/or intercellular bridges. Non-keratinizing squamous cell carcinoma is characterized by the absence of keratinization, pearl formation, and intercellular bridges. Cigarette smoking and arsenic exposure are strongly associated with squamous cell lung carcinoma.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.187 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_1028211_GC	missense_variant	Likely consequential	0.001693		0	0	31	0.9299		0.779	0.1305	0.001693	0

8. ABO - Ischemic stroke

ENSG00000175164 | HP_0002140

L2G score 0.961	Unmet need index 5.27	SAS global MAF 0.04705
Genome India global MAF 0.03973	Variants 2	Interpretations not_missense

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.961 across 8 evidence rows and 8 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
9_133256189_C_T	stop_gained	not_missense	0.0329	0.02446	0.01796	0.01162	51					0.04705	0.03973
9_133256234_G_T	frameshift_variant	not_missense	0.01463	0.01566	0.002961	0.0003027	17.65					0.04705	0.03973

9. HDLBP - Ischemic stroke

ENSG00000115677 | HP_0002140

L2G score 0.763	Unmet need index 5.27	SAS global MAF 0.04532
Genome India global MAF 0	Variants 1	Interpretations Likely consequential

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.763 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_241239714_A_C	missense_variant	Likely consequential	0.04532		0.006525	0.03564	24.6	0.9741		0.575	0.05503	0.04532	0

10. CUX2 - Ischemic stroke

ENSG00000111249 | HP_0002140

L2G score 0.717	Unmet need index 5.27	SAS global MAF 0.006762
Genome India global MAF 0	Variants 2	Interpretations Likely consequential; not_missense

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.717 across 9 evidence rows and 9 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_111347618_T_TGC	frameshift_variant	not_missense	0.002081		0.007879	3.78e-05	32					0.006762	0
12_111347622_A_C	missense_variant	Likely consequential	0.00469		0.02853	0.00159	21.5	0.0743		0.072	0.737	0.006762	0

11. SPPL2A - Ischemic stroke

ENSG00000138600 | HP_0002140

L2G score
0.45

Unmet need index
5.27

SAS global MAF
0.001057

Genome India global MAF
0.0008187

Variants
1

Interpretations
Likely consequential

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.45 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_50722170_A_T	missense_variant	Likely consequential	0.001057	0.0008187	0.002762	0.0001206	24.3	0.9626		0.538	0.08175	0.001057	0.0008187

12. ADH5 - Ischemic stroke

ENSG00000197894 | HP_0002140

L2G score 0.334	Unmet need index 5.27	SAS global MAF 0.0001752
Genome India global MAF 0.001177	Variants 1	Interpretations Likely consequential

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.334 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_99072585_T_G	missense_variant	Likely consequential	0.0001752	0.001177	0	0	22.6	0.7039		0.222	0.4195	0.0001752	0.001177

13. MMP3 - Ischemic stroke

ENSG00000149968 | HP_0002140

L2G score
0.277

Unmet need index
5.27

SAS global MAF
0.09613

Genome India global MAF
0

Variants
2

Interpretations
not_missense

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.277 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
11_102842429_A_T	splice_donor_variant	not_missense	0.06631		0.008394	0.07395	27.2					0.09613	0
11_102842430_C_T	splice_donor_variant	not_missense	0.03193		0.008242	0.05908	29					0.09613	0

14. ICA1L - Ischemic stroke

ENSG00000163596 | HP_0002140

L2G score 0.231	Unmet need index 5.27	SAS global MAF 1.16e-05
Genome India global MAF 0.9999	Variants 1	Interpretations not_missense

Disease Description

Acute ischemic stroke (AIS) is defined by the sudden loss of blood flow to an area of the brain with the resulting loss of neurologic function. It is caused by thrombosis or embolism that occludes a cerebral vessel supplying a specific area of the brain. During a vessel occlusion, there is a core area where damage to the brain is irreversible and an area of penumbra where the brain has lost function owing to decreased blood flow but is not irreversibly injured.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.231 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_20278834_G_GA	frameshift_variant	not_missense	1.16e-05	0.9999	0	0						1.16e-05	0.9999

15. PLCE1 - esophageal squamous cell carcinoma

ENSG00000138193 | EFO_0005922

L2G score
0.294

Unmet need index
5.25

SAS global MAF
0.3109

Genome India global MAF
0.3439

Variants
2

Interpretations
Likely consequential

Disease Description

Esophageal squamous cell carcinoma (ESCC) is a type of esophageal carcinoma (EC) that can affect any part of the esophagus, but is usually located in the upper or middle third.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.294 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
10_94032003_G_C	missense_variant	Likely consequential	0.001762	0.001842	0	0	21.2	0.6314		0.24	0.2131	0.3109	0.3439
10_94171330_G_T	missense_variant	Likely consequential	0.3097	0.3427	0.1691	0.1296	23.8	0.3728	0.5887	0.086	0.01078	0.3109	0.3439

16. WDR72 - chronic kidney disease

ENSG00000166415 | EFO_0003884

L2G score
0.963

Unmet need index
5.24

SAS global MAF
0.2429

Genome India global MAF
0.259

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.963 across 8 evidence rows and 8 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_53615751_G_A	missense_variant	Likely consequential	0.2429	0.259	0.2251	0.257	25.2	0.3382	0.7526	0.528	0.02449	0.2429	0.259

17. SHROOM3 - chronic kidney disease

ENSG00000138771 | EFO_0003884

L2G score
0.885

Unmet need index
5.24

SAS global MAF
0.003458

Genome India global MAF
0.005114

Variants
2

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.885 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_76555619_G_T	missense_variant	Likely consequential	0.001218	0.0006652	0.002349	0.0007865	27.8	0.7298		0.506	0.1086	0.003458	0.005114
4_76741205_T_C	missense_variant	Likely consequential	0.002243	0.004451	0	0	25.5	0.7435		0.493	0.3707	0.003458	0.005114

18. PDILT - chronic kidney disease

ENSG00000169340 | EFO_0003884

L2G score
0.884

Unmet need index
5.24

SAS global MAF
0.003536

Genome India global MAF
0.002712

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.884 across 2e+01 evidence rows and 2e+01 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
16_20362577_G_A	missense_variant	Likely consequential	0.003536	0.002712	0.006165	0.001292	23.5	0.3062		0.586	0.04297	0.003536	0.002712

19. TCF7L2 - chronic kidney disease

ENSG00000148737 | EFO_0003884

L2G score
0.884

Unmet need index
5.24

SAS global MAF
0.002843

Genome India global MAF
0

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.884 across 9 evidence rows and 9 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
10_113165649_T_C	missense_variant	Likely consequential	0.002843		0.001924	0.002653	26.4	0.0881	0.0369	0.521	0.03355	0.002843	0

20. BCAS3 - chronic kidney disease

ENSG00000141376 | EFO_0003884

L2G score
0.639

Unmet need index
5.24

SAS global MAF
0.000116

Genome India global MAF
0.001177

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.639 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_61040883_C_T	missense_variant	Likely consequential	0.000116	0.001177	0	0	27			0.21	0.6974	0.000116	0.001177

21. C6orf118 - chronic kidney disease

ENSG00000112539 | EFO_0003884

L2G score 0.525	Unmet need index 5.24	SAS global MAF 0.001136
Genome India global MAF 0.00133	Variants 1	Interpretations not_missense

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.525 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
6_165309577_C_A	stop_gained	not_missense	0.001136	0.00133	0	0	34					0.001136	0.00133

22. PCSK9 - chronic kidney disease

ENSG00000169174 | EFO_0003884

L2G score
0.458

Unmet need index
5.24

SAS global MAF
0.001832

Genome India global MAF
0

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.458 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_55057433_GC	missense_variant	Likely consequential	0.001832		0	0	23.5	0.3248	0.6684	0.674	0.2314	0.001832	0

23. IVNS1ABP - chronic kidney disease

ENSG00000116679 | EFO_0003884

L2G score
0.452

Unmet need index
5.24

SAS global MAF
0.00371

Genome India global MAF
0.003837

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.452 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_185298225_T_G	missense_variant	Likely consequential	0.00371	0.003837	0	0	27.3	0.6474		0.941	0.3666	0.00371	0.003837

24. CYP1A2 - chronic kidney disease

ENSG00000140505 | EFO_0003884

L2G score 0.402	Unmet need index 5.24	SAS global MAF 0.001171
Genome India global MAF 0.001023	Variants 1	Interpretations Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.402 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_74754877_A_T	missense_variant	Likely consequential	0.001171	0.001023	0	0	23.8	0.3768		0.606	0.6782	0.001171	0.001023

25. DAB2 - chronic kidney disease

ENSG00000153071 | EFO_0003884

L2G score 0.318	Unmet need index 5.24	SAS global MAF 0.002203
Genome India global MAF 0.001637	Variants 1	Interpretations Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.318 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
5_39394311_C_T	missense_variant	Likely consequential	0.002203	0.001637	0.00187	0.0002247	28.6	0.6487		0.219	0.04278	0.002203	0.001637

26. DHR511 - chronic kidney disease

ENSG00000278535 | EFO_0003884

L2G score
0.269

Unmet need index
5.24

SAS global MAF
0.001042

Genome India global MAF
0

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.269 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_36595095_T_G	missense_variant	Likely consequential	0.001042		0.003927	0.0003952	24.2	0.7491			0.9855	0.001042	0

27. OR52N2 - chronic kidney disease

ENSG00000180988 | EFO_0003884

L2G score
0.218

Unmet need index
5.24

SAS global MAF
0.001204

Genome India global MAF
0.0003582

Variants
1

Interpretations
Likely consequential

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.218 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
11_5820697_C_T	missense_variant	Likely consequential	0.001204	0.0003582	0	0	21.5	0.5659		0.403	0.4332	0.001204	0.0003582

28. GAB2 - chronic kidney disease

ENSG00000033327 | EFO_0003884

L2G score 0.197	Unmet need index 5.24	SAS global MAF 0.1638
Genome India global MAF 0	Variants 3	Interpretations Likely consequential; not_missense

Disease Description

The end-stage of CHRONIC RENAL INSUFFICIENCY. It is characterized by the severe irreversible kidney damage (as measured by the level of PROTEINURIA) and the reduction in GLOMERULAR FILTRATION RATE to less than 15 ml per min (Kidney Foundation: Kidney Disease Outcome Quality Initiative, 2002). These patients generally require HEMODIALYSIS or KIDNEY TRANSPLANTATION.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.197 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
11_78226603_T_G	missense_variant	Likely consequential	0.001572		0.0008841	0.001251	26.3	0.9743		0.251	0.3241	0.1638	0
11_78226611_C_G	missense_variant	Likely consequential	0.1604		0.1098	0.064	24.8	0.9349		0.364	0.01756	0.1638	0
11_78226616_T_TG	frameshift_variant	not_misense	0.002415		0.000905	0.001438	34					0.1638	0

29. PSEN1 - intracranial hemorrhage

ENSG00000080815 | EFO_0000551

L2G score 0.751	Unmet need index 5.2	SAS global MAF 0.001961
Genome India global MAF 0	Variants 1	Interpretations Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.751 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
14_73192774_A_C	missense_variant	Likely consequential	0.001961		0.007899	0.0009889	25.1	0.611	0.6764	0.941	0.9803	0.001961	0

30. EXOC4 - intracranial hemorrhage

ENSG00000131558 | EFO_0000551

L2G score 0.711	Unmet need index 5.2	SAS global MAF 0.007772
Genome India global MAF 0.008239	Variants 1	Interpretations Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.711 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_134064365_C_T	missense_variant	Likely consequential	0.007772	0.008239	0.001528	0.000612	29.2	0.4057	0.7702	0.367	0.03259	0.007772	0.008239

31. ANKS1B - intracranial hemorrhage

ENSG00000185046 | EFO_0000551

L2G score 0.621	Unmet need index 5.2	SAS global MAF 8.115e-05
Genome India global MAF 0.001228	Variants 1	Interpretations Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.621 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_99806627_T_C	missense_variant	Likely consequential	8.115e-05	0.001228	0	0	27.7	0.2076			0.7108	8.115e-05	0.001228

32. WWOX - intracranial hemorrhage

ENSG00000186153 | EFO_0000551

L2G score 0.611	Unmet need index 5.2	SAS global MAF 0.08356
Genome India global MAF 0.09746	Variants 2	Interpretations Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.611 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
16_78115103_C_T	missense_variant	Likely consequential	0.01693	0.01202	0.007544	0.01005	31	0.1905	0.5718		0.02518	0.08356	0.09746
16_78425018_C_G	missense_variant	Likely consequential	0.06778	0.08648	0.0002878	0.002808	24	0.1893	0.5105		0.02795	0.08356	0.09746

33. ACACA - intracranial hemorrhage

ENSG00000278540 | EFO_0000551

L2G score 0.594	Unmet need index 5.2	SAS global MAF 0.001516
Genome India global MAF 0	Variants 1	Interpretations Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.594 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37221733_T_G	missense_variant	Likely consequential	0.001516		0.005953	0.0001697	25.7				0.9725	0.001516	0

34. PSPH - intracranial hemorrhage

ENSG00000146733 | EFO_0000551

L2G score
0.504

Unmet need index
5.2

SAS global MAF
0.5026

Genome India global MAF
0.000307

Variants
6

Interpretations
**Likely consequential;
not_missense**

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.504 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_56019598_AC_A	splice_donor_variant	not_missense	0.05696		0.0123	0.07816	27.6					0.5026	0.000307
7_56019607_C_T	missense_variant	Likely consequential	0.06794	0.000307	0.01486	0.08684	24.7	0.4362	0.3032	0.738	0.1532	0.5026	0.000307
7_56019672_A_G	missense_variant	Likely consequential	0.1325		0.01804	0.09756	26.5	0.9648	0.415	0.748	0.1037	0.5026	0.000307
7_56019681_C_T	missense_variant	Likely consequential	0.1195		0.01489	0.09731	28.3	0.9682	0.7713	0.818	0.1607	0.5026	0.000307
7_56019730_G_A	missense_variant	Likely consequential	0.1481		0.02094	0.1173	22.1	0.3969	0.4362	0.595	0.03455	0.5026	0.000307

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_5602118_TC	missense_variant	Likely consequential	0.1304		0.03035	0.1151	33	0.9506	0.7497	0.692	0.06746	0.5026	0.000307

35. TMEM130 - intracranial hemorrhage

ENSG00000166448 | EFO_0000551

L2G score
0.395

Unmet need index
5.2

SAS global MAF
0.001731

Genome India global MAF
0.001995

Variants
1

Interpretations
Likely consequential

Disease Description

Bleeding within the SKULL, including hemorrhages in the brain and the three membranes of MENINGES. The escape of blood often leads to the formation of HEMATOMA in the cranial epidural, subdural, and subarachnoid spaces.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.395 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Common

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_98851615_A_G	missense_variant	Likely consequential	0.001731	0.001995	0	0	26.4	0.8845		0.305	0.09118	0.001731	0.001995

36. RTEL1 - glioblastoma multiforme

ENSG00000258366 | EFO_0000519

L2G score 0.451	Unmet need index 5.07	SAS global MAF 0.001287
Genome India global MAF 0.0008698	Variants 1	Interpretations Likely consequential

Disease Description

The most malignant astrocytic tumor (WHO grade IV). It is composed of poorly differentiated neoplastic astrocytes and it is characterized by the presence of cellular polymorphism, nuclear atypia, brisk mitotic activity, vascular thrombosis, microvascular proliferation and necrosis. It typically affects adults and is preferentially located in the cerebral hemispheres. It may develop from diffuse astrocytoma WHO grade II or anaplastic astrocytoma (secondary glioblastoma, IDH-mutant), but more frequently, it manifests after a short clinical history de novo, without evidence of a less malignant precursor lesion (primary glioblastoma, IDH- wildtype). (Adapted from WHO)

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.451 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
20_63662850_G_A	missense_variant	Likely consequential	0.001287	0.0008698	0	0	25.2	0.6091	0.3284	0.519	0.3006	0.001287	0.0008698

37. HNF1B - ovarian carcinoma

ENSG00000275410 | EFO_0001075

L2G score 0.865	Unmet need index 5.07	SAS global MAF 0.0169
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.865 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37710582_GTCACCAT_G	frameshift_variant	not_missense	0.0169		0.0023	0.02255	35					0.0169	0

38. ABO - ovarian carcinoma

ENSG00000175164 | EFO_0001075

L2G score 0.799	Unmet need index 5.07	SAS global MAF 0.04705
Genome India global MAF 0.03973	Variants 2	Interpretations not_missense

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.799 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
9_133256189_C_T	stop_gained	not_missense	0.0329	0.02446	0.01796	0.01162	51					0.04705	0.03973
9_133256234_G_T_G	frameshift_variant	not_missense	0.01463	0.01566	0.002961	0.0003027	17.65					0.04705	0.03973

39. HOXD3 - ovarian carcinoma

ENSG00000128652 | EFO_0001075

L2G score 0.739	Unmet need index 5.07	SAS global MAF 0.0005333
Genome India global MAF 0.001023	Variants 1	Interpretations Likely consequential

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.739 across 5 evidence rows and 5 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_176172148_C_G	missense_variant	Likely consequential	0.000533	0.001023	0	0	26.2	0.8374			0.4206	0.000533	0.001023

40. POU5F1B - ovarian carcinoma

ENSG00000212993 | EFO_0001075

L2G score 0.54	Unmet need index 5.07	SAS global MAF 0.001041
Genome India global MAF 0.002812	Variants 2	Interpretations Likely consequential; not_missense

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.54 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_127416359_G_C	missense_variant	Likely consequential	0.0008054	0.001586	0	0	20.9	0.6828			0.9479	0.001041	0.002812
8_127416785_G_T	stop_gained	not_misense	0.0002362	0.001228	0	0	35					0.001041	0.002812

41. ATP6V0A1 - ovarian carcinoma

ENSG00000033627 | EFO_0001075

L2G score 0.357	Unmet need index 5.07	SAS global MAF 0.00021
Genome India global MAF 0.001433	Variants 1	Interpretations Likely consequential

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.357 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_42521077_C_G	missense_variant	Likely consequential	0.00021	0.001433	0	0	29.3	0.9754		0.97	0.9982	0.00021	0.001433

42. SPPL2C - ovarian carcinoma

ENSG00000185294 | EFO_0001075

L2G score 0.234	Unmet need index 5.07	SAS global MAF 0.07968
Genome India global MAF 0.05546	Variants 1	Interpretations Likely consequential

Disease Description

A malignant neoplasm originating from the surface ovarian epithelium. It accounts for the greatest number of deaths from malignancies of the female genital tract and is the fifth leading cause of cancer fatalities in women. It is predominantly a disease of older white women of northern European extraction, but it is seen in all ages and ethnic groups. Adenocarcinomas constitute the vast majority of ovarian carcinomas. The pattern of metastatic spread in ovarian carcinoma is similar regardless of the microscopic type. The most common sites of involvement are the contralateral ovary, peritoneal cavity, para-aortic and pelvic lymph nodes, and liver. Lung and pleura are the most common sites of extra-abdominal spread. The primary form of therapy is surgical. The overall prognosis of ovarian carcinoma remains poor, a direct result of its rapid growth rate and the lack of early symptoms. --2002

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.234 across 6 evidence rows and 6 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_45846288_G_C	missense_variant	Likely consequential	0.07968	0.05546	0.2218	0.07211	23.2	0.8506		0.293	0.1048	0.07968	0.05546

43. MAST4 - bullous pemphigoid

ENSG00000069020 | EFO_0007187

L2G score 0.871	Unmet need index 5.07	SAS global MAF 0.001132
Genome India global MAF 0.0008698	Variants 1	Interpretations not_missense

Disease Description

Bullous pemphigoid (BP) is the most common form of autoimmune bullous dermatosis.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.871 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
5_67166133_GA_G	frameshift_variant	not_missense	0.001132	0.0008698	0	0	24					0.001132	0.0008698

44. PARD3 - cardiac arrest

ENSG00000148498 | EFO_0009492

L2G score 0.877	Unmet need index 5.03	SAS global MAF 0.0003711
Genome India global MAF 0.001586	Variants 1	Interpretations Likely consequential

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.877 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
10_34341687_C_T	missense_variant	Likely consequential	0.0003711	0.001586	0	0	24.4			0.124	0.7087	0.0003711	0.001586

45. IGSF5 - cardiac arrest

ENSG00000183067 | EFO_0009492

L2G score 0.758	Unmet need index 5.03	SAS global MAF 0.0007783
Genome India global MAF 0.001279	Variants 1	Interpretations Likely consequential

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.758 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
21_39765840_C_T	missense_variant	Likely consequential	0.0007783	0.001279	0	0	23.9	0.7657		0.362	0.5046	0.0007783	0.001279

46. THSD7B - cardiac arrest

ENSG00000144229 | EFO_0009492

L2G score 0.752	Unmet need index 5.03	SAS global MAF 0.04541
Genome India global MAF 0.05993	Variants 1	Interpretations not_missense

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.752 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_137272664_C_T	splice_donor_variant	not_missense	0.04541	0.05993	0.007994	0.009018	20.5					0.04541	0.05993

47. PREX2 - cardiac arrest

ENSG000000046889 | EFO_0009492

L2G score
0.726

Unmet need index
5.03

SAS global MAF
0.006294

Genome India global MAF
0.0003582

Variants
2

Interpretations
Likely consequential

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.726 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_68044581_G_A	missense_variant	Likely consequential	0.001104	0.0003582	0.002941	0.0001686	27.1	0.6788	0.5794	0.636	0.05465	0.006294	0.0003582
8_68192488_G_C	missense_variant	Likely consequential	0.005195		0.0009588	0.01402	23.3	0.9207	0.4299	0.635	0.04875	0.006294	0.0003582

48. FMN2 - cardiac arrest

ENSG00000155816 | EFO_0009492

L2G score 0.69	Unmet need index 5.03	SAS global MAF 0.03854
Genome India global MAF 0	Variants 3	Interpretations not_missense

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.69 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_240092269_GC_G	frameshift_variant	not_missense	0.01521		0.003124	0.03017	27.3					0.03854	0
1_240207664_CA	frameshift_variant	not_missense	0.01078		0.009574	0.04247	21.5					0.03854	0
1_240207667_CA_C	frameshift_variant	not_missense	0.01306		0.01156	0.05875	14.43					0.03854	0

49. ADCY5 - cardiac arrest

ENSG00000173175 | EFO_0009492

L2G score 0.641	Unmet need index 5.03	SAS global MAF 0.00313
Genome India global MAF 0.0007163	Variants 2	Interpretations Likely consequential

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.641 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
3_123300202_C_T	missense_variant	Likely consequential	0.001982	0.0007163	0.000893	0.002507	25.4	0.736	0.6324	0.45	0.02516	0.00313	0.0007163
3_123447564_C_A	missense_variant	Likely consequential	0.00115		0	0	29.6	0.9746	0.9394	0.377	0.7995	0.00313	0.0007163

50. SPDYE21 - cardiac arrest

ENSG00000230358 | EFO_0009492

L2G score 0.539	Unmet need index 5.03	SAS global MAF 0.1466
Genome India global MAF 0.2538	Variants 2	Interpretations not_missense

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.539 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_67278874_G_A	splice_donor_variant	not_misense		0.08781			17.61					0.1466	0.2538
7_67281505_T_TGGGTGGC	frameshift_variant	not_misense	0.1466	0.182	0.1583	0.2605	22.2					0.1466	0.2538

51. OR10AD1 - cardiac arrest

ENSG00000172640 | EFO_0009492

L2G score 0.226	Unmet need index 5.03	SAS global MAF 0.002723
Genome India global MAF 0.002507	Variants 1	Interpretations not_missense

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.226 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_48202807_A_C	stop_gained	not_missense	0.002723	0.002507	0.001946	0.0007715	33					0.002723	0.002507

52. LMTK3 - cardiac arrest

ENSG00000142235 | EFO_0009492

L2G score
0.214

Unmet need index
5.03

SAS global MAF
0.02655

Genome India global MAF
0

Variants
4

Interpretations
**Likely consequential;
not_missense**

Disease Description

Cessation of breathing and/or cardiac function.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.214 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Ultra-Orphan

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
19_48491500_C_CG	frameshift_variant	not_missense	0.001048		0.0001579	0.001178	35					0.02655	0
19_48498801_C_GG_C	frameshift_variant	not_missense	0.01424		0.004306	0.001102	33					0.02655	0
19_48498801_C_CG	frameshift_variant	not_missense	0.006912		0.01321	0.005314	29.4					0.02655	0
19_48509445_A_C	missense_variant	Likely consequential	0.004569		0.001078	0.008125	26.2	0.996			0.03098	0.02655	0

53. TP63 - lung adenocarcinoma

ENSG00000073282 | EFO_0000571

L2G score
0.906

Unmet need index
5

SAS global MAF
0.001565

Genome India global MAF
0.001433

Variants
1

Interpretations
Likely consequential

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.906 across 1e+01 evidence rows and 1e+01 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
3_189889363_C_A	missense_variant	Likely consequential	0.001565	0.001433	0.003678	0.0008985	24.6	0.0744	0.2006	0.553	0.02005	0.001565	0.001433

54. SECISBP2L - lung adenocarcinoma

ENSG00000138593 | EFO_0000571

L2G score 0.884	Unmet need index 5	SAS global MAF 0.0002435
Genome India global MAF 0.001279	Variants 1	Interpretations Likely consequential

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.884 across 6 evidence rows and 6 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_48992719_T_C	missense_variant	Likely consequential	0.0002435	0.001279	0	0	22.7	0.1125			0.597	0.0002435	0.001279

55. HNF1B - lung adenocarcinoma

ENSG00000275410 | EFO_0000571

L2G score 0.865	Unmet need index 5	SAS global MAF 0.0169
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.865 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37710582_GTCACCAT_G	frameshift_variant	not_missense	0.0169		0.0023	0.02255	35					0.0169	0

56. FOXP4 - lung adenocarcinoma

ENSG00000137166 | EFO_0000571

L2G score 0.575	Unmet need index 5	SAS global MAF 0.001021
Genome India global MAF 0.001279	Variants 1	Interpretations Likely consequential

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.575 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
6_41597932_TC	missense_variant	Likely consequential	0.001021	0.001279	0	0	25.8	0.1071		0.608	0.1747	0.001021	0.001279

57. TCF7L2 - lung adenocarcinoma

ENSG00000148737 | EFO_0000571

L2G score
0.435

Unmet need index
5

SAS global MAF
0.002843

Genome India global MAF
0

Variants
1

Interpretations
Likely consequential

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.435 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
10_113165649_T_C	missense_variant	Likely consequential	0.002843		0.001924	0.002653	26.4	0.0881	0.0369	0.521	0.03355	0.002843	0

58. RTEL1 - lung adenocarcinoma

ENSG00000258366 | EFO_0000571

L2G score 0.277	Unmet need index 5	SAS global MAF 0.001287
Genome India global MAF 0.0008698	Variants 1	Interpretations Likely consequential

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.277 across 4 evidence rows and 4 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
20_63662850_G_A	missense_variant	Likely consequential	0.001287	0.0008698	0	0	25.2	0.6091	0.3284	0.519	0.3006	0.001287	0.0008698

59. CYP2A6 - lung adenocarcinoma

ENSG00000255974 | EFO_0000571

L2G score 0.238	Unmet need index 5	SAS global MAF 0.01779
Genome India global MAF 0.00764	Variants 3	Interpretations Likely consequential; not_missense

Disease Description

A carcinoma that arises from the lung and is characterized by the presence of malignant glandular epithelial cells. There is a male predilection with a male to female ratio of 2:1. Usually lung adenocarcinoma is asymptomatic and is identified through screening studies or as an incidental radiologic finding. If clinical symptoms are present they include shortness of breath, cough, hemoptysis, chest pain, and fever. Tobacco smoke is a known risk factor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.238 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
19_40843845_C_A	missense_variant	Likely consequential	0.001052		0.000346	0.001268	23.1	0.6058		0.147	0.1383	0.01779	0.00764
19_40843881_G_T	stop_gained	not_missense	0.009121		0.01919	0.003588	33					0.01779	0.00764
19_40848304_C_A	missense_variant	Likely consequential	0.007708	0.00764	0	0	29	0.7887		0.312	0.2805	0.01779	0.00764

60. GPC5 - gastric cancer

ENSG00000179399 | MONDO_0001056

L2G score 0.869	Unmet need index 5	SAS global MAF 0.004093
Genome India global MAF 0.002456	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm involving the stomach.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.869 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
13_91693637_C_G	missense_variant	Likely consequential	0.004093	0.002456	0	0	23.8	0.489		0.838	0.2864	0.004093	0.002456

61. POU5F1B - gastric cancer

ENSG00000212993 | MONDO_0001056

L2G score 0.828	Unmet need index 5	SAS global MAF 0.001041
Genome India global MAF 0.002812	Variants 2	Interpretations Likely consequential; not_missense

Disease Description

A primary or metastatic malignant neoplasm involving the stomach.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.828 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_127416359_G_C	missense_variant	Likely consequential	0.0008054	0.001586	0	0	20.9	0.6828			0.9479	0.001041	0.002812
8_127416785_G_T	stop_gained	not_missense	0.0002362	0.001228	0	0	35					0.001041	0.002812

62. HNF1B - gastric cancer

ENSG00000275410 | MONDO_0001056

L2G score 0.71	Unmet need index 5	SAS global MAF 0.0169
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

A primary or metastatic malignant neoplasm involving the stomach.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.71 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37710582_GTCACCAT_G	frameshift_variant	not_missense	0.0169		0.0023	0.02255	35					0.0169	0

63. STPG2 - gastric cancer

ENSG00000163116 | MONDO_0001056

L2G score 0.342	Unmet need index 5	SAS global MAF 0.4415
Genome India global MAF 0.4197	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm involving the stomach.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.342 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_98128442_A_G	missense_variant	Likely consequential	0.4415	0.4197	0.4088	0.2902	24.6	0.6438		0.185	0.03216	0.4415	0.4197

64. NR2E3 - gastric cancer

ENSG00000278570 | MONDO_0001056

L2G score 0.252	Unmet need index 5	SAS global MAF 0.0007573
Genome India global MAF 0.003121	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm involving the stomach.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.252 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_71811966_G_A	missense_variant	Likely consequential	0.0007573	0.003121	0.0004058	0.006587	32	0.9546	0.6169		0.2446	0.0007573	0.003121

65. L1TD1 - neuroblastoma

ENSG00000240563 | EFO_0000621

L2G score 0.931	Unmet need index 4.96	SAS global MAF 0.004045
Genome India global MAF 0.004042	Variants 1	Interpretations not_missense

Disease Description

Neuroblastoma (NB) is the most common solid, extracranial childhood tumor. It is an aggressive pediatric cancer that originates from neural crest tissues of the sympathetic nervous system.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.931 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_62210575_GA ACT_G	frameshift_variant	not_missense	0.004045	0.004042	0.01217	0.00397	23.8					0.004045	0.004042

66. KANSL1L - neuroblastoma

ENSG00000144445 | EFO_0000621

L2G score 0.692	Unmet need index 4.96	SAS global MAF 0.003556
Genome India global MAF 0.004346	Variants 2	Interpretations Likely consequential

Disease Description

Neuroblastoma (NB) is the most common solid, extracranial childhood tumor. It is an aggressive pediatric cancer that originates from neural crest tissues of the sympathetic nervous system.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.692 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_210028899_A_G	missense_variant	Likely consequential	0.001212	0.001177	0.001075	0.0002624	25.1	0.8966		0.437	0.04716	0.003556	0.004346
2_210043964_G_C	missense_variant	Likely consequential	0.002347	0.003173	0	0	22.1	0.8495		0.196	0.2891	0.003556	0.004346

67. WRN - pneumococcal pneumonia

ENSG00000165392 | EFO_1001474

L2G score 0.585	Unmet need index 4.94	SAS global MAF 0.01562
Genome India global MAF 0.01724	Variants 1	Interpretations not_missense

Disease Description

A febrile disease caused by STREPTOCOCCUS PNEUMONIAE.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.585 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_31173019_C_T	stop_gained	not_missense	0.01562	0.01724	0.001536	0.001143	35					0.01562	0.01724

68. ASB5 - pneumococcal pneumonia

ENSG00000164122 | EFO_1001474

L2G score 0.153	Unmet need index 4.94	SAS global MAF 0.0001748
Genome India global MAF 0.001023	Variants 1	Interpretations Likely consequential

Disease Description

A febrile disease caused by STREPTOCOCCUS PNEUMONIAE.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.153 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Uncommon

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_176215715_G_A	missense_variant	Likely consequential	0.0001748	0.001023	0	0	25	0.3069		0.378	0.9872	0.0001748	0.001023

69. ACKR1 - microscopic polyangiitis

ENSG00000213088 | EFO_1000784

L2G score 0.889	Unmet need index 4.91	SAS global MAF 0.0004407
Genome India global MAF 0.0044	Variants 1	Interpretations Likely consequential

Disease Description

Microscopic polyangiitis (MPA) is an inflammatory, necrotizing, systemic vasculitis that affects predominantly small vessels (i.e. small arteries, arterioles, capillaries, venules) in multiple organs.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.889 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_159205834_G_A	missense_variant	Likely consequential	0.0004407	0.0044	0	0	21.3	0.8225	0.579	0.143	0.0569	0.0004407	0.0044

70. ADH1B - oropharynx cancer

ENSG00000196616 | EFO_1001931

L2G score 0.965	Unmet need index 4.86	SAS global MAF 0.00622
Genome India global MAF 0.003426	Variants 2	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.965 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_99307859_C_T	missense_variant	Likely consequential	0.004835	0.002456	0.003268	0.0005805	24.1	0.7715		0.218	0.1022	0.00622	0.003426
4_99307860_G_A	missense_variant	Likely consequential	0.001391	0.0009722	0.001129	5.617e-05	27.2	0.7617		0.194	0.1033	0.00622	0.003426

71. INPP5D - oropharynx cancer

ENSG00000168918 | EFO_1001931

L2G score 0.745	Unmet need index 4.86	SAS global MAF 0.002724
Genome India global MAF 0.001689	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.745 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_233158415_C_T	missense_variant	Likely consequential	0.002724	0.001689	0	0	25.1	0.8393			0.312	0.002724	0.001689

72. ANKS1A - oropharynx cancer

ENSG00000064999 | EFO_1001931

L2G score 0.612	Unmet need index 4.86	SAS global MAF 0.002968
Genome India global MAF 0.001075	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.612 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
6_35078619_G_A	missense_variant	Likely consequential	0.002968	0.001075	0.001008	0.0001319	29.4	0.1623		0.785	0.1089	0.002968	0.001075

73. CAPN13 - oropharynx cancer

ENSG00000162949 | EFO_1001931

L2G score 0.502	Unmet need index 4.86	SAS global MAF 0.003337
Genome India global MAF 0.008084	Variants 1	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.502 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_30763111_G_A	missense_variant	Likely consequential	0.003337	0.008084	0.005711	0.01445	26.2	0.5726		0.62	0.04429	0.003337	0.008084

74. SULT1C4 - oropharynx cancer

ENSG00000198075 | EFO_1001931

L2G score 0.353	Unmet need index 4.86	SAS global MAF 0.001521
Genome India global MAF 0.0009722	Variants 1	Interpretations not_missense

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.353 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_108383512_T_C	splice_donor_variant	not_missense	0.001521	0.0009722	0	0	24.7					0.001521	0.0009722

75. GPR27 - oropharynx cancer

ENSG00000170837 | EFO_1001931

L2G score 0.197	Unmet need index 4.86	SAS global MAF 0.003987
Genome India global MAF 0	Variants 2	Interpretations Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.197 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
3_71755049_A_C	missense_variant	Likely consequential	0.001837		0.006974	0.0002525	26	0.7877		0.876	0.9995	0.003987	0
3_71755050_A_C	missense_variant	Likely consequential	0.002153		0.007921	0.0002732	26.5	0.7119		0.888	0.9984	0.003987	0

76. RTL1 - oropharynx cancer

ENSG00000254656 | EFO_1001931

L2G score
0.168

Unmet need index
4.86

SAS global MAF
0.0008117

Genome India global MAF
0.001893

Variants
1

Interpretations
Likely consequential

Disease Description

A primary or metastatic malignant neoplasm that affects the oropharynx.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.168 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
14_100884608_C_T	missense_variant	Likely consequential	0.000817	0.001893	0	0	23.1	0.092			0.9187	0.000817	0.001893

77. DPY19L4 - polyarteritis nodosa

ENSG00000156162 | MONDO_0019170

L2G score 0.73	Unmet need index 4.85	SAS global MAF 0.01152
Genome India global MAF 0.002149	Variants 4	Interpretations Likely consequential; not_missense

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.73 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_94734629_G_A	splice_acceptor_variant	not_misense	0.001467	0.002149	0	0	32					0.01152	0.002149
8_94781082_A_T	splice_acceptor_variant	not_misense	0.002889		0.000628	0.003751	35					0.01152	0.002149
8_94781083_G_T	splice_acceptor_variant	not_misense	0.004793		0.01267	0.003047	35					0.01152	0.002149
8_94789792_G_A	missense_variant	Likely consequential	0.00242		4.232e-05	0.002081	26.4	0.5924		0.253	0.2076	0.01152	0.002149

78. NEDD1 - polyarteritis nodosa

ENSG00000139350 | MONDO_0019170

L2G score 0.504	Unmet need index 4.85	SAS global MAF 2.372e-05
Genome India global MAF 0.001228	Variants 1	Interpretations Likely consequential

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.504 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_96937212_A_C	missense_variant	Likely consequential	2.372e-05	0.001228	0	0	21	0.3107		0.146	0.7277	2.372e-05	0.001228

79. GCFC2 - polyarteritis nodosa

ENSG00000005436 | MONDO_0019170

L2G score 0.501	Unmet need index 4.85	SAS global MAF 0.000535
Genome India global MAF 0.001126	Variants 1	Interpretations not_missense

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.501 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_75692013_AT_A	frameshift_variant	not_missense	0.000535	0.001126	0	0	24.4					0.000535	0.001126

80. SEMA3C - polyarteritis nodosa

ENSG00000075223 | MONDO_0019170

L2G score
0.48

Unmet need index
4.85

SAS global MAF
0.03783

Genome India global MAF
0.05051

Variants
1

Interpretations
Likely consequential

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.48 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_80798214_C_T	missense_variant	Likely consequential	0.03783	0.05051	0.006466	0.001262	32	0.7772		0.353	0.08897	0.03783	0.05051

81. C7orf33 - polyarteritis nodosa

ENSG00000170279 | MONDO_0019170

L2G score 0.392	Unmet need index 4.85	SAS global MAF 0.05077
Genome India global MAF 0.04226	Variants 1	Interpretations not_missense

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.392 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_148590928_G_T	start_loss	not_missense	0.05077	0.04226	0.03583	0.01586	7.765				0.01417	0.05077	0.04226

82. ELFN1 - polyarteritis nodosa

ENSG00000225968 | MONDO_0019170

L2G score 0.222	Unmet need index 4.85	SAS global MAF 0.05891
Genome India global MAF 0	Variants 4	Interpretations Likely consequential; not_missense

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.222 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_1746231_C_CTTTTTAATGATA	frameshift_variant	not_misense	0.006063		0.001199	0.000262	23.1					0.05891	0
7_1746232_C_CG	frameshift_variant	not_misense	0.01307		0.00324	0.001124	24.3					0.05891	0
7_1746242_AGAGTTCGGTGG_A	frameshift_variant	not_misense	0.01534		0.004613	0.00178	34					0.05891	0
7_1746263_C_A	missense_variant	Likely consequential	0.02569		0.006784	0.003496	24.4	0.7519				0.05891	0

83. TYW1B - polyarteritis nodosa

ENSG00000277149 | MONDO_0019170

L2G score
0.164

Unmet need index
4.85

SAS global MAF
0.002548

Genome India global MAF
0

Variants
1

Interpretations
not_missense

Disease Description

Polyarteritis nodosa (PAN) is a rare, clinically heterogeneous, rheumatologic disease characterized by necrotizing inflammatory lesions affecting small- and medium-sized blood vessels. PAN most commonly affects skin, joints, peripheral nerves, the gut, and the kidney.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.164 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
7_72575498_T_C	stop_loss	not_missense	0.002548		0	0	17.83					0.002548	0

84. BLK - esophageal adenocarcinoma

ENSG00000136573 | EFO_0000478

L2G score
0.831

Unmet need index
4.84

SAS global MAF
0.005038

Genome India global MAF
0.002659

Variants
2

Interpretations
Likely consequential

Disease Description

A malignant tumor with glandular differentiation arising predominantly from Barrett mucosa in the lower third of the esophagus. Rare examples of esophageal adenocarcinoma deriving from ectopic gastric mucosa in the upper esophagus have also been reported. Grossly, esophageal adenocarcinomas are similar to esophageal squamous cell carcinomas. Microscopically, adenocarcinomas arising in the setting of Barrett esophagus are typically papillary and/or tubular. The prognosis is poor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.831 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_11557983_A_C	missense_variant	Likely consequential	0.002864	0.0008187	0.002022	9.361e-05	26.1	0.5985		0.829	0.06809	0.005038	0.002659
8_11561347_C_T	missense_variant	Likely consequential	0.002181	0.001842	0.001095	0.0005805	33	0.8702		0.823	0.1829	0.005038	0.002659

85. NNT - esophageal adenocarcinoma

ENSG00000112992 | EFO_0000478

L2G score 0.325	Unmet need index 4.84	SAS global MAF 0.02007
Genome India global MAF 0.01775	Variants 1	Interpretations Likely consequential

Disease Description

A malignant tumor with glandular differentiation arising predominantly from Barrett mucosa in the lower third of the esophagus. Rare examples of esophageal adenocarcinoma deriving from ectopic gastric mucosa in the upper esophagus have also been reported. Grossly, esophageal adenocarcinomas are similar to esophageal squamous cell carcinomas. Microscopically, adenocarcinomas arising in the setting of Barrett esophagus are typically papillary and/or tubular. The prognosis is poor.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.325 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
5_43700219_A_G	missense_variant	Likely consequential	0.02007	0.01775	0.004595	0.01125	23.5	0.1974		0.691	0.02754	0.02007	0.01775

86. UST - amyotrophic lateral sclerosis

ENSG00000111962 | MONDO_0004976

L2G score 0.771	Unmet need index 4.81	SAS global MAF 2.319e-05
Genome India global MAF 0.9999	Variants 1	Interpretations not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.771 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
6_149074005_GC_G	frameshift_variant	not_missense	2.319e-05	0.9999	0	0						2.319e-05	0.9999

87. DUSP16 - amyotrophic lateral sclerosis

ENSG00000111266 | MONDO_0004976

L2G score
0.751

Unmet need index
4.81

SAS global MAF
0.2318

Genome India global MAF
0.1593

Variants
2

Interpretations
not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.751 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_12477741_GCACGC_G	frameshift_variant	not_missense	0.1236	0.1593	0.00336	0.02682	23.4					0.2318	0.1593
12_12477747_TGGGC_T	frameshift_variant	not_missense	0.1235		0.003357	0.02677	26.9					0.2318	0.1593

88. ANKS1B - amyotrophic lateral sclerosis

ENSG00000185046 | MONDO_0004976

L2G score 0.75	Unmet need index 4.81	SAS global MAF 8.115e-05
Genome India global MAF 0.001228	Variants 1	Interpretations Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.75 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
12_99806627_T_C	missense_variant	Likely consequential	8.115e-05	0.001228	0	0	27.7	0.2076			0.7108	8.115e-05	0.001228

89. ACOXL - amyotrophic lateral sclerosis

ENSG00000153093 | MONDO_0004976

L2G score
0.674

Unmet need index
4.81

SAS global MAF
0.0007883

Genome India global MAF
0.001842

Variants
1

Interpretations
Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.674 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
2_110794129_C_G	missense_variant	Likely consequential	0.0007883	0.001842	0	0	6.057	0.8466		0.601	0.8359	0.0007883	0.001842

90. G2E3 - amyotrophic lateral sclerosis

ENSG00000092140 | MONDO_0004976

L2G score 0.643	Unmet need index 4.81	SAS global MAF 0.02063
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.643 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
14_30605565_T_TA	frameshift_variant	not_missense	0.02063		0.02155	0.008025	27.3					0.02063	0

91. CRYZ - amyotrophic lateral sclerosis

ENSG00000116791 | MONDO_0004976

L2G score 0.608	Unmet need index 4.81	SAS global MAF 0.001601
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.608 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_74706961_GGTTT_G	frameshift_variant	not_missense	0.001601		0	0	29.7					0.001601	0

92. NFIA - amyotrophic lateral sclerosis

ENSG00000162599 | MONDO_0004976

L2G score 0.52	Unmet need index 4.81	SAS global MAF 0.1045
Genome India global MAF 0	Variants 3	Interpretations not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.52 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
1_61406559_CAG_C	splice_acceptor_variant	not_missense	0.007711		0.001255	0.004074	34					0.1045	0
1_61406560_A_C	splice_acceptor_variant	not_missense	0.06294		0.01306	0.02819	35					0.1045	0
1_61406561_G_C	splice_acceptor_variant	not_missense	0.03689		0.009639	0.01882	36					0.1045	0

93. WDR72 - amyotrophic lateral sclerosis

ENSG00000166415 | MONDO_0004976

L2G score 0.459	Unmet need index 4.81	SAS global MAF 0.2429
Genome India global MAF 0.259	Variants 1	Interpretations Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.459 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
15_53615751_G_A	missense_variant	Likely consequential	0.2429	0.259	0.2251	0.257	25.2	0.3382	0.7526	0.528	0.02449	0.2429	0.259

94. NDST4 - amyotrophic lateral sclerosis

ENSG00000138653 | MONDO_0004976

L2G score 0.451	Unmet need index 4.81	SAS global MAF 0.001049
Genome India global MAF 0.003223	Variants 1	Interpretations Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.451 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_114829860_A_G	missense_variant	Likely consequential	0.001049	0.003223	0	0	25.2	0.4293		0.555	0.169	0.001049	0.003223

95. CCNA1 - amyotrophic lateral sclerosis

ENSG00000133101 | MONDO_0004976

L2G score 0.367	Unmet need index 4.81	SAS global MAF 0.003142
Genome India global MAF 0	Variants 2	Interpretations Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.367 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
13_36438732_A_G	missense_variant	Likely consequential	0.002022		0.001043	0.01569	30			0.682	0.1137	0.003142	0
13_36438735_T_G	missense_variant	Likely consequential	0.001122		0.000366	0.002806	27.9			0.739	0.08409	0.003142	0

96. KIAA0232 - amyotrophic lateral sclerosis

ENSG00000170871 | MONDO_0004976

L2G score
0.311

Unmet need index
4.81

SAS global MAF
0.001461

Genome India global MAF
0.002149

Variants
1

Interpretations
Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.311 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
4_6863249_G_A	missense_variant	Likely consequential	0.001461	0.002149	0	0	21.2	0.2358		0.527	0.105	0.001461	0.002149

97. MTNR1B - amyotrophic lateral sclerosis

ENSG00000134640 | MONDO_0004976

L2G score
0.305

Unmet need index
4.81

SAS global MAF
0.001924

Genome India global MAF
0.001842

Variants
1

Interpretations
Likely consequential

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.305 across 1 evidence rows and 1 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pat_hogenicity	EVE_S_CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
11_92981635_C_T	missense_variant	Likely consequential	0.001924	0.001842	0.001504	5.616e-05	18.72	0.7634		0.709	0.1385	0.001924	0.001842

98. GGT5 - amyotrophic lateral sclerosis

ENSG00000099998 | MONDO_0004976

L2G score 0.169	Unmet need index 4.81	SAS global MAF 0.05402
Genome India global MAF 0	Variants 2	Interpretations Likely consequential; not_missense

Disease Description

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by progressive muscular paralysis reflecting degeneration of motor neurons in the primary motor cortex, corticospinal tracts, brainstem and spinal cord.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.169 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
22_24232203_A_G	missense_variant	Likely consequential	0.003823		0.0009495	0.001735	27.7	0.6413		0.403	0.381	0.05402	0
22_24232210_T_G	splice_acceptor_variant	not_misense	0.05039		0.02001	0.01907	29.6					0.05402	0

99. POU5F1B - hepatocellular carcinoma

ENSG00000212993 | EFO_0000182

L2G score 0.828	Unmet need index 4.79	SAS global MAF 0.001041
Genome India global MAF 0.002812	Variants 2	Interpretations Likely consequential; not_missense

Disease Description

A malignant tumor that arises from hepatocytes. Hepatocellular carcinoma is relatively rare in the United States but very common in all African countries south of the Sahara and in Southeast Asia. Most cases are seen in patients over the age of 50 years, but this tumor can also occur in younger individuals and even in children. Hepatocellular carcinoma is more common in males than females and is associated with hepatitis B, hepatitis C, chronic alcohol abuse and cirrhosis. Serum elevation of alpha-fetoprotein occurs in a large percentage of patients with hepatocellular carcinoma. Grossly, hepatocellular carcinoma may present as a single mass, as multiple nodules, or as diffuse liver involvement. Microscopically, there is a wide range of differentiation from tumor to tumor (well differentiated to poorly differentiated tumors). Hepatocellular carcinomas quickly metastasize to regional lymph nodes and lung. The overall median survival of untreated liver cell carcinoma is about 4 months. The most effective treatment of hepatocellular carcinoma is complete resection of the tumor. Lately, an increasing number of tumors have been treated with liver transplantation.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.828 across 3 evidence rows and 3 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_S CORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
8_127416359_G_C	missense_variant	Likely consequential	0.0008054	0.001586	0	0	20.9	0.6828			0.9479	0.001041	0.002812
8_127416785_G_T	stop_gained	not_missense	0.0002362	0.001228	0	0	35					0.001041	0.002812

100. HNF1B - hepatocellular carcinoma

ENSG00000275410 | EFO_0000182

L2G score 0.71	Unmet need index 4.79	SAS global MAF 0.0169
Genome India global MAF 0	Variants 1	Interpretations not_missense

Disease Description

A malignant tumor that arises from hepatocytes. Hepatocellular carcinoma is relatively rare in the United States but very common in all African countries south of the Sahara and in Southeast Asia. Most cases are seen in patients over the age of 50 years, but this tumor can also occur in younger individuals and even in children. Hepatocellular carcinoma is more common in males than females and is associated with hepatitis B, hepatitis C, chronic alcohol abuse and cirrhosis. Serum elevation of alpha-fetoprotein occurs in a large percentage of patients with hepatocellular carcinoma. Grossly, hepatocellular carcinoma may present as a single mass, as multiple nodules, or as diffuse liver involvement. Microscopically, there is a wide range of differentiation from tumor to tumor (well differentiated to poorly differentiated tumors). Hepatocellular carcinomas quickly metastasize to regional lymph nodes and lung. The overall median survival of untreated liver cell carcinoma is about 4 months. The most effective treatment of hepatocellular carcinoma is complete resection of the tumor. Lately, an increasing number of tumors have been treated with liver transplantation.

Gene Summary

Not available in the current local cache. Add Open Targets/UniProt target metadata to populate this section.

Known Drugs / Clinical Investigation

Not available in the current local cache. Add Open Targets known-drug rows to populate this section.

Biological Evidence

Open Targets GWAS credible-set evidence supports this gene-disease pair with L2G score 0.71 across 2 evidence rows and 2 study loci.

SAS / Indian Prevalence

Rare

Alleles and Scores

varid	consequence	missense_interpretation	AF_sas	genome_india_af	AF_nfe	AF_fin	cadd_phred	am_pathogenicity	EVE_SCORE	REVEL	ClinPred	global_maf_af_sas	global_maf_genome_india_af
17_37710582_GTCACCAT_G	frameshift_variant	not_missense	0.0169		0.0023	0.02255	35					0.0169	0