

A store of hand-picked math books and of everything else Stop and shop. http://astore.amazon.com

Ask a question -Leave a comment

Terms of use Awards Interactive Activities What's new

CTK Exchange CTK Wiki Math CTK Insights - a blog Math Help Games & Puzzles What Is What Arithmetic Algebra Geometry Probability **Outline Mathematics** Make an Identity **Book Reviews** Stories for Young Eve Opener **Analog Gadgets** Inventor's Paradox Did you know?... Proofs Math as Language Things Impossible Visual Illusions My Logo Math Poll Cut The Knot! MSET99 Talk Old and nice bookstore Other Math sites Front Page

Movie shortcuts
Personal info
Privacy Policy

Guest book

Recommend this site

Sites for parents

Press collection

Education & Parenting



Pigeonhole Principle and Extensions

The *Pigeonhole Principle* is one of almost obvious mathematical concepts which are both simple and *powerful*:

(1) If n > m pigeons are put into m pigeonholes, there's a hole with more than one pigeon.

For the proof, assume that the statment is wrong: i.e., assume there are m holes each with at most 1 pigeon. If that's really the case, then summing up the birds across the holes, we would have at most 1 + ... + 1 = m pigeons. However, the given number of pigeons n > m. A contradiction that proves the statement.

The proof, as the principle itself, is very simple and embodies an idea that can be used to prove a generalized statement:

(2) If nk + 1 pigeons, where k is a positive integer, have been put into n holes, then at least one of the holes is crowded with at least k+1 pigeons.

Indeed, let's again assume that the statement is wrong. Then each of the holes houses not more than k birds, which means that the total number of birds can't exceed nk. A contradiction.

In the same vein we can establish the following extension:

If $p_1 + p_2 + ... + p_n - n + 1$ pigeons are placed into n holes, then, for some k, hole k has more than p_k pigeons.

Once more, assume that the statement is wrong, i.e., assume that, for k = 1, ..., n, hole k contains at most p_k - 1 birds. Summing up over all n holes, we find that the total number of the pigeons can't

exceed

$$\Sigma(p_k - 1) = \Sigma p_k - n.$$

A contradiction.

Finally, (2) admits a reformulation:

(2') If m pigeons are found in n holes, then at least one of the holes contains at least p = [(m-1)/n] + 1 pigeons,

where [x] is the *floor function*. Indeed, assuming that every hole contains at most p pigeons, we arrive at the contradiction:

$$m \le np$$

 $\le n \cdot (m - 1)/n$
 $= m - 1.$
 $< m.$

A straightforward reformulation of (2') has been given by E. W. Dijkstra

٠

Search:

Keywords:

Google

○ Web **⊙** CTK

Search

For a non-empty, finite bag of numbers, the maximum value is at least the average value.

(To which we can add the obvious: the average and the maximum values coincide iff the bag only contains equal numbers. Also, the above is obviously equivalent to the assertion that, for a non-empty, finite bag of numbers, the minimum value is at most the average value.)

Example

[Sharygin, p. 12]. Assume in a class of students each of the number of committees contains more than half of all the students. Prove that there is a student who is a member in more than half of the committees.

Let's n be the number of students and m the number of committees in the class. The total committee membership T exceeds $m \cdot n/2$: T > nm/2. On average, a student is a member in T/n

committees and we find that T/n > m/2. Since the maximum value is at least the average value, there is indeed a student who is a member in more than m/2 committees.



Reference

- 1. V. K. Balakrishnan, *Theory and Problems of Combinatorics*, Schaum's Outline Series, McGraw-Hill, 1995
- 2. A. Engel, Problem-Solving Strategies, Springer Verlag, 1998
- 3. D. Fomin, S. Genkin, I. Itenberg, Mathematical Circles (Russian Experience), AMS, 1996
- 4. R. Graham, D. Knuth, O. Patashnik, *Concrete Mathematics*, 2nd edition, Addison-Wesley, 1994.
- 5. R. Honsberger, *Ingenuity in Mathematics*, MAA, New Math Library, 1970
- 6. R. Honsberger, Mathematical Morsels, MAA, New Math Library, 1978
- 7. R. Honsberger, More Mathematical Morsels, MAA, New Math Library, 1991
- 8. M. Kac and S. M. Ulam, Mathematics and Logic, Dover Publications, NY, 1992
- 9. K. Kuratowski and A. Mostowski, Set Theory, North-Holand, Amsterdam, 1967.
- 10. S. Savchev, T. Andreescu, Mathematical Miniatures, MAA, 2003
- 11. I. F. Sharygin, Mathematical Vinegrette, Mir, 2002 (in Russian)
- 12. D. Wells, The Penguin Book of Curious and Interesting Puzzles, Penguin Books, 1992
- 13. P. Zeitz, The Art and Craft of Problem Solving, John Wiley & Sons, 1999



Related material

Read more...

- Twenty five boys and twenty five girls
- · Pigeonhole in Chess Training
- · Married Couples at a Party
- Jumping Isn't Everything
- Zeros and Nines
- Teams In a Tournament
- Divisibility of a Repunit
- Pigeonhole in Clubs



|Contact| |Front page | |Contents | |Did you know? | |Algebra | |Store |

Copyright © 1996-2014 Alexander Bogomolny

45132144

CITE THIS PAGE AS: A. Bogomolny, <u>Pigeonhole Principle and Extensions</u> from Interactive Mathematics Miscellany and Puzzles **Healthman Cut-the-knot ora/pigeonhole/PigeonholeExtensions.shtml, Accessed 17 February 2014 🖸 BOOKMARK 🔣 💺 🖂 ...) ALSO ON INTERACTIVE MATHEMATICS MISCELLANY AND PUZZLES WHAT'S THIS? e and pi Spoof 6 comments How to write an equation of the union of two sets? 3 comments The Law of Cosines by Larry Hoehn 2 comments **Angle Bisectors Cross Circumcircle** 2 comments Login 0 Comments **Interactive Mathematics Miscellany and Puzzles** Sort by Best 🔻 Share 🔁 Favorite * Start the discussion... Be the first to comment. Add Disqus to your site Subscribe Subscribe