

SOCIOL 723: Social Statistics II

Week 2, Regression Inference and Robust Standard Errors

Wenhao Jiang

Department of Sociology, Fall 2026



Today

The Sampling Distribution of OLS

Hypothesis Testing

Heteroskedasticity and Robust Standard Errors

Clustered Data

The Bootstrap

What Robust Standard Errors Cannot Fix

★ = essential, you must understand this

★ = for understanding, not examined

Where We Left Off

Last week, for the bivariate model $Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$:

$$\hat{\beta}_1 - \beta_1 = \frac{\sum_i (X_i - \bar{X}) \epsilon_i}{\sum_i (X_i - \bar{X})^2}, \quad \text{Var}(\hat{\beta}_1 | X) = \frac{\sum_i (X_i - \bar{X})^2 \text{Var}(\epsilon_i | X)}{[\sum_i (X_i - \bar{X})^2]^2},$$

which collapses to $\sigma^2 / \sum_i (X_i - \bar{X})^2$ only if $\text{Var}(\epsilon_i | X)$ is the same for every i .

Today's agenda

1. What is the sampling distribution of $\hat{\beta}_1$ as n grows?
2. How do we turn a standard error into a test and a confidence interval?
3. What if $\text{Var}(\epsilon_i | X)$ is *not* constant? (Heteroskedasticity, then clustering.)
4. The bootstrap as a general-purpose alternative.
5. What robust standard errors can and cannot fix.

Reminder: the Two-Star Convention

Frames marked ★ are **essential**: problem sets and exams draw on them directly. Frames marked ☆ are here for *understanding*, not for evaluation: matrix derivations, asymptotic arguments, and the general formulas you will meet in the literature.

- Examinable this week: the scalar variance formula and where it comes from; what heteroskedasticity and clustering *are* and why they matter; how to read and choose a standard error; what a bootstrap does; and the list of problems robust standard errors do not solve.
- Every standard error in this course starts from the variance of a weighted sum of errors. HC estimators estimate the individual *variance* terms; cluster-robust estimators also estimate the within-cluster *covariance* terms. Hold on to that sentence.

Sampling Distribution

The Question, in Plain Words ★

Last week we computed $\hat{\beta}_1 = 0.119$: one more year of schooling goes with about 12% higher earnings in our sample of 3,509 people.

- The GSS could just as easily have interviewed a *different* 3,509 people. That sample would have produced a different number, perhaps 0.112, perhaps 0.127.
- So 0.119 is one draw from a lottery. Before treating it as evidence about the population, we need to know how much it moves from draw to draw.
- The **standard error** is exactly that: the give-or-take attached to the estimate. Formally, it is (an estimate of) the standard deviation of $\hat{\beta}_1$ across hypothetical repeated samples.

This week in one sentence

Everything called “statistical inference”, standard errors, t -statistics, p -values, confidence intervals, is machinery for learning that give-or-take number from the single sample we actually have.

What a Sampling Distribution Is ★

$\hat{\beta}_1$ is a number computed from *your* sample. Had you drawn a different sample from the same population, you would have got a different number.

The thought experiment

Imagine drawing sample after sample from the population and computing $\hat{\beta}_1$ each time. The histogram of those numbers is the **sampling distribution**. Where it is centered tells you about bias; how wide it is is the true sampling standard deviation, and the *standard error* is our estimate of that width.

- You only ever see one draw. Everything in this lecture is about inferring the shape of that histogram from a single sample.
- In lab we run the experiment by simulation, which is the fastest way to make the idea concrete.

Two Theorems We Lean On All Semester ★

Both are statements about what happens to a *sample average* as n grows.

Law of large numbers (LLN)

The average of n independent draws settles down on the population mean:

$\frac{1}{n} \sum_i Z_i \xrightarrow{p} E[Z]$ (read $\xrightarrow{p}$ as “settles down on”). Flip a fair coin 10 times and the share of heads could be anything; flip it 10,000 times and it sits within a whisker of 0.5.

Central limit theorem (CLT)

For an average of *independent* draws, the *shape* of the wobble around the truth becomes a bell curve, $\sqrt{n}(\bar{Z} - E[Z]) \xrightarrow{d} \mathcal{N}(0, \text{Var}(Z))$ (read $\xrightarrow{d}$ as “the distribution approaches”), **whatever the distribution of Z itself**, so long as its variance is finite.

Why we care: $\hat{\beta}_1$ is built from sample averages, so the LLN gives consistency and the CLT approximate normality. Thursday’s lab watches the bell curve emerge.

The Sampling Distribution of $\hat{\beta}_1$, as n Grows

The big picture: keep drawing samples from the population; as n grows, we will show that $\hat{\beta}_1$'s sampling behaviour approximates a normal distribution, without ever assuming the errors are normal (A6):

$$\hat{\beta}_1 - \beta_1 = \frac{\frac{1}{n} \sum_i (X_i - \bar{X}) \epsilon_i}{\frac{1}{n} \sum_i (X_i - \bar{X})^2}.$$

A normal is pinned down by its *mean* and *variance*, so the work here is to find both, and let the CLT supply the *shape*.

The plan:

1. The denominator converges to a constant (LLN).
2. The numerator, magnified by $\sqrt{n}$, converges to a normal distribution with mean 0 (CLT) under strict exogeneity (A4: $E[\epsilon_i | X] = 0$).
3. A normal distribution divided by a settled constant is still normal.

Step 1: The Denominator Settles (LLN)

The denominator is the sample variance of X . By the law of large numbers

$$\begin{aligned}\bar{X} &\xrightarrow{p} E[X] = \mu_X, \\ \frac{1}{n} \sum_i (X_i - \bar{X})^2 &\xrightarrow{p} E[(X_i - \mu_X)^2] = \text{Var}(X_i)\end{aligned}$$

- It settles on the population variance, a *positive constant*.
- As $n \rightarrow \infty$, the observed X 's approach, in aggregate, the population distribution of X .
- From here on we may treat the denominator as the constant $\text{Var}(X_i)$. Every remaining question is about the numerator, whose terms we may likewise write with μ_X in place of $\bar{X}$: it is the average of the $(X_i - \mu_X) \epsilon_i$.

Step 2: The Numerator's Mean

The numerator averages the identically distributed terms $(X_i - \mu_X) \epsilon_i$, so its mean is the terms' common mean. Compute it:

$$\begin{aligned}
 E[(X_i - \mu_X) \epsilon_i] &= E\left[E[(X_i - \mu_X) \epsilon_i \mid X_i]\right] && \text{iterated expectations} \\
 &= E[(X_i - \mu_X) E[\epsilon_i \mid X_i]] && \text{given } X_i, (X_i - \mu_X) \text{ is fixed} \\
 &= E[(X_i - \mu_X) \cdot 0] = 0 && \text{A4: } E[\epsilon_i \mid X_i] = 0
 \end{aligned}$$

- By the LLN the numerator settles on this mean, 0, so $\hat{\beta}_1 - \beta_1 \xrightarrow{p} 0 / \text{Var}(X_i) = 0$: **consistency**. As sample approaches population, $\hat{\beta}_1$ nails β_1 .
- But a mean says nothing about *spread*, and spread is what a standard error measures. Next: the variance.

Step 3: The Numerator's Variance

Take $n = 3$: the numerator averages the three terms $(X_i - \mu_X)\epsilon_i$. The variance of a sum is every term's variance plus a covariance for every pair:

$$\begin{aligned} \text{Var}(\text{numerator}) = \frac{1}{9} & \left[\underbrace{\text{Var}((X_1 - \mu_X)\epsilon_1) + \text{Var}((X_2 - \mu_X)\epsilon_2) + \text{Var}((X_3 - \mu_X)\epsilon_3)}_{\text{one variance per draw}} \right. \\ & \left. + 2 \underbrace{\text{Cov}((X_1 - \mu_X)\epsilon_1, (X_2 - \mu_X)\epsilon_2) + \dots}_{\text{one covariance per pair; 3 pairs}} \right] \end{aligned}$$

- A2 kills every covariance: draws are *independent*, so each Cov is 0. Clustering breaks exactly this step.
- Merging the remaining variance terms, with n draws, $\text{Var}((X_i - \mu_X)\epsilon_i)/n$.
- The spread shrinks like $1/\sqrt{n}$: **magnify by $\sqrt{n}$** to hold it steady. That is where the $\sqrt{n}$ comes from.

Step 4: The CLT Adds the Shape

The magnified numerator has mean 0 and variance $\text{Var}((X_i - \mu_X)\epsilon_i)$; the CLT adds the missing *shape*. Divide by the settled denominator:

$$\sqrt{n}(\hat{\beta}_1 - \beta_1) = \frac{\overbrace{\sqrt{n} \cdot \frac{1}{n} \sum_i (X_i - \bar{X}) \epsilon_i}^{\xrightarrow{d} \mathcal{N}(0, \text{Var}((X_i - \mu_X)\epsilon_i))}}{\underbrace{\frac{1}{n} \sum_i (X_i - \bar{X})^2}_{\xrightarrow{p} \text{Var}(X_i)}} \xrightarrow{d} \mathcal{N}(0, V), \quad V = \frac{\text{Var}((X_i - \mu_X)\epsilon_i)}{[\text{Var}(X_i)]^2}.$$

- Under A5, homoskedasticity, $\text{Var}(\epsilon_i | X_i) = \sigma^2$: the error variance is the same number at every value of X . The numerator of V then collapses and $V = \sigma^2 / \text{Var}(X_i)$, the familiar formula: next frame, slowly.
- Without A5, **estimating V is this lecture's entire subject.**

Under A4 and A5, the Meat Collapses

The meat is a mean square around the unconditional mean, so A4 first certifies that mean is zero; then A5 collapses it:

$$\begin{aligned}
 E[(X_i - \mu_X)^2 \epsilon_i^2] &= E\left[E[(X_i - \mu_X)^2 \epsilon_i^2 \mid X_i] \right] && \text{iterated expectations} \\
 &= E[(X_i - \mu_X)^2 E[\epsilon_i^2 \mid X_i]] && \text{given } X_i, (X_i - \mu_X)^2 \text{ is fixed} \\
 &= E[(X_i - \mu_X)^2 \sigma^2] && \text{A4 + A5: } E[\epsilon_i^2 \mid X_i] = \sigma^2 \\
 &= \sigma^2 E[(X_i - \mu_X)^2] = \sigma^2 \text{Var}(X_i) && \sigma^2 \text{ is one number: pull it out}
 \end{aligned}$$

So $V = \sigma^2 \text{Var}(X_i) / [\text{Var}(X_i)]^2 = \sigma^2 / \text{Var}(X_i)$: the same answer as Week 1.

Testing

The Vocabulary, Pinned Down ★

You met these words in *Social Statistics I*; now they have foundations.

- A **null hypothesis** H_0 is a specific claim about a population quantity, e.g. $\beta_1 = 0$: schooling and earnings are unrelated.
- A **test statistic** measures the distance between estimate and null *in standard-error units*: $t = (\hat{\beta}_1 - 0)/\text{se}(\hat{\beta}_1)$. Big distance, implausible null.
- The ***p*-value**: if H_0 were true, how often would sampling noise alone produce a t this large? It is a statement about the data under the null, **not** the probability the null is true.
- A **95% confidence interval** collects every null value the data cannot reject; across repeated samples, intervals built this way cover the truth 95% of the time. *This* interval either does or does not; the 95% describes the procedure.

Testing a Single Coefficient ★

To test $H_0 : \beta_j = \beta_j^0$, compare

$$t = \frac{\hat{\beta}_j - \beta_j^0}{\text{se}(\hat{\beta}_j)}$$



against t_{n-k} critical values; a 95% confidence interval is $\hat{\beta}_j \pm t_{0.975, n-k} \text{se}(\hat{\beta}_j)$.

- By the last section, t is approximately $\mathcal{N}(0, 1)$ for large n , and t_{n-k} merges into that same curve (the critical values differ by under 2% once $n - k > 100$); under the classical assumption of normal errors, t_{n-k} is even exact at any n , which is where the convention comes from. This is why regression output reports t_{n-k} ; clustered data will later change the right degrees of freedom.
- **Everything depends on whether se is right.** A t of 3.0 computed from a standard error that is half of what it should be is really a t of 1.5.

Heteroskedasticity

What Heteroskedasticity Is ★

Week 1's A5 had *two* parts: constant variance, and zero covariance across observations. This section drops the first while keeping the second; the clustering section then drops the second. **Homoskedasticity** says $\text{Var}(\epsilon_i | X_i) = \sigma^2$: the spread of the outcome around the line is the same everywhere.



Under exogeneity, both lines are estimated without bias: heteroskedasticity by itself does not bias $\hat{\beta}$. Only the *uncertainty* about them differs, and only the right-hand panel needs a different formula.

Heteroskedasticity Is the Norm

- Earnings dispersion rises steeply with education and experience: at 8 years of schooling almost everyone earns a modest wage; at 20 years the range is enormous.
- Group-level outcomes built from different numbers of individuals, a county mean over 50 people is noisier than one over 50,000.
- Binary outcomes: $\text{Var}(Y_i | X_i) = p(X_i)(1 - p(X_i))$ is *necessarily* heteroskedastic, so the linear probability model always violates A5.

What it does and does not do

Does not: bias $\hat{\beta}$ or break consistency (given A4 and random sampling).

Does: invalidate the standard errors your software prints by default, so t -statistics, p -values and confidence intervals are wrong; and cost efficiency (OLS is no longer BLUE).

The Scalar Sandwich ★

The last section already handed us the bell's variance, built from population moments:

$$V = \frac{E[(X_i - \mu_X)^2 \epsilon_i^2]}{[\text{Var}(X_i)]^2} = \frac{\text{"meat"}}{\text{"bread"}^2}.$$

- The meat is a $(X_i - \mu_X)^2$ -weighted average of the squared errors: errors sitting far from the mean of X count the most.
- Under A5 the squared error averages to σ^2 at every value of X , so the meat factors into $\sigma^2 \text{Var}(X_i)$ and $V = \sigma^2 / \text{Var}(X_i)$: the classical formula is the A5 special case.
- Without A5 nothing factors, and the classical formula is simply wrong. But every piece of V is a population mean, and sample averages estimate population means. Next frame.

White's Estimator, in Scalar Form ★

Estimate each population mean by its sample counterpart, with residuals standing in for the unobservable errors: writing $d_i \equiv X_i - \bar{X}$, the bread $\text{Var}(X_i)$ becomes $\frac{1}{n} \sum_i d_i^2$ and the meat $E[(X_i - \mu_X)^2 \epsilon_i^2]$ becomes $\frac{1}{n} \sum_i d_i^2 \hat{e}_i^2$. The squared standard error of $\hat{\beta}_1$ is this estimate of V , divided by n ; the $1/n$'s cancel into

$$\boxed{\hat{V}_{\text{HC0}}(\hat{\beta}_1) = \frac{\sum_i d_i^2 \hat{e}_i^2}{(\sum_i d_i^2)^2}} \quad \text{versus} \quad \hat{V}_{\text{classical}}(\hat{\beta}_1) = \frac{s^2}{\sum_i d_i^2}.$$

- That is the whole of “heteroskedasticity-robust” and “(Eicker–)Huber–White” (White 1980); **Stata's robust** adds HC1 scaling, next frame.
- With controls, replace d_i by Week 1's FWL residual $\tilde{d}_i$:
 $\hat{V}(\hat{\beta}_j) = \sum_i \tilde{d}_i^2 \hat{e}_i^2 / (\sum_i \tilde{d}_i^2)^2$: same formula, residualized weights.
- The classical formula pools *one* s^2 ; the robust one lets each observation carry its own $\hat{e}_i^2$.

“But $\hat{\epsilon}_i^2$ Is a Terrible Estimate of ϵ_i^2 ”

A frequent and reasonable objection: each $\hat{\epsilon}_i^2$ is a single noisy draw. How can plugging in n bad estimates give a good one?

- Because we never try to learn any individual ϵ_i^2 . The target is *one number*, the population mean $E[(X_i - \mu_X)^2 \epsilon_i^2]$, and $\frac{1}{n} \sum_i d_i^2 \hat{\epsilon}_i^2$ is a sample average aimed at it: the law of large numbers averages the individual noise away.
- Same logic that lets you estimate $\text{Var}(\bar{Y})$ from one sample: you never learn any individual's deviation, only the average square.

From Scalar to Matrix: the Dictionary ★

One change of meaning: X_i now denotes the *vector* of observation i 's regressors, in the bivariate case $X_i = (1, X_i)'$. Then

$$X_i X_i' = \begin{pmatrix} 1 & X_i \\ X_i & X_i^2 \end{pmatrix}, \quad X_i \epsilon_i = \begin{pmatrix} \epsilon_i \\ X_i \epsilon_i \end{pmatrix},$$

and their sample averages stack the scalar sums you have used all lecture:

$$\mathbb{E}_n[X_i X_i'] = \begin{pmatrix} 1 & \bar{X} \\ \bar{X} & \frac{1}{n} \sum_i X_i^2 \end{pmatrix}, \quad \mathbb{E}_n[X_i \epsilon_i] = \begin{pmatrix} \bar{\epsilon} \\ \frac{1}{n} \sum_i X_i \epsilon_i \end{pmatrix}.$$

- Population versions: $\mathbf{Q} = E[X_i X_i']$ and the meat $\mathbf{\Omega} = E[\epsilon_i^2 X_i X_i']$, which stacks $E[\epsilon_i^2]$, $E[X_i \epsilon_i^2]$, $E[X_i^2 \epsilon_i^2]$.
- The next frames state the matrix result and then cash it: the slope's entry of the matrix sandwich is *exactly* the scalar V .

The General Statement ★

For the full coefficient vector, writing sample averages as $\mathbb{E}_n$:

$$\hat{\beta} - \beta = (\mathbb{E}_n[X_i X_i'])^{-1} \mathbb{E}_n[X_i \epsilon_i].$$

The LLN gives $\mathbb{E}_n[X_i X_i'] \xrightarrow{p} E[X_i X_i'] \equiv \mathbf{Q}$ and $\mathbb{E}_n[X_i \epsilon_i] \xrightarrow{p} \mathbf{0}$, so $\hat{\beta} \xrightarrow{p} \beta$ (*consistency*). Multiplying by $\sqrt{n}$ and applying the CLT to the second factor,

$$\boxed{\sqrt{n}(\hat{\beta} - \beta) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \mathbf{Q}^{-1} \mathbf{\Omega} \mathbf{Q}^{-1})}, \quad \mathbf{\Omega} = E[\epsilon_i^2 X_i X_i'].$$

Note what is *not* assumed: no normality, no homoskedasticity. Note also the weaker exogeneity requirement, $E[X_i \epsilon_i] = 0$ suffices for consistency; $E[\epsilon_i | X_i] = 0$ is not needed.

Reading the “Sandwich” ★

$$V = \underbrace{Q^{-1}}_{\text{bread}} \underbrace{\Omega}_{\text{meat}} \underbrace{Q^{-1}}_{\text{bread}}, \quad Q = E[X_i X_i'], \quad \Omega = E[\epsilon_i^2 X_i X_i'].$$

- The **bread** depends only on the distribution of the regressors.
- The **meat** is where every assumption about the error structure lives. Classical, robust, clustered, and HAC standard errors differ **only** in how the meat is estimated.
- If $E[\epsilon_i^2 | X_i] = \sigma^2$ then $\Omega = \sigma^2 Q$ and $V = \sigma^2 Q^{-1}$, the classical formula.

The organizing idea for the whole lecture

Keep the estimator $\hat{\beta}$; change how you estimate the meat.

The Dictionary, Cashed: Multiply the Sandwich Out ★

Every symbol below is a population *number*. Abbreviate $v = \text{Var}(X_i)$, $q = E[X_i^2]$, and the meat's three entries $a = E[\epsilon_i^2]$, $b = E[X_i \epsilon_i]$, $c = E[X_i^2 \epsilon_i]$. The ingredients ($\mathbf{Q}^{-1}$ by the 2×2 inverse, $\det \mathbf{Q} = q - \mu_X^2 = v$):

$$\mathbf{Q}^{-1} = \frac{1}{v} \begin{pmatrix} q & -\mu_X \\ -\mu_X & 1 \end{pmatrix}, \quad \mathbf{\Omega} = \begin{pmatrix} a & b \\ b & c \end{pmatrix}.$$

First multiplication, plain row-times-column arithmetic:

$$\mathbf{\Omega} \mathbf{Q}^{-1} = \frac{1}{v} \begin{pmatrix} aq - b\mu_X & b - a\mu_X \\ bq - c\mu_X & c - b\mu_X \end{pmatrix}.$$

One more multiplication by $\mathbf{Q}^{-1}$ from the left: next frame.

The Final 2×2 Matrix ★

Multiply again, row times column; for instance the bottom-right entry is $(-\mu_X)(b - a\mu_X) + 1 \cdot (c - b\mu_X) = a\mu_X^2 - 2b\mu_X + c$. Collecting all four entries:

$$\mathbf{Q}^{-1}\mathbf{\Omega}\mathbf{Q}^{-1} = \frac{1}{v^2} \begin{pmatrix} aq^2 - 2bq\mu_X + c\mu_X^2 & b(q + \mu_X^2) - \mu_X(aq + c) \\ b(q + \mu_X^2) - \mu_X(aq + c) & a\mu_X^2 - 2b\mu_X + c \end{pmatrix}.$$

The highlighted slope entry, with a, b, c substituted back:

$$a\mu_X^2 - 2b\mu_X + c = E[\mu_X^2\epsilon_i^2 - 2\mu_X X_i\epsilon_i^2 + X_i^2\epsilon_i^2] = E[(X_i - \mu_X)^2\epsilon_i^2],$$

so the $(2, 2)$ entry is $E[(X_i - \mu_X)^2\epsilon_i^2]/v^2 = V$: the scalar sandwich, recovered.

Reading the rest: the diagonal holds the intercept's and the slope's variances, the off-diagonal their covariance.

Why the Squared Residuals Are Systematically Too Small

Two frames ago: each $\hat{\epsilon}_i^2$ is a *noisy* stand-in for ϵ_i^2 , and the meat's average kills the noise. True, but **averaging only kills noise; a systematic error survives the average** – and there is one.

- OLS chooses the line that makes the residuals as small as possible, so it partly fits the noise itself: residuals run systematically *smaller* than the errors they stand in for.
- The exact bookkeeping (under A5, for clean arithmetic): $E[\hat{\epsilon}_i^2 | X] = \sigma^2(1 - h_{ii})$, where the **leverage** $h_{ii} = \frac{1}{n} + d_i^2 / \sum_j d_j^2$ measures how unusual X_i is.
- So every $\hat{\epsilon}_i^2$ is deflated, and most at *high-leverage* observations, exactly the ones the meat weights most heavily. The meat estimate, hence HC0, is **biased down in small samples**.
- As n grows, $h_{ii} \rightarrow 0$ and the deflation vanishes: a purely finite-sample issue. The target is still only the meat; the fixes just undo a known bias in its ingredients.

Where $\sigma^2(1 - h_{ii})$ Comes From ★

Write the residual in terms of the errors, step by step:

$$\begin{aligned}
 \hat{e}_i &= Y_i - \hat{\beta}_0 - \hat{\beta}_1 X_i && \text{definition} \\
 &= \epsilon_i - \bar{\epsilon} - (\hat{\beta}_1 - \beta_1) d_i && \text{sub } Y_i = \beta_0 + \beta_1 X_i + \epsilon_i, \hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X} \\
 &= \epsilon_i - \bar{\epsilon} - d_i \frac{\sum_j d_j \epsilon_j}{\sum_j d_j^2} = \epsilon_i - \sum_j w_{ij} \epsilon_j && \text{Week 1: } \hat{\beta}_1 - \beta_1 = \sum_j d_j \epsilon_j / \sum_j d_j^2
 \end{aligned}$$

with $w_{ij} = \frac{1}{n} + d_i d_j / \sum_k d_k^2$: the fitted line absorbs a weighted share of *every* error, with weight $w_{ii} = h_{ii}$ on observation i 's own. Under A5 and uncorrelated errors,

$$E[\hat{e}_i^2 \mid X] = \sigma^2 \left(1 - 2h_{ii} + \sum_j w_{ij}^2 \right) = \sigma^2 (1 - h_{ii}),$$

because $\sum_j w_{ij}^2 = h_{ii}$ (expand the square; $\sum_j d_j = 0$).

Reading: observation i pulls the line toward itself with strength h_{ii} ; the residual keeps the $(1 - h_{ii})$ share. At $n = 2$, $h_{ii} = 1$: the line passes through both points and both residuals are exactly zero.

Finite-Sample Corrections: HC0–HC3 ★

Every $\hat{e}_i^2$ is deflated by its leverage. The corrections rescale each squared residual before it enters the meat, undoing the deflation (factors motivated by the homoskedastic benchmark, used regardless):

	Meat uses	Comment
HC0	$\hat{e}_i^2$	White's original; anticonservative when n is small
HC1	$\frac{n}{n-k} \hat{e}_i^2$	simple d.f. correction; Stata's robust default
HC2	$\frac{\hat{e}_i^2}{1 - h_{ii}}$	exact at the homoskedastic benchmark
HC3	$\frac{\hat{e}_i^2}{(1 - h_{ii})^2}$	jackknife-like; recommended default

All four are consistent under any heteroskedasticity pattern and coincide as $n \rightarrow \infty$. MacKinnon and White (1985), Long and Ervin (2000): with $n < 250$ or high leverage, use **HC3**.

How Much Does It Matter?

- Robust standard errors can be *larger or smaller* than classical ones. There is no general direction.
- The difference is driven by whether the error variance happens to be large where X is extreme. In our GSS earnings regression it is modest: $\text{se}(\hat{\beta}_{\text{educ}})$ goes from 0.00510 (classical) to 0.00547 (HC0) to 0.00549 (HC3), about 7%.
- With a skewed or sparse regressor, a rare binary treatment, a long-tailed count; the difference can be a factor of two.
- Clustering, by contrast, routinely changes standard errors by factors of three to ten. In most sociological data that is the bigger issue.

Should You *Test* for Heteroskedasticity?

Classical tests exist: Breusch–Pagan, White.

Recommendation: no

Do not pre-test and then choose. Just report robust standard errors.

- A pre-test makes your final inference conditional on the pre-test outcome, which distorts the very sampling distribution you are claiming.
- The cost of robust standard errors when the errors happen to be homoskedastic is small, a modest efficiency loss in the *variance estimate*.
- The cost of classical standard errors when they are wrong is invalid inference.
- Exception: the tests are useful *diagnostically*. Heteroskedasticity is often a symptom of a misspecified functional form, which is worth knowing about for its own sake.

Clustering

Why Independence Fails ★

Sociological data are almost never i.i.d. draws:

- Students within schools; workers within firms; siblings within families.
- Repeated observations on the same individual (panel data).
- Multi-stage survey designs, the GSS samples tracts, then households.
- Treatment assigned at the group level (a state law), outcomes measured at the individual level.

Error components model

$$Y_{ig} = \beta_0 + \beta_1 X_{ig} + \underbrace{\nu_g}_{\text{cluster shock}} + \underbrace{\eta_{ig}}_{\text{idiosyncratic}}, \quad \rho_e = \frac{\sigma_\nu^2}{\sigma_\nu^2 + \sigma_\eta^2}$$

Everyone in cluster g shares the shock ν_g , so their errors are correlated. ρ_e is the **intra-class correlation**.

Where the Scalar Formula Breaks ★

Hold the X 's fixed for the bookkeeping. The variance of the error sum then involves *all* the covariances:

$$\text{Var}\left(\sum_i d_i \epsilon_i \mid X\right) = \underbrace{\sum_i d_i^2 \sigma_i^2}_{\text{what HC0 estimates}} + \underbrace{\sum_i \sum_{j \neq i} d_i d_j \text{Cov}(\epsilon_i, \epsilon_j \mid X)}_{\text{assumed zero so far}},$$

with $\sigma_i^2 = \text{Var}(\epsilon_i \mid X_i)$, the error spread at X_i .

- Independent sampling makes the second term vanish. Clustering does not.
- If errors within a cluster are *positively* correlated and d_i has the same sign within clusters, every one of those cross terms is positive, so the true variance is **larger** than the formula we have been using, often by a lot.
- There are on the order of $n(\bar{n} - 1)$ such cross terms versus n terms in the first sum, so even tiny correlations can dominate.

The Moulton Factor: How Bad Is It? ★

With equal cluster sizes $\bar{n}$ and an error-components structure, Moulton (1986) showed

$$\frac{\text{Var}_{\text{clustered}}(\hat{\beta}_1)}{\text{Var}_{\text{OLS}}(\hat{\beta}_1)} = 1 + (\bar{n} - 1) \rho_x \rho_e,$$

where ρ_x is the intra-class correlation of the regressor.

- If the regressor is *constant within clusters*, treatment assigned at the cluster level, then $\rho_x = 1$ and the inflation is $1 + (\bar{n} - 1)\rho_e$. This is the worst case.
- 50 states, 1,000 individuals each, $\rho_e = 0.01$ (small!): $\sqrt{1 + 999(0.01)} \approx 3.3$.
Standard errors are more than three times too small, so a reported t of 6 is really a t of 1.8.
- Moral: the effective sample size is closer to the number of *clusters* than the number of observations.

The Cluster-Robust Fix, in Scalar Form ★

With clusters the true meat also contains the within-cluster cross terms. The fix: **sum within clusters first, then square**:

$$\hat{V}_{\text{CR}}(\hat{\beta}_1) = \frac{\sum_{g=1}^G \left(\sum_{i \in g} d_i \hat{e}_i \right)^2}{\left(\sum_i d_i^2 \right)^2}.$$

- Expand the square: $\left(\sum_{i \in g} d_i \hat{e}_i \right)^2 = \sum_{i \in g} d_i^2 \hat{e}_i^2 + \sum_{i \neq j \in g} d_i d_j \hat{e}_i \hat{e}_j$. The cross terms we had been assuming away are now **estimated**, within clusters; across clusters they are still assumed zero.
- Same logic as White: no single cluster's covariances are estimated well, but the *sum* over many clusters is. Consistency therefore runs in G , the number of clusters, not in n .
- With one observation per cluster this collapses exactly to HC0: clustering strictly generalizes White.

Cluster-Robust Variance (Liang–Zeger) ★

Let $g = 1, \dots, G$ index clusters, with $\mathbf{X}_g$ and $\hat{\mathbf{e}}_g$ the stacked regressors and residuals in cluster g :

$$\hat{V}_{\text{CR}}(\hat{\beta}) = c (\mathbf{X}'\mathbf{X})^{-1} \left(\sum_{g=1}^G \mathbf{X}'_g \hat{\mathbf{e}}_g \hat{\mathbf{e}}'_g \mathbf{X}_g \right) (\mathbf{X}'\mathbf{X})^{-1}$$

with the usual small-sample scaling $c = \frac{G}{G-1} \cdot \frac{n-1}{n-k}$ (CR1; Stata's default).

- Same sandwich; the meat now sums outer products of *cluster* score vectors instead of individual ones, which is exactly how the cross terms get included.
- Allows **arbitrary** correlation within cluster (including serial correlation over time) and any heteroskedasticity, while assuming independence *across* clusters.
- Consistency is in $G \rightarrow \infty$, not $n \rightarrow \infty$.

At What Level Should You Cluster? ★

Abadie et al. (2023): clustering is a statement about the *design*, not a knob to tune.

- **Sampling-based:** cluster at the level of the sampling units, when the clusters in your data are a small share of the clusters in the population.
- **Design-based:** cluster at the level at which *treatment* is assigned. If a policy varies by state, cluster on state, even with individual-level data.
- Cluster at the level the sampling or assignment design justifies. Never finer than the assignment level; going *coarser* needs its own dependence argument, and is not automatically safer.
- “Cluster on everything” is not free: with few clusters the estimator is biased downward and the test over-rejects.
- **Never** choose the level by whichever gives the smallest p -value.

The Few-Clusters Problem

With small G , $\hat{V}_{\text{CR}}$ is biased downward, and even the usual cluster-robust reference distribution (t with G -based degrees of freedom, commonly $G - 1$, not $n - k$) is too thin. As rules of thumb, worry below $G \approx 40$ – 50 and be alarmed below 20 , though performance also depends on cluster-size imbalance and leverage. A typical symptom: rejection rates of 10–15% for a nominal 5% test.

Remedies

1. **CR2 with Satterthwaite degrees of freedom** (Bell and McCaffrey 2002; Imbens and Kolesar 2016): a leverage correction (the cluster analogue of HC2) plus data-dependent d.f. `clubSandwich::coef_test()`.
2. **Wild cluster bootstrap- t** (Cameron et al. 2008): impose H_0 , resample cluster-level ± 1 weights, and build the reference distribution empirically. `fwildclusterboot`.
3. **Aggregate**: collapse to G cluster-level observations and analyze those. Honest, but note it reweights and can change the estimand.
4. **Randomization inference** when treatment is assigned to few units.

Serial Correlation: Bertrand, Duflo, and Mullainathan (2004)

A famous demonstration of what happens when you ignore within-unit correlation.

- They ran *placebo* difference-in-differences: take CPS data, invent a state-level “law” beginning in a random year, and estimate its effect on women’s wages. The true effect is zero by construction.
- With conventional standard errors the fake law came out “significant” at 5% in 45% of the placebo runs.
- Cause: state-level outcomes are highly serially correlated over time, and the standard errors treated 50×21 observations as independent.
- Clustering by state brought rejection rates close to nominal here.

The rule for any policy-adoption design, sociology’s most common clustering decision: **cluster at the level of the adopting unit**. Week 10 returns to this; design and inference are not separable.

Other Dependence Structures ★

- **Two-way / multi-way clustering** (Cameron et al. 2011): firms *and* years, individuals *and* places. Add the two cluster-robust matrices and subtract their intersection.
- **HAC (Newey–West)**: time-series dependence of unknown form, handled with a bandwidth-weighted sum of autocovariances. For one long series rather than many short ones.
- **Spatial HAC (Conley)**: correlation decaying in geographic distance.
- **Network dependence**: correlation along a graph; an active area with fewer settled answers.

All are the same sandwich with a different meat. Once you can read $\hat{V}$, you can read any of these off the formula.

Bootstrap

The Idea

Everything so far derived a *formula* for the sampling distribution. The bootstrap instead **simulates** it.

Plug-in principle (Efron 1979)

We want the sampling distribution of a statistic when samples are drawn from the population F . We do not know F , but the sample *is* our best estimate of F . So draw new samples from the sample.

- In our notation: replace E by $\mathbb{E}_n$, and resample.
- Especially valuable for statistics with no convenient closed-form variance: quantiles, ratios, decompositions, two-step estimators, indirect effects.

The Nonparametric (Pairs) Bootstrap ★

Notation: θ is any target, $\hat{\theta}$ its estimate, B the number of resamples.

1. For $b = 1, \dots, B$: draw n observations (Y_i, X_i) *with replacement* from the sample.
2. Re-estimate on the resample to get $\hat{\theta}^{(b)}$.
3. Use the empirical distribution of $\{\hat{\theta}^{(b)}\}_{b=1}^B$.
 - se_{boot} = the standard deviation of the $\hat{\theta}^{(b)}$.
 - **Percentile CI**: the 2.5th and 97.5th percentiles of $\{\hat{\theta}^{(b)}\}$.
 - Because it resamples (Y_i, X_i) jointly, the pairs bootstrap is automatically heteroskedasticity-robust, given *independent* observations; it assumes nothing about how the error variance moves with X .
 - $B = 1,000$ for standard errors; $B \geq 5,000$ for percentile intervals or p -values.

Variants

Variant	What is resampled	When to use
Pairs	(Y_i, X_i) jointly	default; robust to heteroskedasticity
Residual	$\hat{e}_i$, then $Y_i^* = \hat{Y}_i + \hat{e}_i^*$	requires homoskedasticity; fixed design
Wild	$Y_i^* = \hat{Y}_i + \hat{e}_i v_i, v_i \in \{-1, 1\}$	heteroskedastic, small n
Block / cluster	whole clusters	clustered data
Wild cluster	cluster-level v_g, H_0 imposed	few clusters

Rademacher weights: $v_i = +1$ or -1 with probability $1/2$ each, so $E[v_i] = 0$ and $E[v_i^2] = 1$, the resampled errors keep the right variance observation by observation.

Confidence Intervals: Not All Equal

- **Normal-approximation:** $\hat{\theta} \pm 1.96 \text{se}_{\text{boot}}$. Simple; assumes symmetry.
- **Percentile:** quantiles of $\{\hat{\theta}^{(b)}\}$. Handles skew and respects transformations.
- **BC_a:** bias-corrected and accelerated; second-order accurate and the usual recommendation for a percentile-type interval.
- **Bootstrap-*t* (studentized):** bootstrap the *t statistic* $(\hat{\theta}^{(b)} - \hat{\theta})/\text{se}^{(b)}$ and use its quantiles. Its approximation error shrinks *faster* with n than the other intervals' ★. Requires a standard error at each replication.

The wild cluster bootstrap-*t* is exactly this idea applied to the few-clusters problem, which is why it works so much better than a naive cluster bootstrap.

When the Bootstrap Fails

The bootstrap is not magic; the statistic must be *regular* enough (in a smoothness-like sense; quantiles qualify, the cases below do not).

- **Non-smooth estimators:** the sample maximum or minimum; nearest-neighbour matching estimators (Abadie and Imbens 2008).
- **Parameters on a boundary**, a variance component at zero.
- **Heavy tails** with infinite variance.
- **Dependence the resampling scheme does not preserve.** If you resample individuals when the data are clustered, the bootstrap reproduces the *wrong*, too-small standard error just as faithfully as the analytic formula does.

Bootstrap the design

Resample the units that were independently sampled. That single rule prevents most bootstrap mistakes.

Limits

Freedman's Critique

Freedman (2006) makes an uncomfortable point.

The argument

The sandwich gives a consistent estimate of the variance of $\hat{\beta}$ around the *population regression coefficient* β . If the model is misspecified, if the CEF is not $X_i'\beta$, then β is merely the best linear approximation, and a correct standard error for a quantity you do not care about is small comfort.

- Distinguish three failures: a **nonlinear CEF** (OLS still consistently estimates the BLP from Week 1, so robust inference about *that* is fine); an **exogeneity failure** ($\hat{\beta}$ no longer estimates the causal quantity you wanted); a **wrong estimand** (the BLP may not answer the question).
- Robust standard errors repair neither of the last two: precise inference about the wrong quantity is small comfort.

A Scoreboard ★

Problem	Robust SEs?	What actually helps
Heteroskedasticity	yes	(given independent observations)
Within-cluster correlation	yes (cluster-robust)	design-based clustering
Serial correlation in panels	yes	cluster on unit
Non-normal errors	yes (large- n approx.)	larger n
Omitted variable bias	no	identification strategy (Wks 5–11)
Simultaneity, reverse causation	no	a design (Weeks 7–10)
Measurement error in X	no	IV, validation data
Nonlinear CEF (CEF the target)	no	saturate, splines, flexible ML
Selection into the sample	no	modelling the selection
Wrong estimand	no	say what you are estimating
Influential outliers	no	diagnostics; robust regression
p -hacking, specification search	no	pre-registration

A Practical Protocol ★

1. State the estimand and the design *before* choosing a standard error.
2. **Identify the independently sampled or independently assigned units, and cluster there.** Use HC2 or HC3 only when observations are plausibly independent.
3. Report the number of clusters G . If $G < 40$, use CR2 with Satterthwaite d.f. or a wild cluster bootstrap, and say so.
4. If robust and classical standard errors differ dramatically, treat it as a diagnostic: something about the specification or the influence structure deserves attention.
5. Report the specification you pre-committed to, then robustness checks, not the reverse.

Looking Ahead

- We now have a complete account of OLS: what it estimates, and how to quantify our uncertainty about it.
- **Weeks 3–4:** maximum likelihood. The same asymptotic machinery, score, information, sandwich, returns in a more general form, and OLS turns out to be a special case.
- **Weeks 5+:** the harder question. Not “how precise is $\hat{\beta}$?” but “is β the thing I want?”

Problem Set 1

Assigned today, due Monday September 14 at 11:59PM. Covers Weeks 1–2: scalar OLS algebra, FWL, omitted variable bias, and robust/clustered inference on the GSS extract.

References

- Abadie, Alberto and Guido W. Imbens. 2008. “On the Failure of the Bootstrap for Matching Estimators.” *Econometrica* 76(6): 1537–1557.
- Abadie, Alberto, Susan Athey, Guido W. Imbens, and Jeffrey M. Wooldridge. 2023. “When Should You Adjust Standard Errors for Clustering?” *Quarterly Journal of Economics* 138(1): 1–35.
- Bell, Robert M. and Daniel F. McCaffrey. 2002. “Bias Reduction in Standard Errors for Linear Regression with Multi-Stage Samples.” *Survey Methodology* 28(2): 169–181.
- Bertrand, Marianne, Esther Duflo, and Sendhil Mullainathan. 2004. “How Much Should We Trust Differences-in-Differences Estimates?” *Quarterly Journal of Economics* 119(1): 249–275.
- Cameron, A. Colin, Jonah B. Gelbach, and Douglas L. Miller. 2008. “Bootstrap-Based Improvements for Inference with Clustered Errors.” *Review of Economics and Statistics* 90(3): 414–427.
- Cameron, A. Colin, Jonah B. Gelbach, and Douglas L. Miller. 2011. “Robust Inference with Multiway Clustering.” *Journal of Business & Economic Statistics* 29(2): 238–249.
- Efron, Bradley. 1979. “Bootstrap Methods: Another Look at the Jackknife.” *Annals of Statistics* 7(1): 1–26.
- Freedman, David A. 2006. “On the So-Called ‘Huber Sandwich Estimator’ and ‘Robust Standard Errors.’” *The American Statistician* 60(4): 299–302.
- Imbens, Guido W. and Michal Kolesár. 2016. “Robust Standard Errors in Small Samples: Some Practical Advice.” *Review of Economics and Statistics* 98(4): 701–712.

References (cont.)

- Long, J. Scott and Laurie H. Ervin. 2000. “Using Heteroscedasticity Consistent Standard Errors in the Linear Regression Model.” *The American Statistician* 54(3): 217–224.
- MacKinnon, James G. and Halbert White. 1985. “Some Heteroskedasticity-Consistent Covariance Matrix Estimators with Improved Finite Sample Properties.” *Journal of Econometrics* 29(3): 305–325.
- Moulton, Brent R. 1986. “Random Group Effects and the Precision of Regression Estimates.” *Journal of Econometrics* 32(3): 385–397.
- White, Halbert. 1980. “A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity.” *Econometrica* 48(4): 817–838.