



Model Validation in Extraction of the Inclusive Muon Neutrino Charged Current Cross Section at MicroBooNE

Xiangpan Ji and Wenqiang Gu (BNL)
On behalf of the MicroBooNE collaboration
APS April Meeting 2021


MicroBooNE Overview

- Micro Booster Neutrino Experiment


- Accelerator ν experiment at Fermilab
- LArTPC with 85 ton active mass
- Near-surface operation

- Main physics goals:

- Investigate MiniBooNE low-energy excess
- Measure ν -Ar interaction cross-sections




MicroBooNE detector




There are 16 presentations given by MicroBooNE at this meeting!

Measurements of ν_μ CC Inclusive Cross Section



Few measurements at low neutrino energy, especially in Argon target





Wire-Cell

Selection of Inclusive ν_μ CC Interactions

- The Wire-Cell 3D event reconstruction for LArTPC are used, achieving a **high rejection** of cosmic-ray backgrounds, a **high-performance** generic neutrino detection ([arXiv:2011.01375](#), [arXiv:2011.01375](#), [arXiv:2101.05076](#))
- The **high-statistics** event selection allows for high-precision cross-section measurements ([MICROBOONE-NOTE-1095-PUB](#))
- The selected events of inclusive ν_μ CC interactions: 93% purity and 64% efficiency

MicroBooNE Preliminary





Wire-Cell

Selection of Inclusive ν_μ CC Interactions


London Cooper-Troendle's talk on
Wire-Cell generic neutrino detection
(E18.2)

Hai Wang Yu's talk on
Wire-Cell pattern recognition (Z19.1)

Lee Hagman's talk on
Wire-Cell ν_e CC selection (H13.1)
Jay Hyun Jo's talk on
Wire-Cell eLEE search (H13.2)

Giacomo Scanavini's talk on
Wire-Cell ν_μ CC selection (S11.8)


MicroBooNE Preliminary



Validation of Neutrino Energy Modeling: E_ν to E_ν^{rec}

- Neutrino energy modeling is crucial to neutrino oscillation measurement
- Key challenge: understanding ν -Ar cross section as a function of energy
- A method: validate $E_{\text{had}}^{\text{rec}}$ after applying the constraint of muon kinematics distribution
 - Common systematics are suppressed, providing a more stringent validation

$$E_\nu^{\text{rec}} = E_\mu^{\text{rec}} + E_{\text{had}}^{\text{rec}}$$



Method Description*

Given the vector of variable X, Y with their mean and covariance matrix

$$\mu_{X,Y} = \begin{pmatrix} \mu_X \\ \mu_Y \end{pmatrix} \quad \Sigma = \begin{pmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{pmatrix}$$


We can calculate the conditional mean and conditional variance

$$\begin{aligned} \mu_{Y|X}^{\text{constrained}} &= \mu_Y + \Sigma_{YX} \Sigma_{XX}^{-1} (m_X - \mu_X) & \text{Expectations on } Y \\ \Sigma_{Y|X}^{\text{constrained}} &= \Sigma_{YY} - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{YX} & \text{Uncertainties on } Y \end{aligned}$$

*E. L. Morris "Multivariate Statistics: a Vector Space Approach" 1983

Muon Kinematics:

Total Reconstructed Energy of the Muon (E_{μ}^{rec})



FC: fully contained events in the fiducial volume

PC: partially contained events in the fiducial volume


Goodness-of-fit test:

$$\chi^2 = (m - \mu)^T \cdot \Sigma^{-1} \cdot (m - \mu)$$

Error band for statistic, flux and cross section systematic uncertainties

The reasonable value of goodness-of-fit test show good agreement between data and model prediction

Muon Kinematics: Reconstructed Polar Angle between of Muon and Neutrino Beam ($\cos\theta_{\mu}^{\text{rec}}$)



FC: fully contained events in the fiducial volume

PC: partially contained events in the fiducial volume

Goodness-of-fit test:


$$\chi^2 = (m - \mu)^T \cdot \Sigma^{-1} \cdot (m - \mu)$$

Error band for statistic, flux and cross section systematic uncertainties

The reasonable value of goodness-of-fit test show good agreement between data and model prediction

Validation of Hadronic Energy Reconstruction

MicroBooNE Preliminary




* Similar result of PC events

- **Before constraint: Excess observed at low hadronic energy**
 - Mis-modeling of missing energy in the hadron final states?
- **After constraint with E_{μ}^{rec} and $\cos\theta_{\mu}^{\text{rec}}$: No more excess at low hadronic energy**
 - Significant reduction in overall uncertainties (20% \rightarrow 5%)
 - No sign of mis-modeling of the hadron missing energy

Validation of Hadronic Energy Reconstruction

MicroBooNE Preliminary



* Similar result of PC events

- Before constraint: Excess observed at low hadronic energy
 - Mis-modeling of missing energy in the hadron final states?
- After constraint with E_{μ}^{rec} and $\cos\theta_{\mu}^{\text{rec}}$:
No more excess at low hadronic energy
 - Significant reduction in overall uncertainties (20% \rightarrow 5%)
 - No sign of mis-modeling of the **hadron missing energy**


$$\mu_{Y|X}^{\text{constrained}} = \mu_Y + \Sigma_{YX} \Sigma_{XX}^{-1} (m_X - \mu_X) \quad Y: E_{\text{had}}^{\text{rec}}$$

$$\Sigma_{Y|X}^{\text{constrained}} = \Sigma_{YY} - \Sigma_{YX} \Sigma_{XX}^{-1} \Sigma_{YX} \quad X: E_{\mu}^{\text{rec}} \text{ and } \cos\theta_{\mu}^{\text{rec}}$$

To measure the cross section


Reco neutrino energy

MicroBooNE Preliminary




Reco muon energy

MicroBooNE Preliminary




Reco polar angle

MicroBooNE Preliminary



Reco hadrons energy

MicroBooNE Preliminary



- Measure the cross section:

- σ_{CC}/E_ν : total cross section as a function of neutrino energy
- $d\sigma_{CC}/dE_\mu$: differential cross section as a function of muon energy
- $d\sigma_{CC}/d\nu$: differential cross section as a function of energy transfer to Ar

- It enables us to do the multi-dimension differential cross-section measurements

- E.g. $d\sigma_{CC}/dE_\mu d\theta_\mu$

Summary

- A high-performance inclusive ν_μ CC selection (**93% purity, 64% efficiency**) has been achieved using Wire-Cell reconstruction at MicroBooNE
- New technique with conditional constraint allows for more stringent validations of the **cross-section modeling** and **neutrino energy reconstruction** for oscillation and cross section measurements
 - Examination of hadronic energy distribution after constraint by muon kinematics explains the observed low-hadronic-energy excess
- High-statistics ν_μ CC event selection (**$\approx 225\text{k}$ expected for 1.2×10^{21} POT**) for precision cross-section measurements
 - Stay tuned for the initial measurement of cross section (by Wiener-SVD unfolding method [arXiv:1705.03568](https://arxiv.org/abs/1705.03568))

Thank you!