

Optimization

Wenxiao Yang*

*Department of Mathematics, University of Illinois at Urbana-Champaign

2022

目录

1	Unconstrained Optimization	3
1.1	Conditions for Optimality	3
1.2	Global minimizer, Local minimizer	3
1.3	Optimization in \mathbb{R}	3
1.3.1	Theorem: local minimizer $\Rightarrow f'(x^*) = 0$	3
1.3.2	Theorem: $f'(x^*) = 0, f''(x^*) \geq 0 \Rightarrow$ local minimizer	3
1.4	Optimization in \mathbb{R}^n	3
1.4.1	Necessary Conditions for Optimality: Local Extremum $\Rightarrow \nabla f(x^*) = 0$	3
1.4.2	Stationary Point, Saddle Point	4
1.4.3	Second Order Necessary Condition	5
1.4.4	Sufficient Conditions for Optimality	5
1.4.5	Using Optimality Conditions to Find Minimum	6
1.4.6	Fix Conditions for Global Optimality	6
1.5	Optimization in a Set	7
1.5.1	Existence of Global-min	7
1.6	Method of finding-global-min-among-stationary-points (FGMSP)	8
2	Convexity	8
2.1	Definition	8
2.2	Optimal Conditions	9
2.3	μ -strongly convexity	9
2.4	Unconstrained Quadratic Optimization	9
3	MATH 484	10
3.0.1	The first-derivative test in \mathbb{R}^n : $\phi'_u(t) = \nabla f(x + tu) \cdot u$	11
3.0.2	Theorem 4: ∇f is continuous, x^* is a global minimizer of $f \Rightarrow \nabla f(x^*) = 0$. . .	11

3.0.3	The second-derivative test in \mathbb{R}^n	11
3.0.4	Hessian matrix	11
3.0.5	Theorem 5: Hf is continuous, $\nabla f(x^*) = 0$, $u^T Hf(x^*)u \geq 0, \forall u \Rightarrow x^*$ is a global minimizer of f	12
3.1	Minimizing over other sets	12
3.1.1	Theorem 7: ∇f is continuous, x^* (interior of D) is a local minimizer of f $\Rightarrow \nabla f(x^*) = 0$	12
3.1.2	Theorem 8: Hf is continuous, x^* (interior of D) $\nabla f(x^*) = 0$, $\exists r$ s.t. $u^T Hf(x^*)u \geq$ $0, \forall x \in B(x^*, r), \forall u \Rightarrow x^*$ is a local minimizer of f	12

1 Unconstrained Optimization

1.1 Conditions for Optimality

Function: $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $x \in \mathcal{X}$, $\mathcal{X} \subseteq \mathbb{R}^n$.

Terminology: x^* will always be the optimal input at some function.

1.2 Global minimizer, Local minimizer

Definition 1.

Say x^* is a global minimizer(minimum) of f if $f(x^*) \leq f(x), \forall x \in \mathcal{X}$.

Say x^* is a unique global minimizer(minimum) of f if $f(x^*) < f(x), \forall x \neq x^*$.

Say x^* is a local minimizer(minimum) of f if $\exists r > 0$ so that $f(x^*) \leq f(x)$ when $\|x - x^*\| < r$.

A minimizer is strict if $f(x^*) < f(x)$ for all relevant x .

1.3 Optimization in \mathbb{R}

1.3.1 Theorem: local minimizer $\Rightarrow f'(x^*) = 0$

Theorem 1. If $f(x)$ is differentiable function and x^* is a local minimizer, then $f'(x^*) = 0$.

证明.

Def of $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h)-f(x)}{h}$

Def of local minimizer: $f(x^*) - f(x) \geq 0, |x^* - x| < r$

when $0 < h < r$, $\frac{f(x+h)-f(x)}{h} \geq 0$; when $-r < h < 0$, $\frac{f(x+h)-f(x)}{h} \leq 0$. Then $f'(x) = 0$. \square

1.3.2 Theorem: $f'(x^*) = 0, f''(x^*) \geq 0 \Rightarrow$ local minimizer

Theorem 2. If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a function with a continuous second derivative and x^* is a critical point of f (i.e. $f'(x) = 0$), then:

(1): If $f''(x) \geq 0, \forall x \in \mathbb{R}$, then x^* is a global minimizer on \mathbb{R} .

(2): If $f''(x) \geq 0, \forall x \in [a, b]$, then x^* is a global minimizer on $[a, b]$.

(3): If we only know $f''(x^*) \geq 0$, x^* is a local minimizer.

证明.

(1) $f(x) = f(x^*) + f'(x^*)(x - x^*) + \frac{1}{2}f''(\xi)(x - x^*)^2 = f(x^*) + 0 + \text{something non negative} \geq f(x^*) \forall x$

(2) Similar to (1)

(3) $f''(x^*) \geq 0, f''$ continuous $\Rightarrow \exists r$ s.t. $f''(x) \geq 0 \forall x \in [x^* - \frac{r}{2}, x^* + \frac{r}{2}]$, then x is a local minimizer. \square

1.4 Optimization in \mathbb{R}^n

1.4.1 Necessary Conditions for Optimality: Local Extremum $\Rightarrow \nabla f(x^*) = 0$

A base point x , we consider an arbitrary direction u . $\{x + tu | t \in \mathbb{R}\}$

For $\alpha > 0$ sufficiently small:

1. $f(x^*) \leq f(x^* + \alpha u)$
2. $g(\alpha) = f(x^* + \alpha u) - f(x^*) \geq 0$
3. $g(\beta)$ is continuously differentiable for $\beta \in [0, \alpha]$

By chain rule,

$$g'(\beta) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i$$

By Mean Value Theorem,

$$g(\alpha) = g(0) + g'(\beta)\alpha \text{ for some } \beta \in [0, \alpha]$$

Thus

$$\begin{aligned} g(\alpha) &= \alpha \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i \geq 0 \\ &\Rightarrow \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i \geq 0 \end{aligned}$$

Letting $\alpha \rightarrow 0$ and hence $\beta \rightarrow 0$, we get

$$\sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^*) u_i \geq 0 \text{ for all } u \in \mathbb{R}^n$$

By choosing $u = [1, 0, \dots, 0]^T$, $u = [-1, 0, \dots, 0]^T$, we get

$$\frac{\partial f(x^*)}{\partial x_1} \geq 0, \quad \frac{\partial f(x^*)}{\partial x_1} \leq 0 \Rightarrow \frac{\partial f(x^*)}{\partial x_1} = 0$$

Similarly, we can get

$$\nabla f(x^*) = \left[\frac{\partial f(x^*)}{\partial x_1}, \frac{\partial f(x^*)}{\partial x_2}, \dots, \frac{\partial f(x^*)}{\partial x_n} \right]^T = 0$$

Theorem 3. *If f is continuously differentiable and x^* is a local extremum. Then $\nabla f(x^*) = 0$.*

1.4.2 Stationary Point, Saddle Point

All points x^* s.t. $\nabla f(x^*) = 0$ are called stationary points.

Thus, all extrema are stationary points.

But not all stationary points have to be extrema.

Saddle points are the stationary points neither local minimum nor local maximum.

Example 1. $f(x) = x^3$, $x = 0$ is a stationary point but not extrema. (saddle point)

1.4.3 Second Order Necessary Condition

Definition 2. The Hessian of f at point x is an $n \times n$ symmetric matrix denoted by $\nabla^2 f(x)$ with $[\nabla^2 f(x)]_{ij} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}$

Theorem 4. Suppose f is twice continuously differentiable and x^* is local minimum. Then

$$\nabla f(x^*) = 0 \text{ and } \nabla^2 f(x^*) \geq 0$$

证明.

$\nabla f(x^*) = 0$ already proved before.

Let α be small enough so that $g(\alpha) = f(x^* + \alpha u) - f(x^*) \geq 0$.

By Taylor series expansion,

$$\begin{aligned} g(\alpha) &= g(0) + \alpha g'(0) + \frac{\alpha^2}{2} g''(0) + O(\alpha^2) \\ g'(\alpha) &= \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i = \nabla f(x^* + \alpha u)^T u \\ g''(\alpha) &= \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j}(x^* + \beta u) u_i u_j = u^T \nabla^2 f(x^* + \alpha u) u \end{aligned}$$

$$g'(0) = \nabla f(x^*)^T u = 0; \quad g''(0) = u^T \nabla^2 f(x^*) u$$

$$g(\alpha) = \frac{\alpha^2}{2} u^T \nabla^2 f(x^*) u + O(\alpha^2) \geq 0$$

$$\begin{aligned} \text{When } \alpha \rightarrow 0, \text{ we get } u^T \nabla^2 f(x^*) u &\geq 0, \quad \forall u \in \mathbb{R}^n \\ &\Rightarrow \nabla^2 f(x^*) \geq 0 \end{aligned}$$

□

1.4.4 Sufficient Conditions for Optimality

Theorem 5. Suppose f is twice continuously differentiable in a neighborhood of x^* and (1) $\nabla f(x^*) = 0$; (2) $\nabla^2 f(x^*)$ is PD ($u^T \nabla^2 f(x^*) u > 0, \forall u \in \mathbb{R}^n$). Then x^* is local minimum.

证明.

Consider $u \in \mathbb{R}^n, \alpha > 0$ and let

$$\begin{aligned} g(\alpha) &= f(x^* + \alpha u) - f(x^*) \\ &= \frac{\alpha^2}{2} u^T \nabla^2 f(x^*) u + O(\alpha^2) \geq 0 \\ &= \frac{\alpha^2}{2} [u^T \nabla^2 f(x^*) u + 2 \frac{O(\alpha^2)}{\alpha^2}] \\ &u^T \nabla^2 f(x^*) u > 0; \quad \frac{O(\alpha^2)}{\alpha^2} \rightarrow 0 \\ &\Rightarrow g(\alpha) > 0 \text{ for } \alpha \text{ sufficiently small for all } u \neq 0 \\ &\Rightarrow x^* \text{ is local minimum.} \end{aligned}$$

(specially if $\|u\| = 1$, $u^T \nabla^2 f(x^*) u \geq \lambda_{\min}(\nabla^2 f(x^*))$, $\lambda_{\min}(\nabla^2 f(x^*))$ is the minimal eigenvalues of $\nabla^2 f(x^*)$.) \square

1.4.5 Using Optimality Conditions to Find Minimum

1. Find all points satisfying necessary condition $\nabla f(x) = 0$ (all stationary points)
2. Filter out points that don't satisfy $\nabla^2 f(x) \geq 0$
3. Points with $\nabla^2 f(x) > 0$ are strict local minimum.
4. Among all points with $\nabla^2 f(x) \geq 0$, declare a global minimum, one with the smallest value of f , assuming that global minimum exists.

Example 2. $f(x) = 2x^2 - x^4$

$$f'(x) = 4x - 4x^3 = 0$$

$\Rightarrow x = 0, x = 1, x = -1$ are stationary points

$$f''(x) = 4 - 12x^2 = \begin{cases} 4 & \text{if } x = 0 \\ -8 & \text{if } x = 1, -1 \end{cases}$$

$\Rightarrow x = 0$ is the only local min, and it is strict

But $-f(x) \rightarrow \infty$ as $|x| \rightarrow \infty \Rightarrow$ no global min, but global max exists. $f(1), f(-1)$ are strict local max and both global max.

1.4.6 Fix Conditions for Global Optimality

Claim 1: Consider a differentiable function f . Suppose:

(C1) f has at least one global minimizer;

(C2) The set of stationary points is S , and $f(x^*) \leq f(x), \forall x \in S$.

Then x^* is a global minimizer of f .

证明.

Suppose \hat{x} is a global minimizer of f , i.e.,

$$f(\hat{x}) \leq f(x), \forall x.$$

By the necessary optimality condition, we have $\nabla f(\hat{x}) = 0$, thus $\hat{x} \in S$. By (C2), we have

$$f(x^*) \leq f(\hat{x}).$$

Combining the two inequalities, we have $f(\hat{x}) \leq f(x^*) \leq f(\hat{x})$, thus $f(\hat{x}) = f(x^*)$. Plugging into the second inequality, we have $f(x^*) \leq f(x), \forall x$. Thus x^* is a global minimizer of f . \square

1.5 Optimization in a Set

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & x \in X\end{array}$$

- Objective function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuous function

- Optimization variable $x \in X$

- Local minimum of f on X : $\exists \epsilon > 0$ s.t. $f(x) \geq f(\hat{x})$, for all $x \in X$ such that $\|x - \hat{x}\| \leq \epsilon$;

i.e., x^* is the best in the intersection of a small neighborhood and X

- Global minimum of f on X : $f(x) \geq f(x^*)$ for all $x \in X$

"Strict global minimum", "strict local minimum" "local maximum", "global maximum" of f on X are defined accordingly

1.5.1 Existence of Global-min

Theorem 6 (Bolzano-Weierstrass Theorem (compact domain)). *Any continuous function f has at least one global minimizer on any **compact set** X .*

That is, there exists an $x^ \in X$ such that $f(x) \geq f(x^*)$, $\forall x \in X$.*

Corollary 1 (bounded level sets). *Suppose $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuous function. If for a certain c , the level set*

$$\{x \mid f(x) \leq c\}$$

*is **non-empty** and **compact**, then the global minimizer of f exists, i.e., there exists $x^* \in \mathbb{R}^d$ s.t.*

$$f(x^*) = \inf_{x \in \mathbb{R}^d} f(x)$$

Example 3. $f(x) = x^2$. Level set $\{x \mid x^2 \leq 1\}$ is $\{x \mid -1 \leq x \leq 1\}$: non-empty compact. Thus there exists a global minimum.

Corollary 2 (coercive). *Suppose $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuous function. If $f(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$, then the global minimizer of f over \mathbb{R}^d exists.*

証明. Let $\alpha \in \mathbb{R}^d$ be chosen so that the set $S = \{x \mid f(x) \leq \alpha\}$ is non-empty. By coercivity, this set is compact. □

Coercive \Rightarrow one non-empty bounded level set; but not the other way.

Claim (all level sets bounded \Leftrightarrow coercive): Let f be a continuous function, then f is coercive iff $\{x \mid f(x) \leq \alpha\}$ is compact for any α .

1.6 Method of finding-global-min-among-stationary-points (FGMSP)

Method of finding-global-min-among-stationary-points (FGMSP):

Step 0: Verify coercive or bounded level set:

- Case 1: success, go to Step 1.
- Case 2: otherwise, try to show non-existence of global-min. If success, exit and report "no global-min exists".
- Case 3: cannot verify coercive or bounded level set; cannot show non-existence of global-min. Exit and report "cannot decide".

Step 1: Find all stationary points (candidates) by solving $\nabla f(\mathbf{x}) = 0$;

Step 2 (optional): Find all candidates s.t. $\nabla^2 f(\mathbf{x}) \succeq 0$.

Step 3: Among all candidates, find one candidate with the minimal value. Output this candidate, and report "find a global min".

2 Convexity

2.1 Definition

Convex set C : $x, y \in C$ implies $\lambda x + (1 - \lambda)y \in C$, for any $\lambda \in [0, 1]$.

Convex function (0-th order): f is convex in a convex set C iff $f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y), \forall x, y \in C, \forall \alpha \in [0, 1]$.

Property (1st order) If f is differentiable, then f is convex iff $f(z) \geq f(x) + (z - x)' \nabla f(x), \forall x, z \in C$.

Property (2nd order): If f is twice differentiable, then f is convex iff

$$\nabla^2 f(x) \succeq 0, \forall x \in C.$$

Strictly convex: when \geq becomes $>$ in any of the above relations.

A function f is a **concave function** iff $-f$ is a convex function.

Convex set graph:




图 1:

2.2 Optimal Conditions

Proposition 1. Let $f : X \mapsto \mathbb{R}$ be a convex function over the convex set X .

(a) A local-min of f over X is also a global-min over X .

(b) If X is open (e.g. \mathbb{R}^n), then $\nabla f(x^*) = 0$ is a necessary and sufficient condition for x^* to be a global minimum.

证明.

Proof based on a property: If f is differentiable over C , then f is convex iff

$$f(z) \geq f(x) + (z - x)' \nabla f(x), \quad \forall x, z \in C.$$

□

Corollary 3. Let $f : X \mapsto \mathbb{R}$ be a concave function over the convex set X .

(a) A local-max of f over X is also a global-max over X .

(b) If X is open (e.g. \mathbb{R}^n), then $\nabla f(x^*) = 0$ is a necessary and sufficient condition for x^* to be a global maximum.

2.3 μ -strongly convexity

Definition: We say $f : C \rightarrow \mathbb{R}$ is a μ -strongly convex function in a convex set C if f is differentiable and

$$\langle \nabla f(w) - \nabla f(v), w - v \rangle \geq \mu \|w - v\|^2, \quad \forall w, v \in C.$$

If f is twice differentiable, then f is μ -strongly convex iff

$$\nabla^2 f(x) \succeq \mu I, \quad \forall x \in C.$$

Namely, all eigenvalues of the Hessian at any point is at least μ .

if $f(w)$ is convex, then $f(w) + \frac{\mu}{2} \|w\|^2$ is μ -strongly convex.

- In machine learning, easy to change a convex function to a strongly convex function: just add a regularizer

2.4 Unconstrained Quadratic Optimization

$$\begin{aligned} \text{minimize} \quad & f(\mathbf{w}) = \frac{1}{2} \mathbf{w}^T \mathbf{Q} \mathbf{w} - \mathbf{b}^T \mathbf{w} \\ \text{subject to} \quad & \mathbf{w} \in \mathbb{R}^d \end{aligned}$$

where \mathbf{Q} is a symmetric $d \times d$ matrix. (what if non-symmetric?)

- Necessary condition for (local) optimality

$$\mathbf{Q} \mathbf{w} = \mathbf{b}, \quad \mathbf{Q} \succeq 0$$

Case 1: $\mathbf{Q}\mathbf{w} = \mathbf{b}$ has no solution, i.e. $\mathbf{b} \notin R(\mathbf{Q})$. No stationary point, no lower bound (f can achieve $-\infty$).

Case 2: \mathbf{Q} is not PSD (f is non-convex) No local-min, no lower bound (f can achieve $-\infty$).

Case 3: $\mathbf{Q} \succeq 0$ (PSD) and $\mathbf{b} \in R(\mathbf{Q})$. Convex, has global-min, any stationary point is a global optimal solution.

Example 4. *Toy Problem 1:* $\min_{x,y \in \mathbb{R}} f(x,y) \triangleq x^2 + y^2 + \alpha xy$.

1. Step 1: First order condition: $2x^* + \alpha y^* = 0, 2y^* + \alpha x^* = 0$.

- We get $4x^* = -2\alpha y^* = \alpha^2 x^*$. So $(4 - \alpha^2) x^* = 0$.

- Case 1: $\alpha^2 = 4$. If $x^* = -\alpha y^*/2$, then (x^*, y^*) is a stationary point.

- Case 2: $\alpha^2 \neq 4$. Then $x^* = 0; y^* = -\alpha x^*/2 = 0$. So $(0, 0)$ is stat-pt.

2. Step 2: Check convexity. Hessian $\nabla^2 f(x,y) = \begin{pmatrix} 2 & \alpha \\ \alpha & 2 \end{pmatrix}$.

Eigenvalues λ_1, λ_2 satisfy $(\lambda_i - 2)^2 = \alpha^2, i = 1, 2$. Thus $\lambda_{1,2} = 2 \pm |\alpha|$.

- If $|\alpha| \leq 2$, then $\lambda_i \geq 0, \forall i$. Thus f is convex. Any stat-pt is global-min.

- If $|\alpha| > 2$, at least one $\lambda_i < 0$, thus f is not convex.

3. Step 3: For non-convex case ($|\alpha| > 2$), prove no lower bound. $f(x,y) = (x + \alpha y/2) + (1 - \alpha^2/4) y^2$. Pick $y = M, x = -\alpha M/2$, then $f(x,y) = (1 - \alpha^2/4) M^2 \rightarrow -\infty$ as $M \rightarrow \infty$.

Summary: If $|\alpha| > 2$, no global-min, $(0,0)$ is stat-pt; if $|\alpha| = 2$, any $(-0.5\alpha t, t), t \in \mathbb{R}$ is a stat-pt and global-min; if $|\alpha| < 2$, $(0,0)$ is the unique stat-pt and global-min.

3 MATH 484

A base point x , we consider an arbitrary direction u . $\{x + tu | t \in \mathbb{R}\}$

We define the restriction of f to the line through x in the direction of u to be the function:

$$\phi_u(t) = f(x + tu)$$

Lemma 1. x^* is a global minimizer of f iff for all u , $t = 0$ is the global minimizer of $\phi_u(t)$

证明.

$(\Rightarrow) \phi_u(0) = f(x^*) \leq f(x^* + tu) = \phi_u(t)$

(\Leftarrow) Let $X \in \mathbb{R}^n, u = X - x^*$. $\phi_u(0) \leq \phi_u(1) \Rightarrow f(x^*) \leq f(x^* + u) = f(x)$

□

3.0.1 The first-derivative test in \mathbb{R}^n : $\phi'_u(t) = \nabla f(x + tu) \cdot u$

First derivative of $f : \mathbb{R}^n \rightarrow \mathbb{R}$, Easier: $\phi'_u(t)$?

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}^n$:

$$\frac{\partial f(g(t))}{\partial t} = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(g(t)) \frac{d}{dt} g_i(t)$$

$$\frac{\partial \phi_u(t)}{\partial t} = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x + tu) u_i$$

The gradient of f : $\nabla f(x) = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_d})^T \Rightarrow \phi'_u(t) = \nabla f(x + tu) \cdot u$

Fine print: Chain rule only works when all $\frac{\partial f}{\partial x_k}$ exists and are continuous.

Example 5. $f(x, y) = x^2 + 3xy - 1$, $x^* = (0, 0)$, $u = (3, 2)$

$$\phi_u(t) = f(x^* + tu) = f(3t, 2t) = 27t^2 - 1$$

$$\phi'_u(t) = 54t$$

$$\nabla f(x, y) = (2x + 3y, 3x)$$

$$\phi'_u(t) = \nabla f(x + tu) \cdot u = 54t$$

3.0.2 Theorem 4: ∇f is continuous, x^* is a global minimizer of $f \Rightarrow \nabla f(x^*) = 0$

Theorem 7 (Theorem 2.1). *Given a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, if ∇f is continuous and x^* is a global minimizer of f , then $\nabla f(x^*) = 0$. (When $\nabla f(x^*) = 0$, we call x^* a critical point of f .)*

x^* is a global minimizer $\Rightarrow x^*$ is a critical point, inverse may not true.

3.0.3 The second-derivative test in \mathbb{R}^n

$$\phi'_u(t) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x + tu) u_i$$

$$\phi''_u(t) = \sum_{i=1}^n \sum_{j=1}^n u_i u_j \frac{\partial^2 f}{\partial x_i \partial x_j}(x + tu)$$

3.0.4 Hessian matrix

Define Hessian matrix of f and write Hf . That is,

$$\phi''_u(t) = u^T Hf(x + tu) u$$

Fine print: Chain rule only works when all $\frac{\partial^2 f}{\partial x_i \partial x_j}$ exists and are continuous. ($\Rightarrow Hf$ is continuous)

3.0.5 Theorem 5: Hf is continuous, $\nabla f(x^*) = 0$, $u^T Hf(x^*)u \geq 0, \forall u \Rightarrow x^*$ is a global minimizer of f

Theorem 8. Given a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, if Hf is continuous and x^* is a critical point of f . If for any u , that $u^T Hf(x^*)u \geq 0$. Then x^* is a global minimizer of f .

proved by Taylor

Theorem 9 (Taylor). Given a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, if Hf is continuous and x^* is a critical point of f , then

$$f(x) = f(x^*) = \nabla f(x^*)(x - x^*) + \frac{1}{2}(x - x^*)^T Hf(z)(x - x^*)$$

for some z on the line between x and x^*

3.1 Minimizing over other sets

What if the domain of f : $D \subset \mathbb{R}^n$

(1): want x^* to be in the interior of D , not on the boundary (want to be able to "look" from x^* in any direction.)

(2): want x^* to "see" all other points in D using straight line u .

Convexity

good domain e.g. Ball: $B(x^*, r) = \{x \mid \|x - x^*\| < r\}$

3.1.1 Theorem 7: ∇f is continuous, x^* (interior of D) is a local minimizer of $f \Rightarrow \nabla f(x^*) = 0$

Theorem 10 (Theorem 4.1, 类似 Theorem 2.1). Suppose $f : D \rightarrow \mathbb{R}$ has continuous ∇f and x^* is not on the boundary of D . If x^* is a local minimizer of f , then x^* is a critical point of f : $\nabla f(x^*) = 0$

3.1.2 Theorem 8: Hf is continuous, x^* (interior of D) $\nabla f(x^*) = 0$, $\exists r$ s.t. $u^T Hf(x^*)u \geq 0, \forall x \in B(x^*, r), \forall u \Rightarrow x^*$ is a local minimizer of f

Theorem 11. Given a function $f : D \rightarrow \mathbb{R}$, if Hf is continuous and x^* is a critical point of f in the interior of D . Suppose $\exists r$ s.t. for any u , that $u^T Hf(x^*)u \geq 0$ whenever $x \in B(x^*, r) \subset D$. Then x^* is a local minimizer of f .