

# Optimization

Wenxiao Yang\*

\*Department of Mathematics, University of Illinois at Urbana-Champaign

2022

## 目录

<b>1</b>	<b>Unconstrained Optimization</b>	<b>2</b>
1.1	Conditions for Optimality . . . . .	2
1.2	Global minimizer, Local minimizer . . . . .	2
1.3	Optimization in $\mathbb{R}$ . . . . .	2
1.3.1	Theorem 1: differentiable $f$ , $x^*$ is a local minimizer $\Rightarrow f'(x^*) = 0$ . . . . .	2
1.3.2	Theorem 2: $f'(x^*) = 0, f''(x^*) \geq 0, \forall x \in [a, b] \Rightarrow x^*$ is a global minimizer on $[a, b]$ ; $f'(x^*) = 0, f''(x^*) \geq 0 \Rightarrow x^*$ is a local minimizer . . . . .	2
1.4	Optimization in $\mathbb{R}^n$ . . . . .	3
1.4.1	Necessary Conditions for Optimality: Local Minimum $\Rightarrow \nabla f(x^*) = 0$ . . . . .	3
1.4.2	The first-derivative test in $\mathbb{R}^n$ : $\phi'_u(t) = \nabla f(x + tu) \cdot u$ . . . . .	4
1.4.3	Theorem 4: $\nabla f$ is continuous, $x^*$ is a global minimizer of $f \Rightarrow \nabla f(x^*) = 0$ . . . . .	4
1.4.4	The second-derivative test in $\mathbb{R}^n$ . . . . .	4
1.4.5	Hessian matrix . . . . .	4
1.4.6	Theorem 5: $Hf$ is continuous, $\nabla f(x^*) = 0, u^T Hf(x^*)u \geq 0, \forall u \Rightarrow x^*$ is a global minimizer of $f$ . . . . .	5
1.5	Minimizing over other sets . . . . .	5
1.5.1	Theorem 7: $\nabla f$ is continuous, $x^*$ (interior of $D$ ) is a local minimizer of $f \Rightarrow \nabla f(x^*) = 0$ . . . . .	5
1.5.2	Theorem 8: $Hf$ is continuous, $x^*$ (interior of $D$ ) $\nabla f(x^*) = 0, \exists r$ s.t. $u^T Hf(x^*)u \geq 0, \forall x \in B(x^*, r), \forall u \Rightarrow x^*$ is a local minimizer of $f$ . . . . .	5

# 1 Unconstrained Optimization

## 1.1 Conditions for Optimality

Function:  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ,  $x \in \mathcal{X}$ ,  $\mathcal{X} \subseteq \mathbb{R}^n$ .

Terminology:  $x^*$  will always be the optimal input at some function.

## 1.2 Global minimizer, Local minimizer

**Definition 1.**

Say  $x^*$  is a global minimizer(minimum) of  $f$  if  $f(x^*) \leq f(x), \forall x \in \mathcal{X}$ .

Say  $x^*$  is a unique global minimizer(minimum) of  $f$  if  $f(x^*) < f(x), \forall x \neq x^*$ .

Say  $x^*$  is a local minimizer(minimum) of  $f$  if  $\exists r > 0$  so that  $f(x^*) \leq f(x)$  when  $\|x - x^*\| < r$ .

A minimizer is strict if  $f(x^*) < f(x)$  for all relevant  $x$ .

## 1.3 Optimization in $\mathbb{R}$

**1.3.1 Theorem 1: differentiable  $f$ ,  $x^*$  is a local minimizer  $\Rightarrow f'(x^*) = 0$**

**Theorem 1.** If  $f(x)$  is differentiable function and  $x^*$  is a local minimizer, then  $f'(x^*) = 0$ .

证明.

Def of  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

Def of local minimizer:  $f(x^*) - f(x) \geq 0, |x^* - x| < r$

when  $0 < h < r$ ,  $\frac{f(x+h) - f(x)}{h} \geq 0$ ; when  $-r < h < 0$ ,  $\frac{f(x+h) - f(x)}{h} \leq 0$ . Then  $f'(x) = 0$ .  $\square$

**1.3.2 Theorem 2:  $f'(x^*) = 0, f''(x^*) \geq 0, \forall x \in [a, b] \Rightarrow x^*$  is a global minimizer on  $[a, b]$ ;  
 $f'(x^*) = 0, f''(x^*) \geq 0 \Rightarrow x^*$  is a local minimizer**

**Theorem 2.** If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a function with a continuous second derivative and  $x^*$  is a critical point of  $f$  (i.e.  $f'(x) = 0$ ), then:

(1): If  $f''(x) \geq 0, \forall x \in \mathbb{R}$ , then  $x^*$  is a global minimizer on  $\mathbb{R}$ .

(2): If  $f''(x) \geq 0, \forall x \in [a, b]$ , then  $x^*$  is a global minimizer on  $[a, b]$ .

(3): If we only know  $f''(x^*) \geq 0$ ,  $x^*$  is a local minimizer.

proof of theorem 2.

(1)  $f(x) = f(x^*) + f'(x^*)(x - x^*) + \frac{1}{2}f''(\xi)(x - x^*)^2 = f(x^*) + 0 + \text{something non negative} \geq f(x^*) \forall x$

(2) Similar to (1)

(3)  $f''(x^*) \geq 0, f''$  continuous  $\Rightarrow \exists r$  s.t.  $f''(x) \geq 0 \forall x \in [x^* - \frac{r}{2}, x^* + \frac{r}{2}]$ , then  $x$  is a local minimizer.  $\square$

## 1.4 Optimization in $\mathbb{R}^n$

### 1.4.1 Necessary Conditions for Optimality: Local Minimum $\Rightarrow \nabla f(x^*) = 0$

A base point  $x$ , we consider an arbitrary direction  $u$ .  $\{x + tu | t \in \mathbb{R}\}$

For  $\alpha > 0$  sufficiently small:

1.  $f(x^*) \leq f(x^* + \alpha u)$
2.  $g(\alpha) = f(x^* + \alpha u) - f(x^*) \geq 0$
3.  $g(\beta)$  is continuously differentiable for  $\beta \in [0, \alpha]$

By chain rule,

$$g'(\beta) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i$$

By Mean Value Theorem,

$$g(\alpha) = g(0) + g'(\beta)\alpha \text{ for some } \beta \in [0, \alpha]$$

Thus

$$\begin{aligned} g(\alpha) &= \alpha \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i \geq 0 \\ &\Rightarrow \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^* + \beta u) u_i \geq 0 \end{aligned}$$

Letting  $\alpha \rightarrow 0$  and hence  $\beta \rightarrow 0$ , we get

$$\sum_{i=1}^n \frac{\partial f}{\partial x_i}(x^*) u_i \geq 0 \text{ for all } u \in \mathbb{R}^n$$

By choosing  $u = [1, 0, \dots, 0]^T$ ,  $u = [-1, 0, \dots, 0]^T$ , we get

$$\frac{\partial f(x^*)}{\partial x_1} \geq 0, \quad \frac{\partial f(x^*)}{\partial x_1} \leq 0 \Rightarrow \frac{\partial f(x^*)}{\partial x_1} = 0$$

Similarly, we can get

$$\nabla f(x^*) = \left[ \frac{\partial f(x^*)}{\partial x_1}, \frac{\partial f(x^*)}{\partial x_2}, \dots, \frac{\partial f(x^*)}{\partial x_n} \right]^T = 0$$

A base point  $x$ , we consider an arbitrary direction  $u$ .  $\{x + tu | t \in \mathbb{R}\}$

We define the restriction of  $f$  to the line through  $x$  in the direction of  $u$  to be the function:

$$\phi_u(t) = f(x + tu)$$

**Lemma 1.**  $x^*$  is a global minimizer of  $f$  iff for all  $u$ ,  $t = 0$  is the global minimizer of  $\phi_u(t)$

证明.

$$(\Rightarrow) \phi_u(0) = f(x^*) \leq f(x^* + tu) = \phi_u(t)$$

$$(\Leftarrow) \text{ Let } X \in \mathbb{R}^n, u = X - x^*. \phi_u(0) \leq \phi_u(1) \Rightarrow f(x^*) \leq f(x^* + u) = f(x)$$

□

### 1.4.2 The first-derivative test in $\mathbb{R}^n$ : $\phi'_u(t) = \nabla f(x + tu) \cdot u$

First derivative of  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , Easier:  $\phi'_u(t)$ ?

Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}^n$ :

$$\frac{\partial f(g(t))}{\partial t} = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(g(t)) \frac{d}{dt} g_i(t)$$

$$\frac{\partial \phi_u(t)}{\partial t} = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x + tu) u_i$$

The gradient of  $f$ :  $\nabla f(x) = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_d})^T \Rightarrow \phi'_u(t) = \nabla f(x + tu) \cdot u$

Fine print: Chain rule only works when all  $\frac{\partial f}{\partial x_k}$  exists and are continuous.

**Example 1.**  $f(x, y) = x^2 + 3xy - 1$ ,  $x^* = (0, 0)$ ,  $u = (3, 2)$

$$\phi_u(t) = f(x^* + tu) = f(3t, 2t) = 27t^2 - 1$$

$$\phi'_u(t) = 54t$$

$$\nabla f(x, y) = (2x + 3y, 3x)$$

$$\phi'_u(t) = \nabla f(x + tu) \cdot u = 54t$$

### 1.4.3 Theorem 4: $\nabla f$ is continuous, $x^*$ is a global minimizer of $f \Rightarrow \nabla f(x^*) = 0$

**Theorem 3** (Theorem 2.1). *Given a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , if  $\nabla f$  is continuous and  $x^*$  is a global minimizer of  $f$ , then  $\nabla f(x^*) = 0$ . (When  $\nabla f(x^*) = 0$ , we call  $x^*$  a critical point of  $f$ .)*

$x^*$  is a global minimizer  $\Rightarrow x^*$  is a critical point, inverse may not true.

### 1.4.4 The second-derivative test in $\mathbb{R}^n$

$$\phi'_u(t) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(x + tu) u_i$$

$$\phi''_u(t) = \sum_{i=1}^n \sum_{j=1}^n u_i u_j \frac{\partial^2 f}{\partial x_i \partial x_j}(x + tu)$$

### 1.4.5 Hessian matrix

Define Hessian matrix of  $f$  and write  $Hf$ . That is,

$$\phi''_u(t) = u^T Hf(x + tu) u$$

Fine print: Chain rule only works when all  $\frac{\partial^2 f}{\partial x_i \partial x_j}$  exists and are continuous. ( $\Rightarrow Hf$  is continuous)

**1.4.6 Theorem 5:**  $Hf$  is continuous,  $\nabla f(x^*) = 0$ ,  $u^T Hf(x^*)u \geq 0, \forall u \Rightarrow x^*$  is a global minimizer of  $f$

**Theorem 4.** Given a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , if  $Hf$  is continuous and  $x^*$  is a critical point of  $f$ . If for any  $u$ , that  $u^T Hf(x^*)u \geq 0$ . Then  $x^*$  is a global minimizer of  $f$ .

proved by Taylor

**Theorem 5** (Taylor). Given a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , if  $Hf$  is continuous and  $x^*$  is a critical point of  $f$ , then

$$f(x) = f(x^*) = \nabla f(x^*)(x - x^*) + \frac{1}{2}(x - x^*)^T Hf(z)(x - x^*)$$

for some  $z$  on the line between  $x$  and  $x^*$

## 1.5 Minimizing over other sets

What if the domain of  $f$ :  $D \subset \mathbb{R}^n$

(1): want  $x^*$  to be in the interior of  $D$ , not on the boundary (want to be able to "look" from  $x^*$  in any direction.)

(2): want  $x^*$  to "see" all other points in  $D$  using straight line  $u$ .

Convexity

good domain e.g. Ball:  $B(x^*, r) = \{x \mid \|x - x^*\| < r\}$

**1.5.1 Theorem 7:**  $\nabla f$  is continuous,  $x^*$ (interior of  $D$ ) is a local minimizer of  $f \Rightarrow \nabla f(x^*) = 0$

**Theorem 6** (Theorem 4.1, 类似 Theorem 2.1). Suppose  $f : D \rightarrow \mathbb{R}$  has continuous  $\nabla f$  and  $x^*$  is not on the boundary of  $D$ . If  $x^*$  is a local minimizer of  $f$ , then  $x^*$  is a critical point of  $f$ :  $\nabla f(x^*) = 0$

**1.5.2 Theorem 8:**  $Hf$  is continuous,  $x^*$ (interior of  $D$ )  $\nabla f(x^*) = 0$ ,  $\exists r$  s.t.  $u^T Hf(x^*)u \geq 0, \forall x \in B(x^*, r), \forall u \Rightarrow x^*$  is a local minimizer of  $f$

**Theorem 7.** Given a function  $f : D \rightarrow \mathbb{R}$ , if  $Hf$  is continuous and  $x^*$  is a critical point of  $f$  in the interior of  $D$ . Suppose  $\exists r$  s.t. for any  $u$ , that  $u^T Hf(x^*)u \geq 0$  whenever  $x \in B(x^*, r) \subset D$ . Then  $x^*$  is a local minimizer of  $f$ .