



Inefficiency of Equilibria

- Prisoner's dilemma
 - Equilibrium outcome are bad for both the players
- We want to quantify how bad is an equilibrium solution than a centralized/coordinated/dictatorial solution

Inefficiency of Equilibria

- Prisoner's dilemma
 - Equilibrium outcome are bad for both the players
- We want to quantify how bad is an equilibrium solution than a centralized/coordinated/dictatorial solution
- Objective
 - What is the quantity we want to maximize/minimize?
 - What is the meaning of optimal solution?
- Notion of equilibria (Nash (pure, mixed), correlated, ...)

Inefficiency of Equilibria

- Prisoner's dilemma
 - Equilibrium outcome are bad for both the players
- We want to quantify how bad is an equilibrium solution than a centralized/coordinated/dictatorial solution
- Objective
 - What is the quantity we want to maximize/minimize?
 - What is the meaning of optimal solution?
- Notion of equilibria (Nash (pure, mixed), correlated, ...)
- Choice of objective function and choice of equilibrium concept



How to measure the quality of solution?

- **Utilitarian:** maximum social welfare (total utility/payoff)

How to measure the quality of solution?

- Utilitarian: maximum social welfare (total utility/payoff)
- Egalitarian: maximizes the minimum payoff (max min)

Why we want to measure inefficiency of equilibria?

The answer to this may help

- We may take some measure to improve it
- If it is not too bad, then we may not bother for the selfish behavior because it is expensive and sometimes impossible to change it

Why we want to measure inefficiency of equilibria?

The answer to this may help

- We may take some measure to improve it
- If it is not too bad, then we may not bother for the selfish behavior because it is expensive and sometimes impossible to change it
- How to quantify the selfish behavior (equilibrium)?
 - When there are multiple equilibria, which one to choose?

Price of Anarchy (PoA)

- PoA is defined as the ratio between the value of optimal solution and the value of the **worst** equilibrium solution

Price of Anarchy (PoA)

- PoA is defined as the ratio between the value of optimal solution and the value of the **worst** equilibrium solution

$$\text{PoA} = \frac{\text{value of optimal solution}}{\text{value of worst equilibrium}} \geq 1 \quad (\text{payoff/utility})$$

Price of Anarchy (PoA)

- PoA is defined as the ratio between the value of optimal solution and the value of the **worst** equilibrium solution

$$\text{PoA} = \frac{\text{value of optimal solution}}{\text{value of worst equilibrium}} \geq 1 \quad (\text{payoff/utility})$$

$$\text{PoA} = \frac{\text{worst equilibrium cost}}{\text{optimal cost}} \geq 1 \quad (\text{cost})$$

Keep it always greater than or equal to 1.

Price of Anarchy (PoA)

- PoA is defined as the ratio between the value of optimal solution and the value of the **worst** equilibrium solution

$$\text{PoA} = \frac{\text{value of optimal solution}}{\text{value of worst equilibrium}} \geq 1 \quad (\text{payoff/utility})$$

$$\text{PoA} = \frac{\text{worst equilibrium cost}}{\text{optimal cost}} \geq 1 \quad C \quad (\text{cost})$$

- If PoA is close to 1, then ?

$$S \begin{bmatrix} 0, 0 & (5, 1) \\ (1, 5) & 2, 2 \end{bmatrix}$$

$$NE = \left\{ (C, S), (S, C), \left(\left(\frac{3}{4}S, \frac{1}{4}C \right), \left(\frac{3}{4}S, \frac{1}{4}C \right) \right) \right\}$$

(Utilitarian)

$$PoA = \frac{6}{\frac{3}{16} \times 8 \times 2 + \frac{1}{16} \times 4} = \frac{12}{5}$$

Price of Stability (PoS)

$$PoS = \frac{6}{6} = 1$$

- PoS is defined as the ratio between the value of optimal solution and the value of the **best** equilibrium solution

$$PoS = \frac{\text{value of optimal solution}}{\text{value of best equilibrium}} \geq 1 \quad (\text{payoff/utility})$$

$$PoS = \frac{\text{best equilibrium cost}}{\text{optimal cost}} \geq 1 \quad (\text{cost})$$


- PoS quantifies the necessary degradation of solution quality due to imposition of the game theoretic constraint of stability
- $1 \leq PoS \leq PoA$



Example

Selfish Routing

- **Applications:** road traffic, communication networks, etc.
- Recall Braess' paradox



Selfish Routing

- **Applications:** road traffic, communication networks, etc.

- Recall Braess' paradox

☐ Atomic setting


☐ Non-atomic setting (each individual has negligible size)

each player is not negligible

Millions of people.

100 people.

$$P_{0A} = P_{0S} = \frac{2}{1.5} = \frac{4}{3}$$




$$\frac{x}{100} \leq 1 \Rightarrow 2 \text{ hours each,}$$

Pigou's Example (1920)

Non-atomic setting. (total 1 unit flow).

- A simple selfish routing network

- ☐ Two paths from s to t with delay (cost) of 1 and x



$$PoA = PoS = \frac{4}{3}$$

Equilibrium solution: \Rightarrow $x \leq 1$. All choose " x ". $\Rightarrow \underline{x = 1}$.


Optimal solution: $\min_x \{x^2 + (1-x)\}$
 $\Rightarrow x = \frac{1}{2}$

half choose "1" half choose " x ".

Pigou's Example (1920)

- A simple selfish routing network

- What happens when the paths have affine cost function $ax + b$ where $a, b \geq 0$ and r units of flow need to go from s to t



$$PoA = PoS = \frac{ar^2 + br}{\frac{3}{4}ar^2 + br} \leq \frac{4}{3}$$

Equilibrium sol: $ax+b \leq ar+b \Rightarrow$ all choose "ax+b"
 $\Rightarrow x = r$.


optimal sol: $\min_x \{(ar+b)(1-x) + x(ax+b)\}$
 $\Rightarrow x = \frac{r}{2} \Rightarrow$ half "ar+b"
 half "ax+b".

PoA of General Network

- Let $G = (V, E)$ be network and we have affine cost function on edges, i.e., $c_e(x) = a_e x + b_e$
- Let $f = \{f_e\}_{e \in E}$ be an equilibrium flow

$$\sum_{e \in E} f = \text{total flow}$$

1 or r
in former
examples.



Cost of equilibrium: $C(f) = \sum_{e \in E} (a_e f_e + b_e) f_e$.

Cost of optimal: $C(f^*)$.


PoA of General Network

- Let $G = (V, E)$ be network and we have affine cost function on edges, i.e., $c_e(x) = a_e x + b_e$
- Let $f = \{f_e\}_{e \in E}$ be an equilibrium flow
- Let $C(f)$ be the cost of f
- $C(f) = ?$

PoA of General Network

- Let $G = (V, E)$ be network and we have affine cost function on edges, i.e., $c_e(x) = a_e x + b_e$
 - Let $f = \{f_e\}_{e \in E}$ be an equilibrium flow
 - Let $C(f)$ be the cost of f
 - $C(f) = ?$
-
- Let $f^* = \{f_e^*\}_{e \in E}$ be an optimal solution
-
- PoA = ? $\frac{C(f)}{C(f^*)} \leq \frac{4}{3}$

Equilibrium: $f = \{f_e\}_{e \in E}$ total cost of each path.



$$C_{P_1}(f) = \sum_{e \in P_1} (a_e f_e + b_e)$$

$a_e f_e + b_e$
可以表示任一段的 cost.

$$C_{P_2}(f) = \sum_{e \in P_2} (a_e f_e + b_e)$$

$$C_{P_3}(f) = \sum_{e \in P_3} (a_e f_e + b_e)$$

① if there are positive amount of flow both on P_1 and P_2 , then $C_{P_1}(f) = C_{P_2}(f)$

otherwise one would switch from its path to the path with the lowest cost until they are same.

② if positive flow on P_1 , zero flow on P_2 , then $C_{P_1}(f) \leq C_{P_2}(f)$.

otherwise one would choose P_2 instead of P_1 .

$$P_0 A = \frac{C(f)}{C(f^*)}.$$

Prove $PoA \leq \frac{4}{3}$

Proof: Set Equilibrium flow $f = \{f_e\}_{e \in E}$
Let $X = \{x_e\}_{e \in E}$ be an arbitrary flow.

$$\begin{aligned} C^f(x) &= \sum_{e \in E} (a_e f_e + b_e) x_e \\ &= \sum (a_e f_e x_e + b_e x_e). \end{aligned}$$

$$f_e x_e - x_e^2 \leq \frac{1}{4} f_e^2$$

$$\leq \sum (a_e x_e^2 + b_e x_e) + \sum \frac{a_e}{4} f_e^2$$

$$\leq C(x) + \frac{1}{4} C(f)$$

In equilibrium, the flow only lies on smallest cost paths, hence $C^f(x) \begin{cases} = C^f(f) & \text{still only lies on} \\ & \text{smallest cost paths} \\ > C^f(f) & \text{some flow lie on} \\ & \text{other paths.} \end{cases}$
Hence,

$$\underline{C^f(x) \geq C^f(f) \quad \forall x.}$$


$$\Rightarrow C(f^*) + \frac{1}{4} C(f) \geq C^f(f^*) \geq C^f(f) = C(f)$$

$$\Rightarrow C(f^*) \geq \frac{3}{4} C(f)$$

$$\Rightarrow PoA \leq \frac{4}{3}.$$

Atomic Setting

- Few number of players, so the size is not negligible
- **Example:** Two players want to send 1 unit of flow from s to t in the Pigou-like network



$$\text{Equilibrium} \left\{ \begin{array}{ccc} x_1 & x_2 & \text{cost.} \\ (0, 2) & 4 \\ (1, 1) & 3 \end{array} \right.$$

$$PoA = \frac{\text{worst equilibrium}}{\text{optimal}} = \frac{4}{3}$$

$$PoS = \frac{\text{best equilibrium}}{\text{optimal}} = 1.$$

Atomic Setting

- Few number of players, so the size is not negligible
- **Example:** Two players want to send 1 unit of flow from s to t in the Pigou-like network
- Equilibrium?
- Difference with non-atomic setting?

{ ①. the cost of every player can be different.
②. can have multiple equilibriums

Atomic Setting

- Few number of players, so the size is not negligible
- **Example:** Two players want to send 1 unit of flow from s to t in the Pigou-like network
- Equilibrium?


Atomic Setting

- **General problem:** k players, each player i wants to send 1 unit of flow from s_i to t_i on a single path in a network G

different players have different
starts and terminals.

PoA

- PoA can be larger than $4/3$



- PoA = $5/2$ optimal: —
- worst equilibrium: —

$$P_0 A = \frac{10}{4} = \frac{5}{2}.$$