

Problem 1.

$$i) m_x(t) = E[e^{tx}] = \sum_{k=1}^n e^{tk} \cdot \frac{1}{n} = \frac{1}{n} \sum_{k=1}^n e^{tk}$$

$$ii) E[X] = \left. \frac{dm_x(t)}{dt} \right|_{t=0} = \frac{1}{n} \sum_{k=1}^n k e^{tk} \Big|_{t=0} = \frac{1}{n} \sum_{k=1}^n k$$

$$\text{Var}[X] = E[X^2] - E[X]^2$$

$$E[X^2] = \left. \frac{d^2 m_x(t)}{dt^2} \right|_{t=0} = \frac{1}{n} \sum_{k=1}^n k^2 e^{tk} \Big|_{t=0} = \frac{1}{n} \sum_{k=1}^n k^2$$

$$\Rightarrow \text{Var}[X] = \frac{1}{n} \sum_{k=1}^n k^2 - \left(\frac{1}{n} \sum_{k=1}^n k \right)^2$$

$$= \frac{1}{n} \sum_{k=1}^n k^2 - \frac{1}{n^2} \left(\sum_{k=1}^n k \right)^2$$

Problem 2.

$$m_x(t) = m_Y(t) \Rightarrow \sum_{k=0}^n e^{kt} f_X(k) = \sum_{k=0}^n e^{kt} f_Y(k) \Rightarrow \sum_{k=0}^n e^{kt} [f_X(k) - f_Y(k)] = 0$$

since for $\forall t \in (-\epsilon, \epsilon)$, $m_X(t) = m_Y(t)$

$$\Rightarrow \text{for } \forall e^t \in (e^{-\epsilon}, e^{\epsilon}), \sum_{k=0}^n (e^t)^k [f_X(k) - f_Y(k)] = 0$$

$$\Rightarrow \text{the only solution should be all the coefficients} = 0 \Rightarrow f_X(k) = f_Y(k) \text{ for } \forall 0 \leq k \leq n$$

$$\Rightarrow f_X(k) = f_Y(k) \text{ for } k = 0, \dots, n.$$

Problem 3.

$$i) P[Z=z] = \sum_{x+y=z} P[X=x \text{ and } Y=y] \text{ since } Z = X+Y$$

$$= \sum_{x+y=z} P[X=x] \cdot P[Y=y] \text{ since } X \text{ and } Y \text{ are independent.}$$

ii) Suppose X and Y are two independent and identical geometric random variables

$$\text{s.t. } X: S \rightarrow \mathcal{N} \setminus \{0\}, f_X(x) = (1-p)^{x-1} p$$

$$Y: S \rightarrow \mathcal{N} \setminus \{0\}, f_Y(y) = (1-p)^{y-1} p$$

denote the sum of them as $Z = X+Y$ where $Z: S \rightarrow \mathcal{N} \setminus \{0\}$.

$$\text{then } P[Z=z] = \sum_{x+y=z} P[X=x] \cdot P[Y=y] = \sum_{x+y=z} f_X(x) \cdot f_Y(y) \text{ using the conclusion above.}$$

$$= \sum_{x+y=z} (1-p)^{x-1} p \cdot (1-p)^{y-1} p = \sum_{x+y=z} (1-p)^{x+y-2} p^2 = \sum_{x+y=z} (1-p)^{z-2} p^2$$

$$\text{since } z \geq 1, \text{ then } x \leq z-1 \text{ since } x \geq 1 \Rightarrow 1 \leq x \leq z-1 \in \mathcal{N} \setminus \{0\}.$$



$\Rightarrow X$ can have $z-1$ possible values ranging from 1 to $z-1$ and we can get the corresponding

$$y = z - x$$

therefore, the ways of getting $x+y = z$ are in total $\binom{z-1}{1}$

$\Rightarrow f_z(z) = \binom{z-1}{1} (1-p)^{z-2} p^2$, which is a Pascal distribution with $r=2$.

Problem 4.

the initial value should be $P_0(0) = 1$

$P_0(t)$ means in the time period $[0, t]$, the probability of no arrival should be 1, which means there must be no arrival.

then with conditions: $P_0'(t) = -\lambda P_0(t)$, $P_k'(t) + \lambda P_k(t) = \lambda P_{k-1}(t)$, we use induction to prove that $P_k(t) = \frac{(\lambda t)^k e^{-\lambda t}}{k!}$

① when $k=0$, since $\frac{dP_0}{dt} = -\lambda P_0 \Rightarrow \frac{dP_0}{P_0} = -\lambda dt \Rightarrow \ln|P_0| = -\lambda t + C \Rightarrow P_0 = C e^{-\lambda t}$

since $P_0(0) = 1$, then $P_0(t) = C e^0 = C = 1 \Rightarrow P_0(t) = e^{-\lambda t}$

since $P_0(t) = \frac{(\lambda t)^0 e^{-\lambda t}}{0!} = e^{-\lambda t}$, then the conclusion is right.

② for $k=k, k \geq 0$, suppose $P_k(t) = \frac{(\lambda t)^k e^{-\lambda t}}{k!}$

then since $P_{k+1}'(t) + \lambda P_{k+1}(t) = \lambda P_k(t) = \frac{\lambda^{k+1} t^k e^{-\lambda t}}{k!}$

$$\mu = e^{\int \lambda dt} = e^{\lambda t}, \int \frac{\lambda^{k+1} t^k e^{-\lambda t}}{k!} \times e^{\lambda t} dt = \int \frac{\lambda^{k+1} t^k}{k!} dt = \frac{\lambda^{k+1}}{(k+1)!} t^{k+1} = \frac{(\lambda t)^{k+1}}{(k+1)!}$$

$$\Rightarrow P_{k+1} = e^{-\lambda t} \left(\frac{(\lambda t)^{k+1}}{(k+1)!} + C \right) = \frac{(\lambda t)^{k+1} e^{-\lambda t}}{(k+1)!} + C e^{-\lambda t}$$

since $P_{k+1}(0) = 0$ because in $[0, 0]$, the probability of $k+1$ arrivals should be 0.

then $P_{k+1}(0) = C = 0$

$$\Rightarrow P_{k+1} = \frac{(\lambda t)^{k+1} e^{-\lambda t}}{(k+1)!}, \text{ which accords with } P_k(t) = \frac{(\lambda t)^k e^{-\lambda t}}{k!}$$

Based on ① and ②, we can conclude that $P_k(t) = \frac{(\lambda t)^k e^{-\lambda t}}{k!}$ holds true for $\forall k \in \mathbb{N}$, then it gets proved.

Problem 5.

$$f(x) = \binom{n}{x} p^x (1-p)^{n-x} = \frac{n!}{x!(n-x)!} \left(\frac{k}{n}\right)^x \left(1 - \frac{k}{n}\right)^{n-x}$$

when $n \rightarrow \infty$, using Stirling's formula: $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$ as $n \rightarrow \infty$.

$$\begin{aligned} \text{then } f(x) &= \frac{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n}{x!(n-x)!} \left(\frac{k}{n}\right)^x \left(1 - \frac{k}{n}\right)^{n-x} \text{ as } n \rightarrow \infty \\ &= \frac{k^x}{x!} \frac{\sqrt{2\pi n} \left(\frac{n}{e}\right)^n}{\sqrt{2\pi(n-x)} \left(\frac{n-x}{e}\right)^{n-x}} \cdot n^{-x} \cdot \left(1 - \frac{k}{n}\right)^n \cdot \left(1 - \frac{k}{n}\right)^{-x} \end{aligned}$$

$$\text{since } \lim_{n \rightarrow \infty} \left(1 - \frac{k}{n}\right)^n = e^{-k}, \text{ then } \lim_{n \rightarrow \infty} f(x) = \frac{e^{-k} k^x}{x!} \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n-x}} \cdot \left(\frac{n}{n-x}\right)^{n-x} \cdot e^{-x} \cdot \left(1 - \frac{k}{n}\right)^{-x}$$



when $n \rightarrow \infty$, it's obvious that $\sqrt{\frac{n}{n-k}} = \sqrt{\frac{1}{1-\frac{k}{n}}} \rightarrow 1$ and $(1-\frac{k}{n})^{-k} \rightarrow 1$.

$$\text{then } \lim_{n \rightarrow \infty} f(x) = \frac{e^{-k} k^x}{x!} \left(\lim_{n \rightarrow \infty} \left(\frac{n}{n-k} \right)^{n-k} \right) \cdot e^{-k}.$$

$$\text{since } \lim_{n \rightarrow \infty} \left(\frac{n}{n-k} \right)^{n-k} = \lim_{n \rightarrow \infty} \left(1 + \frac{k}{n-k} \right)^{n-k} = \lim_{n \rightarrow \infty} \left(1 + \frac{k}{n-k} \right)^{n-k} = e^k.$$


$$\begin{aligned} \text{then } \lim_{n \rightarrow \infty} f(x) &= \frac{e^{-k} k^x}{x!} e^k \cdot e^{-k} \\ &= \frac{e^{-k} k^x}{x!} \end{aligned}$$

Problem 6.

i) since $\frac{1}{b-a} > 0$, then $f(x) \geq 0$ for all $x \in \mathbb{R}$.

$$\text{and } \int_{-\infty}^{\infty} f(x) dx = 0 + \int_a^b \frac{1}{b-a} dx = \left[\frac{x}{b-a} \right]_a^b = \frac{b-a}{b-a} = 1.$$

therefore, it's a density for a continuous random variable.



$$\text{iii) } P\left[X \leq \frac{a+b}{2}\right] = \int_{-\infty}^{\frac{a+b}{2}} f(x) dx = 0 + \int_a^{\frac{a+b}{2}} \frac{1}{b-a} dx = \left[\frac{x}{b-a} \right]_a^{\frac{a+b}{2}} = \frac{\frac{a+b}{2} - a}{b-a} = \frac{\frac{b-a}{2}}{b-a} = \frac{1}{2}.$$

$$\text{iv) } P[c \leq X \leq d] = \int_c^d f(x) dx = \int_c^d \frac{1}{b-a} dx = \left[\frac{x}{b-a} \right]_c^d = \frac{d-c}{b-a}, \text{ since } (c, d) \text{ is subinterval of } (a, b).$$

$$P[e \leq X \leq f] = \int_e^f f(x) dx = \int_e^f \frac{1}{b-a} dx = \left[\frac{x}{b-a} \right]_e^f = \frac{f-e}{b-a}, \text{ since } (e, f) \text{ is subinterval of } (a, b).$$

since (c, d) and (e, f) are of equal length, then $d-c = f-e \Rightarrow \frac{d-c}{b-a} = \frac{f-e}{b-a}$

$$\text{then } P[c \leq X \leq d] = P[e \leq X \leq f]$$

$$\text{v) } F(x) = P[X \leq x] = \int_{-\infty}^x f(y) dy$$

if $x \leq a$, then $f(y) = 0 \Rightarrow F(x) = 0$.

$$\text{if } a < x < b, \text{ then } F(x) = \int_{-\infty}^a 0 dy + \int_a^x \frac{1}{b-a} dy = 0 + \left[\frac{y}{b-a} \right]_a^x = \frac{x-a}{b-a}$$

$$\text{if } x \geq b, \text{ then } F(x) = \int_{-\infty}^a 0 dy + \int_a^b \frac{1}{b-a} dy + \int_b^x 0 dy = 0 + \frac{b-a}{b-a} + 0 = 1.$$

$$\Rightarrow F(x) = \begin{cases} 0 & , x \leq a \\ \frac{x-a}{b-a} & , a < x < b \\ 1 & , x \geq b. \end{cases}$$




$$\begin{aligned}
 \text{vii). } m_x &= E[e^{tx}] = \int_{-\infty}^{\infty} e^{tx} f(x) dx = \int_{-\infty}^a e^{tx} \cdot 0 dx + \int_a^b e^{tx} \cdot \frac{1}{b-a} dx + \int_b^{\infty} e^{tx} \cdot 0 dx \\
 &= 0 + \left[\frac{e^{tx}}{b-a} \right]_a^b + 0 \\
 &= \frac{e^{tb} - e^{ta}}{t(b-a)}
 \end{aligned}$$

$$\Rightarrow E[X] = \frac{dm_x}{dt} \Big|_{t=0} = \frac{e^{tb} - e^{ta}}{t(b-a)} \Big|_{t=0} = \frac{b^1 - a^1}{b-a}$$

$$\text{vi). } E[X] = \int_{-\infty}^{\infty} x \cdot f(x) dx = \int_a^b x \cdot \frac{1}{b-a} dx = \frac{1}{b-a} \cdot \left[\frac{1}{2} x^2 \right]_a^b = \frac{1}{b-a} \cdot \frac{1}{2} \cdot (b^2 - a^2) = \frac{(b-a)(b+a)}{2(b-a)} = \frac{a+b}{2}$$

$$E[X^2] = \int_{-\infty}^{\infty} x^2 \cdot f(x) dx = \int_a^b x^2 \cdot \frac{1}{b-a} dx = \frac{b^3 - a^3}{3(b-a)} = \frac{b^2 + ab + a^2}{3}$$

$$\text{Var}[X] = E[X^2] - E[X]^2 = \frac{b^2 + ab + a^2}{3} - \frac{a^2 + b^2 + 2ab}{4} = \frac{4b^2 + 4ab + 4a^2 - 3a^2 - 3b^2 - 6ab}{12} = \frac{(b-a)^2}{12}$$



$$\frac{d}{dx} \left(\frac{1}{b-a} \right) = \frac{d}{dx} \left(\frac{1}{b-a} \right) = 0$$

$$\text{For } x < a, f(x) = 0 \Rightarrow \frac{d}{dx} f(x) = 0$$

$$\text{For } x > b, f(x) = 0 \Rightarrow \frac{d}{dx} f(x) = 0$$

$$\frac{d}{dx} f(x) = \frac{d}{dx} \left(\frac{1}{b-a} \right) = 0$$

$$[f(x) \cdot x] = \left[\frac{1}{b-a} \cdot x \right]$$

$$[f(x) \cdot x^2] = \left[\frac{1}{b-a} \cdot x^2 \right]$$

$$\frac{d}{dx} [f(x) \cdot x] = \frac{d}{dx} \left(\frac{x}{b-a} \right) = \frac{1}{b-a}$$

$$\frac{d}{dx} [f(x) \cdot x^2] = \frac{d}{dx} \left(\frac{x^2}{b-a} \right) = \frac{2x}{b-a}$$

$$1 = 0 + \frac{1}{b-a} + 0 = \frac{1}{b-a}$$

