

Below, we can see the original image that was described in the question:




Figure 1: The original 201×201 black image with a white row in the middle.

A plot of the logarithm of the modulus of the DFT of the image can be found below:




Figure 2: The log plot of the modulus of the DFT of Figure (1).

0.1 Analytical Derivation

We know that the formula for the DFT of a 2D signal is as follows:

$$F_d(u, v) = \frac{1}{\sqrt{W_1 W_2}} \sum_{x=0}^{W_1-1} \sum_{y=0}^{W_2-1} f(x, y) \exp \left(-i2\pi \left(\frac{ux}{W_1} + \frac{vy}{W_2} \right) \right)$$

Since we know that in our image, $W_1 = W_2 = 201$, we can substitute this into the formula:

$$F_d(u, v) = \frac{1}{201} \sum_{x=0}^{200} \sum_{y=0}^{200} f(x, y) \exp \left(-i2\pi \left(\frac{ux}{201} + \frac{vy}{201} \right) \right) \quad (1)$$

In our case, our 2D signal is the image shown in Figure 1, and its definition is

$$f(x, y) = \begin{cases} 0, y \in \{0, 1, 2, \dots, 200\} \text{ and } y \neq 100, & x \in \{0, 1, 2, \dots, 200\} \\ 1, y = 100, & x \in \{0, 1, 2, \dots, 200\} \end{cases}$$

Observe that we normalized the image intensities, so the range of intensity values is $[0, 1]$, instead of the usual $\{0, 1, \dots, 255\}$. Moreover, due to the indices starting from 0 instead of 1 (like in MATLAB), the index of the white pixel row is 100 instead of 101.

Since most of the image consists of 0-intensity pixels, Equation (1) simplifies to

$$\begin{aligned} F_d(u, v) &= \frac{1}{201} \sum_{x=0}^{200} f(x, 100) \exp \left(\frac{-2\pi i(ux + 100v)}{201} \right) \\ &= \frac{1}{201} \sum_{x=0}^{200} \exp \left(\frac{-2\pi i(ux + 100v)}{201} \right) \end{aligned}$$