

Patient Name: **TEST, DAVID1**Case #: **D-08-0013908**DOB/Age/Sex: 1/1/1951 57 years Male  
MRN: 545454545Collected: 2/29/2008 12:01:00 PM  
Received: 2/29/2008 12:01:00 PMClient Name: TEST CLIENT CLIENT TEST%  
Provider: DOC TEST1 MD  
Consulting:Deliver to: - 12345678910,TEST DOCTOR  
1923 S UTICA  
TULSA, OK 74104

---

**SURGICAL PATHOLOGY REPORT**

---

**Diagnosis**Skin, chest, punch biopsy --  
Grover's disease

Test, Pathologist Pathologist (Electronic Signature)

PT 02/29/2008

---

**Microscopic Examination**

There are focal areas of suprabasal clefting and a few acantholytic cells are seen. There is villous formation and individual dyskeratotic cells are present in the upper layers of the epidermis.

Corps ronds and grains are present. Grover's disease is favored.

**Gross Examination**

Punch biopsy of skin: chest

Size: 0.4 x 0.4 cm

Excision depth: 0.6 cm

Specimen is bisected and entirely submitted in one cassette for microscopic examination.

PT /PT

**Specimen**

From chest

**Pertinent History**

Grover's disease vs. other