

Modeling PaLEON biomass

Wesley Brooks

UW-Madison

May 27, 2013

Outline

Goal

- Produce a model of per-species biomass at time of settlement

Table of Contents

Data

- Computed from settlement-era survey
- Working with composition, biomass, and stem density

Models

There are two divisions for modeling biomass data:

		Estimation method		
		<i>Splines</i>	<i>GMRF</i>	<i>GLM</i>
Model	One-stage	Zero-inflated gamma		
		Tweedie		
	Two-stage	Bernoulli-Gamma		

Two-stage models

- First stage: zero/non-zero
 - ▶ Logistic regression
 - ▶ $Z \sim \text{Bernoulli}(\gamma)$
 - ▶ $\text{logit}(\gamma) = f(x, y, p_k)$
- Second stage: distribution of positive biomass
 - ▶ $Y|Z = 1 \sim \text{Gamma}(\alpha, \beta)$
 - ▶ $E(Y|Z = 1) = \mu = \alpha\beta = f(x, y, p_k)$

Two-stage models

Mean and variance of the two-stage model:

- $E(Y) = \gamma\mu$
- $\text{var}(Y) = \gamma\alpha\beta^2(1 + \alpha(1 - \gamma))$

Tweedie model

The Tweedie model is a Gamma-Poisson mixture.

- Draw $N \sim \text{Poisson}(\lambda)$
- Now make N iid draws: $V_\ell \sim \text{Gamma}(\alpha, \beta)$
- $Y = \sum_{\ell=1}^N V_\ell$

Tweedie model

With θ given, the Tweedie distribution is in the exponential family with natural parameter ν :

- $f(y) = \exp \{ \phi^{-1} [y\nu - \kappa(\nu)] + c(y, \phi, \nu) \}$
- ϕ is a scale parameter
- $\nu = \mu^{1-\theta} / (1 - \theta)$
- $\kappa(\nu) = [(1 - \theta)\nu]^{(2-\theta)/(1-\theta)} / (2 - \theta) = \mu^{2-\theta} / (2 - \theta)$

Table of Contents