

1. Selection

Theorem 1.1. *If $hn^{-1/2}b_n \xrightarrow{p} \infty$ then $P\left\{\hat{\beta}_{(b)}(s) = 0\right\} \rightarrow 1$.*

Proof. The proof is by contradiction.

Recall that the objective to be minimized by $\hat{\beta}_{(p)}(s)$ is

$$Q\left\{\hat{\beta}(s)\right\} = (1/2)\left\{Y - Z(s)\hat{\beta}(s)\right\}^T W(s)\left\{Y - Z(s)\hat{\beta}(s)\right\} + \sum_{j=1}^p \lambda_j \|\hat{\beta}_p(s)\| \quad (1)$$

Let $\hat{\beta}_p(s) \neq 0$. Then (1) is differentiable w.r.t. $\beta_p(s)$ and Q is maximized at

$$\begin{aligned} 0 &= Z_{(p)}^T(s)W(s)\left\{Y - Z_{(-p)}(s)\hat{\beta}_{(-p)}(s) - Z_{(p)}(s)\hat{\beta}_{(p)}(s)\right\} - \lambda_p \frac{\hat{\beta}_{(p)}(s)}{\|\hat{\beta}_{(p)}(s)\|} \\ &= Z_{(p)}^T(s)W(s)\left[Y - Z(s)\beta(s) - \frac{h^2\kappa_2}{2\kappa_0}\{\beta_{uu}(s) + \beta_{vv}(s)\}\right] \\ &\quad + Z_{(p)}^T(s)W(s)Z_{(-p)}(s)\left[\beta_{(-p)}(s) + \frac{h^2\kappa_2}{2\kappa_0}\{\beta_{(-p),uu}(s) + \beta_{(-p),vv}(s)\} - \hat{\beta}_{(-p)}(s)\right] \\ &\quad + Z_{(p)}^T(s)W(s)Z_{(p)}(s)\left[\beta_{(p)}(s) + \frac{h^2\kappa_2}{2\kappa_0}\{\beta_{(p),uu}(s) + \beta_{(p),vv}(s)\} - \hat{\beta}_{(p)}(s)\right] \\ &\quad - \lambda_p \frac{\hat{\beta}_{(p)}(s)}{\|\hat{\beta}_{(p)}(s)\|} \end{aligned} \quad (2)$$

So

$$\begin{aligned}
\frac{h}{\sqrt{n}} \lambda_p \frac{\hat{\beta}_{(p)}(\mathbf{s})}{\|\hat{\beta}_{(p)}(\mathbf{s})\|} &= \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \frac{h}{\sqrt{n}} \left[\mathbf{Y} - \mathbf{Z}(\mathbf{s}) \boldsymbol{\beta}(\mathbf{s}) - \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{uu}(\mathbf{s}) + \boldsymbol{\beta}_{vv}(\mathbf{s})\} \right] \\
&+ \left\{ n^{-1} \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(-p)}(\mathbf{s}) \right\} h\sqrt{n} \left[\boldsymbol{\beta}_{(-p)}(\mathbf{s}) + \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{(-p),uu}(\mathbf{s}) + \boldsymbol{\beta}_{(-p),vv}(\mathbf{s})\} - \hat{\boldsymbol{\beta}}_{(-p)}(\mathbf{s}) \right] \\
&+ \left\{ n^{-1} \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(-p)}(\mathbf{s}) \right\} h\sqrt{n} \left[\boldsymbol{\beta}_{(-p)}(\mathbf{s}) + \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{(p),uu}(\mathbf{s}) + \boldsymbol{\beta}_{(p),vv}(\mathbf{s})\} - \hat{\boldsymbol{\beta}}_{(p)}(\mathbf{s}) \right]
\end{aligned} \tag{3}$$

From Lemma 2 of ?, $\left\{ n^{-1} \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(-p)}(\mathbf{s}) \right\} = O_p(1)$ and $\left\{ n^{-1} \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(p)}(\mathbf{s}) \right\} = O_p(1)$.

From Theorem 3 of ?, we have that $h\sqrt{n} \left[\hat{\boldsymbol{\beta}}_{(-p)}(\mathbf{s}) - \boldsymbol{\beta}_{(-p)}(\mathbf{s}) - \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{(p),uu}(\mathbf{s}) + \boldsymbol{\beta}_{(p),vv}(\mathbf{s})\} \right] = O_p(1)$ and $h\sqrt{n} \left[\hat{\boldsymbol{\beta}}_{(p)}(\mathbf{s}) - \boldsymbol{\beta}_{(p)}(\mathbf{s}) - \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{(p),uu}(\mathbf{s}) + \boldsymbol{\beta}_{(p),vv}(\mathbf{s})\} \right] = O_p(1)$.

So the second and third terms of the sum in (3) are $O_p(1)$.

We showed in the proof of ?? that $h\sqrt{n} \mathbf{Z}_{(p)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \left[\mathbf{Y} - \mathbf{Z}(\mathbf{s}) \boldsymbol{\beta}(\mathbf{s}) - \frac{h^2 \kappa_2}{2\kappa_0} \{\boldsymbol{\beta}_{uu}(\mathbf{s}) + \boldsymbol{\beta}_{vv}(\mathbf{s})\} \right] = O_p(1)$.

The three terms of the sum to the right of the equals sign in (3) are $O_p(1)$, so for $\hat{\boldsymbol{\beta}}_{(p)}(\mathbf{s})$ to be a solution, we must have that $hn^{-1/2} \lambda_p \frac{\hat{\beta}_{(p)}(\mathbf{s})}{\|\hat{\beta}_{(p)}(\mathbf{s})\|} = O_p(1)$.

But since by assumption $\hat{\beta}_{(p)}(\mathbf{s}_i) \neq 0$, there must be some $k \in \{1, \dots, d_p\}$ such that $|\hat{\beta}_{(p),k}(\mathbf{s})| = \max\{|\hat{\beta}_{(p),k'}(\mathbf{s})| : 1 \leq k' \leq d_p\}$. And for this k , we have that $|\hat{\beta}_{(p),k}(\mathbf{s})|/\|\hat{\boldsymbol{\beta}}_{(p)}(\mathbf{s})\| \geq 1/\sqrt{d_p} > 0$.

Now since $hn^{-1/2} b_n \rightarrow \infty$, we have that $hn^{-1/2} \lambda_p \frac{\hat{\beta}_{(p)}(\mathbf{s})}{\|\hat{\beta}_{(p)}(\mathbf{s})\|} \geq hn^{-1/2} b_n d_p^{-1/2} \rightarrow \infty$ and therefore the term to the left of the equals sign dominates the sum to the right of the equals sign in (3). So for large enough n , $\hat{\boldsymbol{\beta}}_{(p)}(\mathbf{s}) \neq 0$ cannot maximize Q .

So $P \left\{ \hat{\beta}_{(b)}(\mathbf{s}) = 0 \right\} \rightarrow 1$. □

Theorem 1.2. *If $h\sqrt{na_n} \xrightarrow{p} 0$ then $P\left\{\hat{\beta}_{(a)}(\mathbf{s}) \neq 0\right\} \rightarrow 1$.*

Proof. Again, the proof is by contradiction.

Assume that $\hat{\beta}_{(k)} = 0$ for some $k \leq p_0$. For the adaptive group lasso, the covariate group k is shrunk to zero if

$$\left\| \left\{ \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(k)}^T(\mathbf{s}) \right\}^{-1} \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{r}_{(k)}(\mathbf{s}) \right\|^2 \leq \lambda_k^2$$

where $\mathbf{r}_{(k)}(\mathbf{s})$ is the residual after accounting for all covariate groups except group k . That is,

$\mathbf{r}_{(k)}(\mathbf{s}) = \mathbf{Y} - \mathbf{Z}_{(-k)}(\mathbf{s}) \boldsymbol{\beta}_{(-k)}(\mathbf{s})$. But $\|\tilde{\boldsymbol{\beta}}_{(k)}\| > 0$ implies that $\left\| \left\{ \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(k)}^T(\mathbf{s}) \right\}^{-1} \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{r}_{(k)}(\mathbf{s}) \right\|^2 > 0$ and $\frac{h^2 n \lambda^2}{\|\tilde{\boldsymbol{\beta}}_{(k)}\|^2} \leq h^2 n a_n^2 \rightarrow 0$. So

$$P\left\{\hat{\beta}_{(k)}(\mathbf{s}) \neq 0\right\} \leq P\left[\left\| \left\{ \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{Z}_{(k)}^T(\mathbf{s}) \right\}^{-1} \mathbf{Z}_{(k)}^T(\mathbf{s}) \mathbf{W}(\mathbf{s}) \mathbf{r}_{(k)}(\mathbf{s}) \right\|^2 \leq h^2 n a_n^2 \right] \rightarrow 0. \quad (4)$$

□