

# SPECTRAL METHODS IN GAUSSIAN MODELLING

## TOPIC 2: KERNEL DESIGN

**James Requiema and Wessel Bruinsma**

University of Cambridge and Invenia Labs

21 January 2019

- + RFFs alleviate the  $O(N^3)$  scaling.

- + RFFs alleviate the  $O(N^3)$  scaling.
- RFFs do not help with choice of kernel.

- + RFFs alleviate the  $O(N^3)$  scaling.
- RFFs do not help with choice of kernel.

**How to parametrise a flexible kernel?**

- Bochner's Theorem:

$$k(\tau) \xleftrightarrow{\mathcal{F}} s(\omega) = \text{PSD}.$$

- Bochner's Theorem:

$$k(\tau) \xleftrightarrow{\mathcal{F}} s(\omega) = \text{PSD}.$$

- PSD:
  - distribution of power contained in frequencies,
  - must be **nonnegative** and **symmetric**.

- Bochner's Theorem:

$$k(\tau) \xleftrightarrow{\mathcal{F}} s(\omega) = \text{PSD}.$$

- PSD:
  - distribution of power contained in frequencies,
  - must be **nonnegative** and **symmetric**.
- Easier to flexibly parametrise PSD!

- SSA (Lázaro-Gredilla et al., 2010) models PSD with **symmetric average of lines**:

$$s(\omega) = \frac{1}{2Q} \sum_{q=1}^Q (\delta(\omega - \mu^{(q)}) + \delta(\omega + \mu^{(q)})).$$



- SSA (Lázaro-Gredilla et al., 2010) models PSD with **symmetric average of lines**:

$$s(\omega) = \frac{1}{2Q} \sum_{q=1}^Q (\delta(\omega - \mu^{(q)}) + \delta(\omega + \mu^{(q)})).$$

- Inverse Fourier transform gives kernel:

$$k(\tau) = \frac{1}{Q} \sum_{q=1}^Q \cos(\mu^{(q)\top} \tau).$$

- SSA (Lázaro-Gredilla et al., 2010) models PSD with **symmetric average of lines**:

$$s(\omega) = \frac{1}{2Q} \sum_{q=1}^Q (\delta(\omega - \mu^{(q)}) + \delta(\omega + \mu^{(q)})).$$

- Inverse Fourier transform gives kernel:

$$k(\tau) = \frac{1}{Q} \sum_{q=1}^Q \cos(\mu^{(q)\top} \tau).$$

- Strong parametric assumption:  $f(t)$  = sum of sines.

- SMK (Wilson and Adams, 2013) models PSD with **symmetric mixture of Gaussians**:

$$s(\omega) = \frac{1}{2} \sum_{q=1}^Q w^{(q)} \left( \mathcal{N}(\omega; \mu^{(q)}, \Sigma^{(q)}) + \mathcal{N}(\omega; -\mu^{(q)}, \Sigma^{(q)}) \right).$$

- SMK (Wilson and Adams, 2013) models PSD with **symmetric mixture of Gaussians**:

$$s(\omega) = \frac{1}{2} \sum_{q=1}^Q w^{(q)} \left( \mathcal{N}(\omega; \mu^{(q)}, \Sigma^{(q)}) + \mathcal{N}(\omega; -\mu^{(q)}, \Sigma^{(q)}) \right).$$

- $w^{(q)} = 1/Q$  and  $\Sigma^{(q)} \rightarrow 0$  recovers SSA.

- SMK (Wilson and Adams, 2013) models PSD with **symmetric mixture of Gaussians**:

$$s(\omega) = \frac{1}{2} \sum_{q=1}^Q w^{(q)} \left( \mathcal{N}(\omega; \mu^{(q)}, \Sigma^{(q)}) + \mathcal{N}(\omega; -\mu^{(q)}, \Sigma^{(q)}) \right).$$

- $w^{(q)} = 1/Q$  and  $\Sigma^{(q)} \rightarrow 0$  recovers SSA.
- Inverse Fourier transform gives kernel:

$$k^{(\text{SMK})}(\tau) = \sum_{q=1}^Q w^{(q)} \exp\left(-\frac{1}{2}\tau^\top \Sigma^{(q)} \tau\right) \cos\left(\mu^{(q)\top} \tau\right),$$

- Equivalent generative model as a truncated Fourier series:

$$f^{(\text{SMK})}(t) = \sum_{q=1}^Q \sqrt{w^{(q)}} (c_1^{(q)}(t) \cos(\mu^{(q)\top} t) + c_2^{(q)}(t) \sin(\mu^{(q)\top} t)),$$

$$c_1^{(q)}, c_2^{(q)} \sim \mathcal{GP}(0, \exp(-\frac{1}{2} \tau^\top \Sigma^{(q)} \tau)).$$

- Equivalent generative model as a truncated Fourier series:

$$f^{(\text{SMK})}(t) = \sum_{q=1}^Q \sqrt{w^{(q)}} (c_1^{(q)}(t) \cos(\mu^{(q)\top} t) + c_2^{(q)}(t) \sin(\mu^{(q)\top} t)),$$

$$c_1^{(q)}, c_2^{(q)} \sim \mathcal{GP}(0, \exp(-\frac{1}{2} \tau^\top \Sigma^{(q)} \tau)).$$

- In SSA,  $(c_1^{(q)}, c_2^{(q)})_{q=1}^Q$  are constant.

- Equivalent generative model as a truncated Fourier series:

$$f^{(\text{SMK})}(t) = \sum_{q=1}^Q \sqrt{w^{(q)}} (c_1^{(q)}(t) \cos(\mu^{(q)\top} t) + c_2^{(q)}(t) \sin(\mu^{(q)\top} t)),$$

$$c_1^{(q)}, c_2^{(q)} \sim \mathcal{GP}(0, \exp(-\frac{1}{2} \tau^\top \Sigma^{(q)} \tau)).$$

- In SSA,  $(c_1^{(q)}, c_2^{(q)})_{q=1}^Q$  are constant.
- SMK fattens spectral lines by allowing  $c_1^{(q)}$  and  $c_2^{(q)}$  to vary with time.



- + Flexible, drop-in replacement

- + Flexible, drop-in replacement
- + Can recover many standard kernels

- + Flexible, drop-in replacement
- + Can recover many standard kernels
- + Models negative covariances

- + Flexible, drop-in replacement
- + Can recover many standard kernels
- + Models negative covariances
- Unclear how many components needed

- + Flexible, drop-in replacement
- + Can recover many standard kernels
- + Models negative covariances
- Unclear how many components needed
- Hyperparameters difficult to optimise

- MOSMK (Parra and Tobar, 2017) generalises SMK to multiple outputs.

- MOSMK (Parra and Tobar, 2017) generalises SMK to multiple outputs.
- Uses multivariate extension of Bochner's Theorem: Cramér's Theorem.

- MOSMK (Parra and Tobar, 2017) generalises SMK to multiple outputs.
- Uses multivariate extension of Bochner's Theorem: Cramér's Theorem.
- Multivariate PSD  $S: \mathbb{R}^D \rightarrow \mathbb{C}^{P \times P}$ .



- MOSMK (Parra and Tobar, 2017) generalises SMK to multiple outputs.
- Uses multivariate extension of Bochner's Theorem: Cramér's Theorem.
- Multivariate PSD  $S: \mathbb{R}^D \rightarrow \mathbb{C}^{P \times P}$ .
  - Must be **symmetric**:  $S(\omega) = S^\dagger(-\omega)$ ,  $S_{ii}(\omega) = S_{ii}(-\omega)$ .

- MOSMK (Parra and Tobar, 2017) generalises SMK to multiple outputs.
- Uses multivariate extension of Bochner's Theorem: Cramér's Theorem.
- Multivariate PSD  $S: \mathbb{R}^D \rightarrow \mathbb{C}^{P \times P}$ .
  - Must be **symmetric**:  $S(\omega) = S^\dagger(-\omega)$ ,  $S_{ii}(\omega) = S_{ii}(-\omega)$ .
  - Must be **nonnegative**:  $S(\omega) \geq 0$ .

- MOSMK models PSD with **symmetric mixture of outer products of vectors of Gaussians**:

$$S(\omega) = \frac{1}{2} \sum_{q=1}^Q \left( R^{(q)}(\omega) R^{(q)\dagger}(\omega) + R^{(q)}(-\omega) R^{(q)\dagger}(-\omega) \right),$$

$$R_i^{(q)}(\omega) = w^{(q)} \exp \left( -\frac{1}{4}(\omega - \mu_i^{(q)}) \Sigma_i^{(q)-1} (\omega - \mu_i^{(q)}) - \iota(\theta_i^{(q)\top} \omega + \phi_i^{(q)}) \right).$$

- Inverse Fourier transform gives kernel:

$$K_{ij}^{(\text{MOSMK})}(\tau) = \sum_{q=1}^Q \alpha_{ij}^{(q)} \exp\left(-\frac{1}{2}(\tau + \theta_{ij}^{(q)})^\top \Sigma_{ij}^{(q)} (\tau + \theta_{ij}^{(q)})\right) \\ \times \cos\left((\tau + \theta_{ij}^{(q)})^\top \mu_{ij}^{(q)} + \phi_{ij}^{(q)}\right).$$

- Equivalent generative model as truncated Fourier series:

$$\begin{aligned}
f_i^{(\text{MOSMK})}(t) &= \sum_{q=1}^Q w_i^{(q)} \left( c_{i1}^{(q)}(t - \theta_i^{(q)}) \cos\left(\mu_i^{(q)\top}(t - \theta_i^{(q)}) + \phi_i^{(q)}\right) \right. \\
&\quad \left. + c_{i2}^{(q)}(t - \theta_i^{(q)}) \sin\left(\mu_i^{(q)\top}(t - \theta_i^{(q)}) + \phi_i^{(q)}\right) \right), \\
\mathbb{E}[c_{ik}^{(p)}(t)c_{jl}^{(q)}(t')] &= \begin{cases} \frac{\alpha_{ij}^{(q)}}{w_i^{(q)}w_j^{(q)}} \exp\left(-\frac{1}{2}(t-t')^\top \Sigma_{ij}^{(q)}(t-t')\right) & \text{if } k=l, p=q, \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

- GSMK (Chen et al., 2018) generalises SMK to nonstationary signals.

- GSMK (Chen et al., 2018) generalises SMK to nonstationary signals.
- Uses the *Gibbs kernel* (Gibbs, 1997):

$$k^{(\text{Gibbs})}(t, t') = \prod_{d=1}^D \sqrt{\frac{2\ell_d(t)\ell_d(t')}{\ell_d^2(t) + \ell_d^2(t')}} \exp\left(-\sum_{d=1}^D \frac{(t_d - t'_d)^2}{\ell_d^2(t) + \ell_d^2(t')}\right).$$

## Nonstationary EQ Kernel

- Cannot simply make length scale input dependent.



## Nonstationary EQ Kernel

- Cannot simply make length scale input dependent.
- Construction of EQ from basis functions:

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{\ell^2}(t - c)^2\right),$$

## Nonstationary EQ Kernel

- Cannot simply make length scale input dependent.
- Construction of EQ from basis functions:

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{\ell^2}(t - c)^2\right),$$
$$f(t) | n = \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')),$$

## Nonstationary EQ Kernel

- Cannot simply make length scale input dependent.
- Construction of EQ from basis functions:

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{\ell^2}(t - c)^2\right),$$

$$f(t) | n = \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')),$$

$$\mathbb{E}[f(t)f(t')] = \int_{-\infty}^{\infty} \phi(t; c)\phi(t'; c) \, dc$$

## Nonstationary EQ Kernel

- Cannot simply make length scale input dependent.
- Construction of EQ from basis functions:

$$\begin{aligned}\phi(t; c) &= \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{\ell^2}(t - c)^2\right), \\ f(t) | n &= \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')), \\ \mathbb{E}[f(t)f(t')] &= \int_{-\infty}^{\infty} \phi(t; c)\phi(t'; c) \, dc \\ &= \exp\left(-\frac{1}{2\ell^2}(t - t')^2\right).\end{aligned}$$

## Nonstationary EQ Kernel

- Make length scale of  $\phi$  dependent on  $t$ :

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell(t)} \right)^{\frac{1}{2}} \exp\left( -\frac{1}{\ell^2(t)} (t - c)^2 \right),$$

## Nonstationary EQ Kernel

- Make length scale of  $\phi$  dependent on  $t$ :

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell(t)} \right)^{\frac{1}{2}} \exp\left( -\frac{1}{\ell^2(t)} (t - c)^2 \right),$$
$$f(t) | n = \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')),$$

## Nonstationary EQ Kernel

- Make length scale of  $\phi$  dependent on  $t$ :

$$\phi(t; c) = \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell(t)} \right)^{\frac{1}{2}} \exp\left(-\frac{1}{\ell^2(t)}(t - c)^2\right),$$

$$f(t) | n = \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')),$$

$$\mathbb{E}[f(t)f(t')] = \int_{-\infty}^{\infty} \phi(t; c)\phi(t'; c) \, dc$$

## Nonstationary EQ Kernel

- Make length scale of  $\phi$  dependent on  $t$ :

$$\begin{aligned}\phi(t; c) &= \left( \sqrt{\frac{2}{\pi}} \frac{1}{\ell(t)} \right)^{\frac{1}{2}} \exp\left( -\frac{1}{\ell^2(t)} (t - c)^2 \right), \\ f(t) | n &= \int_{-\infty}^{\infty} \phi(t; c) n(c) \, dc, \quad n(t) \sim \mathcal{GP}(0, \delta(t - t')), \\ \mathbb{E}[f(t)f(t')] &= \int_{-\infty}^{\infty} \phi(t; c) \phi(t'; c) \, dc \\ &= \sqrt{\frac{2\ell(t)\ell(t')}{\ell^2(t) + \ell^2(t')}} \exp\left( -\frac{(t - t')^2}{\ell^2(t) + \ell^2(t')} \right).\end{aligned}$$



- GSMK replaces the EQs with Gibbs kernels:

$$k^{(\text{GSMK})}(t, t') = \sum_{q=1}^Q w^{(q)}(t) w^{(q)}(t') k_q^{(\text{Gibbs})}(t, t') \\ \times \cos\left(\mu^{(q)\top}(t)t - \mu^{(q)\top}(t')t'\right).$$

- GSMK replaces the EQs with Gibbs kernels:

$$k^{(\text{GSMK})}(t, t') = \sum_{q=1}^Q w^{(q)}(t) w^{(q)}(t') k_q^{(\text{Gibbs})}(t, t') \\ \times \cos\left(\mu^{(q)\top}(t)t - \mu^{(q)\top}(t')t'\right).$$

- $(w^{(q)}, \ell^{(q)} \mu^{(q)})_{q=1}^Q$  given log-GP priors.

- GSMK replaces the EQs with Gibbs kernels:

$$k^{(\text{GSMK})}(t, t') = \sum_{q=1}^Q w^{(q)}(t) w^{(q)}(t') k_q^{(\text{Gibbs})}(t, t') \\ \times \cos\left(\mu^{(q)\top}(t)t - \mu^{(q)\top}(t')t'\right).$$

- $(w^{(q)}, \ell^{(q)} \mu^{(q)})_{q=1}^Q$  given log-GP priors.
- Estimated using MAP.

- Equivalent generative model as truncated Fourier series:

$$f^{(\text{GSMK})}(t) = \sum_{q=1}^Q w^{(q)}(t) (c_1^{(q)}(t) \cos(\mu^{(q)\top}(t)t) + c_2^{(q)}(t) \sin(\mu^{(q)\top}(t)t)),$$
$$c_1^{(q)}, c_2^{(q)} \sim \mathcal{GP}(0, k^{(\text{Gibbs})}(t, t')).$$

- SMK and extensions assume **parametric** model.

- SMK and extensions assume **parametric** model.
- More flexible to use **nonparametric** model:

$$s(\omega) = |\hat{h}(\omega)|^2.$$

- SMK and extensions assume **parametric** model.
- More flexible to use **nonparametric** model:

$$s(\omega) = |\hat{h}(\omega)|^2.$$

- Inverse Fourier transform gives kernel:

$$k(t, t') = \int_{-\infty}^{\infty} h(t - z)h(t' - z) \, dz = h * R(h)(t - t').$$

- SMK and extensions assume **parametric** model.
- More flexible to use **nonparametric** model:

$$s(\omega) = |\hat{h}(\omega)|^2.$$

- Inverse Fourier transform gives kernel:

$$k(t, t') = \int_{-\infty}^{\infty} h(t - z)h(t' - z) \, dz = h * R(h)(t - t').$$

- GPCM (Tobar et al., 2015) models  $h \sim \mathcal{GP}(0, k_h)$ .



- SMK and extensions assume **parametric** model.
- More flexible to use **nonparametric** model:


$$s(\omega) = |\hat{h}(\omega)|^2.$$

- Inverse Fourier transform gives kernel:

$$k(t, t') = \int_{-\infty}^{\infty} h(t - z)h(t' - z) \, dz = h * R(h)(t - t').$$


- GPCM (Tobar et al., 2015) models  $h \sim \mathcal{GP}(0, k_h)$ .
  - $\int_{-\infty}^{\infty} k_h(t, t) \, dt < \infty$  (finite trace).

- Nonparametric prior over kernels and PSDs.



- Interpretation as linear system:


- Interpretation as linear system:



$$\text{white noise} \sim \mathcal{GP}(0, \delta(t - t')),$$

$$h \sim \mathcal{GP}(0, k_h).$$

- Interpretation as linear system:



$$\text{white noise} \sim \mathcal{GP}(0, \delta(t - t')),$$

$$h \sim \mathcal{GP}(0, k_h).$$

- Inference complicated.

- Instead of designing kernel, design PSD.

- Instead of designing kernel, design PSD.
- Parametric approaches:
  - line spectrum (SSA),
  - mixture of Gaussians (SMK, MOSMK, GSMK).

- Instead of designing kernel, design PSD.
- Parametric approaches:
  - line spectrum (SSA),
  - mixture of Gaussians (SMK, MOSMK, GSMK).
- Nonparametric approach also possible (GPCM).