

Gaussian Process Convolution Model

Wessel Bruinsma

University of Cambridge, CBL

9 June 2019

$$f \sim \mathcal{GP}(0, k).$$

$$f \sim \mathcal{GP}(0, ???).$$

$$f \sim \mathcal{GP}(0, ???).$$

AA^T is P.S.D.

$$f \sim \mathcal{GP}(0, ???).$$

$$AA^T \text{ is P.S.D.}$$

$$“hh^T” = h * Rh \text{ is P.S.D.}$$

$$(Rh)(t) = h(-t)$$

$$f \sim \mathcal{GP}(0, ???).$$

$$AA^T \text{ is P.S.D.}$$

$$“hh^T” = h * Rh \text{ is P.S.D.}$$

$$(Rh)(t) = h(-t)$$


$$h \sim \mathcal{GP}(0, k_h),$$

$$f | h \sim \mathcal{GP}(0, h * Rh).$$

$$h \sim \mathcal{GP}(0, k_h),$$

$$k \mid h = h * Rh,$$

$$f \mid k \sim \mathcal{GP}(0, k):$$



Model (GPCM (Tobar et al., 2015))

$$h \sim \mathcal{GP}(0, k_h), \quad f | h \sim \mathcal{GP}(0, h * Rh).$$

Model (GPCM (Tobar et al., 2015))

$$h \sim \mathcal{GP}(0, k_h), \quad f | h \sim \mathcal{GP}(0, h * Rh).$$

$$x \sim \mathcal{N}(0, I) \implies Ax \sim \mathcal{N}(0, AA^T)$$

Model (GPCM (Tobar et al., 2015))

$$h \sim \mathcal{GP}(0, k_h), \quad f | h \sim \mathcal{GP}(0, h * Rh).$$

$$\begin{aligned} x \sim \mathcal{N}(0, I) &\implies Ax \sim \mathcal{N}(0, AA^T) \\ x \sim \mathcal{GP}(0, \delta) &\implies "hx" \sim \mathcal{GP}(0, "hh^T") \end{aligned}$$

Model (GPCM (Tobar et al., 2015))

$$h \sim \mathcal{GP}(0, k_h), \quad f | h \sim \mathcal{GP}(0, h * Rh).$$

$$\begin{aligned} x &\sim \mathcal{N}(0, I) \implies Ax \sim \mathcal{N}(0, AA^\top) \\ x &\sim \mathcal{GP}(0, \delta) \implies h * x \sim \mathcal{GP}(0, h * Rh) \end{aligned}$$

Model (GPCM (Tobar et al., 2015))

$$h \sim \mathcal{GP}(0, k_h), \quad f | h \sim \mathcal{GP}(0, h * Rh).$$

$$\begin{aligned} x \sim \mathcal{N}(0, I) &\implies Ax \sim \mathcal{N}(0, AA^\top) \\ x \sim \mathcal{GP}(0, \delta) &\implies h * x \sim \mathcal{GP}(0, h * Rh) \end{aligned}$$

Model (GPCM (Tobar et al., 2015), Equivalent Formulation)

$$h \sim \mathcal{GP}(0, k_h), \quad x \sim \mathcal{GP}(0, \delta), \quad f | h, x = h * x.$$

Model (GPCM (Tobar et al., 2015), Equivalent Formulation)

$$h \sim \mathcal{GP}(0, k_h), \quad x \sim \mathcal{GP}(0, \delta), \quad f | h, x = h * x.$$

Model (GPCM (Tobar et al., 2015), Equivalent Formulation)

$$h \sim \mathcal{GP}(0, k_h), \quad x \sim \mathcal{GP}(0, \delta), \quad f | h, x = h * x.$$

- Joint distribution:

$$p(f, h, \underset{\uparrow}{u}, x, \underset{\uparrow}{z}) = p(f | h, x) p(h | \textcolor{red}{u}) p(\textcolor{red}{u}) p(x | \textcolor{red}{z}) p(\textcolor{red}{z}).$$

inducing points for h and x resp.

Model (GPCM (Tobar et al., 2015), Equivalent Formulation)

$$h \sim \mathcal{GP}(0, k_h), \quad x \sim \mathcal{GP}(0, \delta), \quad f | h, x = h * x.$$

- Joint distribution:

$$p(f, h, \underset{\uparrow}{u}, x, \underset{\uparrow}{z}) = p(f | h, x) p(h | \underset{\uparrow}{u}) p(\underset{\uparrow}{u}) p(x | \underset{\uparrow}{z}) p(\underset{\uparrow}{z}).$$

inducing points for h and x resp.

- Approximate posterior:

$$q(f, h, \underset{\uparrow}{u}, x, \underset{\uparrow}{z}) = p(f | h, x) p(h | \underset{\uparrow}{u}) q(\underset{\uparrow}{u}) p(x | \underset{\uparrow}{z}) q(\underset{\uparrow}{z}).$$

- Mean-field approximate posterior:

$$q(f, h, u, x, z) = p(f | h, x)p(h | u)q(u)p(x | z)q(z).$$

- Mean-field approximate posterior:

$$q(f, h, \mathbf{u}, x, \mathbf{z}) = p(f | h, x) p(h | \mathbf{u}) q(\mathbf{u}) p(x | \mathbf{z}) q(\mathbf{z}).$$

- Structured mean-field approximate posterior:

$$q(f, h, \mathbf{u}, x, \mathbf{z}) = p(f | h, x) p(h | \mathbf{u}) p(x | \mathbf{z}) q(\mathbf{u}, \mathbf{z}).$$

- Mean-field approximate posterior:


$$q(f, h, \mathbf{u}, x, \mathbf{z}) = p(f | h, x)p(h | \mathbf{u})q(\mathbf{u})p(x | \mathbf{z})q(\mathbf{z}).$$

- Structured mean-field approximate posterior:


$$q(f, h, \mathbf{u}, x, \mathbf{z}) = p(f | h, x)p(h | \mathbf{u})p(x | \mathbf{z})q(\mathbf{u}, \mathbf{z}).$$

- MCMC to sample from q^* .


Prior over kernel:




Prior over PSD:



Prior over kernel:




Prior over PSD:



Extension: Causality


8/12




Extension: Multiple Outputs

9/12


$\mathcal{K}_{H_{1,1}}$




$\mathcal{K}_{H_{1,2}}$




Observation 1




Observation 2




$\mathcal{K}_{H_{2,1}}$




$\mathcal{K}_{H_{2,2}}$



Observation 3



Observation 4



But what about the *kernel of the kernel*?

But what about the *kernel of the kernel*?

And the *kernel of the kernel of the kernel*?

Model (N -Deep Kernel Model)

$$\begin{aligned}h_0 &\sim \mathcal{GP}(0, k_h), \\h_1 | h_0 &\sim \mathcal{GP}(0, h_0 * Rh_0), \\&\vdots \\h_N | h_{N-1} &\sim \mathcal{GP}(0, h_{N-1} * Rh_{N-1}), \\f | h_N &= h_N.\end{aligned}$$

Extension: Deep Kernel Model

12/12

