

SPECTRAL METHODS IN GAUSSIAN MODELLING

TOPIC 4: SPECTRUM ESTIMATION


James Requiema and Wessel Bruinsma

University of Cambridge and Invenia Labs


21 January 2019

- PSD is of wide interest.

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.




- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.




- Estimators:

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.




- Estimators:
 - parametric methods

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.




- Estimators:
 - parametric methods (SSA, SMK)

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.




- Estimators:
 - parametric methods (SSA, SMK),
 - nonparametric methods

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.



- Estimators:
 - parametric methods (SSA, SMK),
 - nonparametric methods (GPCM).

- PSD is of wide interest.
- Estimation difficult: samples limited, noisy, and nonuniform.



- Estimators:
 - parametric methods (SSA, SMK),
 - nonparametric methods (GPCM).
- Novel model by Tobar (2018).

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')), \quad s(\xi) \mid f = |\hat{f}(\xi)|^2.$$

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')), \quad s(\xi) \mid f = |\hat{f}(\xi)|^2.$$

- Estimator: $\hat{s}(\xi) = \mathbb{E}[s(\xi) \mid e]$.

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')), \quad s(\xi) \mid f = |\hat{f}(\xi)|^2.$$

- Estimator: $\hat{s}(\xi) = \mathbb{E}[s(\xi) \mid e]$.
- Does not work: samples from f not integrable!

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')), \quad s(\xi) \mid f = |\hat{f}(\xi)|^2.$$

- Estimator: $\hat{s}(\xi) = \mathbb{E}[s(\xi) \mid e]$.
- Does not work: samples from f not integrable!
- Windowed version of f :


$$f_w(t) \mid f = f(t)w(t).$$

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')), \quad s(\xi) | f = |\hat{f}(\xi)|^2.$$

- Estimator: $\hat{s}(\xi) = \mathbb{E}[s(\xi) | e]$.
- Does not work: samples from f not integrable!
- Windowed version of f :

$$f_w(t) | f = f(t)w(t).$$




- Effect of window:

$$\hat{f}_w(\xi) \mid f = (\hat{f} * \hat{w})(\xi).$$


- Effect of window:

$$\hat{f}_w(\xi) | f = (\hat{f} * \hat{w})(\xi).$$



- Effect of window:


$$\hat{f}_w(\xi) | f = (\hat{f} * \hat{w})(\xi).$$



- Window functions:
 - algorithmic purposes (here: ensure integrability),

- Effect of window:

$$\hat{f}_w(\xi) | f = (\hat{f} * \hat{w})(\xi).$$



- Window functions:
 - algorithmic purposes (here: ensure integrability),
 - acquisition devices (e.g., sampling: $w = \mathbb{I}$).

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')),$$

$$f_w(t) \mid f = f(t)w(t),$$

$$s_w(\xi) \mid f_w = |\hat{f}_w(\xi)|^2.$$

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')),$$

$$f_w(t) \mid f = f(t)w(t),$$

$$s_w(\xi) \mid f_w = |\hat{f}_w(\xi)|^2.$$

- \hat{f}_w called **local spectrum**.

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')),$$

$$f_w(t) \mid f = f(t)w(t),$$

$$s_w(\xi) \mid f_w = |\hat{f}_w(\xi)|^2.$$

- \hat{f}_w called **local spectrum**.
- Estimator:

$$\hat{s}_w(\xi) = \mathbb{E}[s_w(\xi) \mid e] = \mathbb{E}[(\operatorname{Re} \hat{f}_w(\xi))^2 \mid e] + \mathbb{E}[(\operatorname{Im} \hat{f}_w(\xi))^2 \mid e].$$

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')),$$

$$f_w(t) \mid f = f(t)w(t),$$

$$s_w(\xi) \mid f_w = |\hat{f}_w(\xi)|^2.$$

- \hat{f}_w called **local spectrum**.

- Estimator:

$$\hat{s}_w(\xi) = \mathbb{E}[s_w(\xi) \mid e] = \mathbb{E}[(\operatorname{Re} \hat{f}_w(\xi))^2 \mid e] + \mathbb{E}[(\operatorname{Im} \hat{f}_w(\xi))^2 \mid e].$$

- Choices:

$$\bullet \quad w(t) = \exp(-\alpha\pi^2 t^2), \quad (\text{tractability})$$

- Generative model:

$$f \sim \mathcal{GP}(0, k(t - t')),$$

$$f_w(t) \mid f = f(t)w(t),$$

$$s_w(\xi) \mid f_w = |\hat{f}_w(\xi)|^2.$$

- \hat{f}_w called **local spectrum**.

- Estimator:

$$\hat{s}_w(\xi) = \mathbb{E}[s_w(\xi) \mid e] = \mathbb{E}[(\operatorname{Re} \hat{f}_w(\xi))^2 \mid e] + \mathbb{E}[(\operatorname{Im} \hat{f}_w(\xi))^2 \mid e].$$

- Choices:

- $w(t) = \exp(-\alpha\pi^2 t^2),$ (tractability)

- $k = \text{SMK or EQ in simple cases.}$

- Notation:
 - $k_{fg}(t, t') = \mathbb{E}[f^*(t)g(t')]$ with $k_{ff} = k_f$,

- Notation:
 - $k_{fg}(t, t') = \mathbb{E}[f^*(t)g(t')]$ with $k_{ff} = k_f$,
 - $\mathcal{F}_t\{f(t)\}(\xi) = \int_{-\infty}^{\infty} f(t)e^{-2\pi i \xi t} dt$.

- Notation:
 - $k_{fg}(t, t') = \mathbb{E}[f^*(t)g(t')]$ with $k_{ff} = k_f$,
 - $\mathcal{F}_t\{f(t)\}(\xi) = \int_{-\infty}^{\infty} f(t)e^{-2\pi i \xi t} dt$.
- We desire

$$k_{\text{Re } \hat{f}_w}(\xi, \xi')$$

$$k_{y(\text{Re } \hat{f}_w)}(t, \xi)$$

$$k_{\text{Im } \hat{f}_w}(\xi, \xi')$$

$$k_{y(\text{Im } \hat{f}_w)}(t, \xi)$$

- Notation:

- $k_{fg}(t, t') = \mathbb{E}[f^*(t)g(t')] \text{ with } k_{ff} = k_f,$

- $\mathcal{F}_t\{f(t)\}(\xi) = \int_{-\infty}^{\infty} f(t)e^{-2\pi i \xi t} dt.$

- We desire

$$k_{\text{Re } \hat{f}_w}(\xi, \xi') = \frac{1}{2}(k_{\hat{f}_w}(\xi, \xi') + k_{\hat{f}_w}(\xi, -\xi')),$$

$$k_{y(\text{Re } \hat{f}_w)}(t, \xi) = \text{Re } k_{y\hat{f}_w}(t, \xi),$$

$$k_{\text{Im } \hat{f}_w}(\xi, \xi') = \frac{1}{2}(k_{\hat{f}_w}(\xi, \xi') - k_{\hat{f}_w}(\xi, -\xi')),$$

$$k_{y(\text{Im } \hat{f}_w)}(t, \xi) = \text{Im } k_{y\hat{f}_w}(t, \xi).$$

$$k_{\hat{f}_w}(\xi, \xi')$$

$$k_{\hat{f}_w}(\xi, \xi') = \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')]$$

$$\begin{aligned}k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\ &= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')]\end{aligned}$$

$$\begin{aligned}k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\&= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\&= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'),\end{aligned}$$

$$\begin{aligned}
k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\
&= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\
&= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'), \\
&= (\hat{k}(u)\delta(u+u') * \hat{r}_w(u, u'))(-\xi, \xi').
\end{aligned}$$

$$\begin{aligned}
k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\
&= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\
&= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'), \\
&= (\hat{k}(u)\delta(u+u') * \hat{r}_w(u, u'))(-\xi, \xi').
\end{aligned}$$

$$\hat{r}_w(\xi + u, \xi' + u) = \mathcal{N}(\xi - \xi'; 0, \alpha) \mathcal{N}(u; \tfrac{1}{2}(\xi + \xi'), \tfrac{1}{4}\alpha).$$

$$\begin{aligned}
 k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\
 &= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\
 &= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'), \\
 &= (\hat{k}(u)\delta(u+u') * \hat{r}_w(u, u'))(-\xi, \xi').
 \end{aligned}$$

$$\hat{r}_w(\xi + u, \xi' + u) = \mathcal{N}(\xi - \xi'; 0, \alpha) \mathcal{N}(u; \frac{1}{2}(\xi + \xi'), \frac{1}{4}\alpha).$$

$$k_{\hat{f}_w}(\xi, \xi') = \int \hat{k}(u)\delta(u+u')\hat{r}_w(-\xi-u, \xi'-u') \, du \, du'$$

$$\begin{aligned}
 k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\
 &= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\
 &= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'), \\
 &= (\hat{k}(u)\delta(u+u') * \hat{r}_w(u, u'))(-\xi, \xi').
 \end{aligned}$$

$$\hat{r}_w(\xi + u, \xi' + u) = \mathcal{N}(\xi - \xi'; 0, \alpha) \mathcal{N}(u; \frac{1}{2}(\xi + \xi'), \frac{1}{4}\alpha).$$

$$\begin{aligned}
 k_{\hat{f}_w}(\xi, \xi') &= \int \hat{k}(u)\delta(u+u')\hat{r}_w(-\xi-u, \xi'-u') \, du \, du' \\
 &= \int \hat{k}(u)\hat{r}_w(\xi+u, \xi'+u) \, du
 \end{aligned}$$

$$\begin{aligned}
 k_{\hat{f}_w}(\xi, \xi') &= \mathbb{E}[\hat{f}_w^*(\xi)\hat{f}_w(\xi')] \\
 &= \mathbb{E}[\mathcal{F}_{t,t'}\{f(t)f(t')w(t)w(t')\}(-\xi, \xi')] \\
 &= (\mathcal{F}_{t,t'}\{k(t-t')\}(u, u') * \mathcal{F}_{t,t'}\{w(t)w(t')\}(u, u'))(-\xi, \xi'), \\
 &= (\hat{k}(u)\delta(u+u') * \hat{r}_w(u, u'))(-\xi, \xi').
 \end{aligned}$$

$$\hat{r}_w(\xi + u, \xi' + u) = \mathcal{N}(\xi - \xi'; 0, \alpha)\mathcal{N}(u; \frac{1}{2}(\xi + \xi'), \frac{1}{4}\alpha).$$

$$\begin{aligned}
 k_{\hat{f}_w}(\xi, \xi') &= \int \hat{k}(u)\delta(u+u')\hat{r}_w(-\xi - u, \xi' - u') \, du \, du' \\
 &= \int \hat{k}(u)\hat{r}_w(\xi + u, \xi' + u) \, du \\
 &= \mathcal{N}(\xi - \xi'; 0, \alpha) \left(\hat{k}(u) * \mathcal{N}(u; 0, \frac{1}{4}\alpha) \right) \left(\frac{1}{2}(\xi + \xi') \right).
 \end{aligned}$$

$$k_{y\hat{f}_w}(t, \xi)$$

$$k_{y\hat{f}_w}(t, \xi) = \mathbb{E}[y^*(t)\mathcal{F}_{t'}\{f(t')w(t')\}(\xi)]$$

$$\begin{aligned}k_{y\hat{f}_w}(t, \xi) &= \mathbb{E}[y^*(t)\mathcal{F}_{t'}\{f(t')w(t')\}(\xi)] \\ &= \mathcal{F}_{t'}\{k(t-t')w(t')\}(\xi)\end{aligned}$$

$$\begin{aligned}k_{y\hat{f}_w}(t, \xi) &= \mathbb{E}[y^*(t)\mathcal{F}_{t'}\{f(t')w(t')\}(\xi)] \\&= \mathcal{F}_{t'}\{k(t-t')w(t')\}(\xi) \\&= (\hat{k}(u)e^{-2\pi\iota ut} * \mathcal{F}\{w\}(u))(\xi)\end{aligned}$$

$$\begin{aligned}k_{y\hat{f}_w}(t, \xi) &= \mathbb{E}[y^*(t)\mathcal{F}_{t'}\{f(t')w(t')\}(\xi)] \\&= \mathcal{F}_{t'}\{k(t-t')w(t')\}(\xi) \\&= (\hat{k}(u)e^{-2\pi\iota ut} * \mathcal{F}\{w\}(u))(\xi) \\&= \left(\hat{k}(u)e^{-2\pi\iota ut} * \mathcal{N}(u; 0, \tfrac{1}{2}\alpha)\right)(\xi).\end{aligned}$$

$$k_{\hat{f}_w}(\xi, \xi') = \mathcal{N}(\xi - \xi'; 0, \alpha) \left(\hat{k}(u) * \mathcal{N}(u; 0, \tfrac{1}{4}\alpha) \right) \left(\tfrac{1}{2}(\xi + \xi') \right)$$

$$k_{y\hat{f}_w}(t, \xi) = \left(\hat{k}(u) e^{-2\pi i u t} * \mathcal{N}(u; 0, \tfrac{1}{2}\alpha) \right) (\xi)$$

$$\begin{aligned}k_{\hat{f}_w}(\xi, \xi') &= \mathcal{N}(\xi - \xi'; 0, \alpha) \left(\hat{k}(u) * \mathcal{N}(u; 0, \tfrac{1}{4}\alpha) \right) \left(\tfrac{1}{2}(\xi + \xi') \right) \\ &\approx \mathcal{N}(\xi - \xi'; 0, \alpha) \hat{k}\left(\tfrac{1}{2}(\xi + \xi')\right),\end{aligned}$$

$$k_{y\hat{f}_w}(t, \xi) = \left(\hat{k}(u) e^{-2\pi i u t} * \mathcal{N}(u; 0, \tfrac{1}{2}\alpha) \right) (\xi)$$

$$\begin{aligned}
k_{\hat{f}_w}(\xi, \xi') &= \mathcal{N}(\xi - \xi'; 0, \alpha) \left(\hat{k}(u) * \mathcal{N}(u; 0, \tfrac{1}{4}\alpha) \right) \left(\tfrac{1}{2}(\xi + \xi') \right) \\
&\approx \mathcal{N}(\xi - \xi'; 0, \alpha) \hat{k}\left(\tfrac{1}{2}(\xi + \xi')\right),
\end{aligned}$$

$$\begin{aligned}
k_{y\hat{f}_w}(t, \xi) &= \left(\hat{k}(u) e^{-2\pi i u t} * \mathcal{N}(u; 0, \tfrac{1}{2}\alpha) \right) (\xi) \\
&\approx \hat{k}(\xi) e^{-2\pi i \xi t}.
\end{aligned}$$

$$\mathbb{E}[\hat{f}_w(\xi) \mid e] = \int \hat{k}(u) \left(\sum_{i=1}^N e^{-2\pi i u t_i} (K_e^{-1} e)_i \right) \mathcal{N}(u; \xi, \tfrac{1}{2}\alpha) \, du.$$

$$\mathbb{E}[\hat{f}_w(\xi) | e] = \int \hat{k}(u) \left(\sum_{i=1}^N e^{-2\pi i u t_i} (K_e^{-1} e)_i \right) \mathcal{N}(u; \xi, \tfrac{1}{2}\alpha) \, du.$$

- Interpretation: **DFT** of **whitened** observations, **weighted** by prior, then **smoothed** due to window.

$$\mathbb{E}[\hat{f}_w(\xi) | e] = \int \hat{k}(u) \left(\sum_{i=1}^N e^{-2\pi i u t_i} (K_e^{-1} e)_i \right) \mathcal{N}(u; \xi, \tfrac{1}{2}\alpha) \, du.$$

- Interpretation: **DFT** of **whitened** observations, **weighted** by prior, then **smoothed** due to window.
- If prior uninformative, $K_e \approx I$, then weighted DFT in the limit:

$$\lim_{\alpha \rightarrow 0} \mathbb{E}[\hat{f}_w(\xi) | e] \approx \hat{k}(\xi) \sum_{i=1}^N e^{-2\pi i \xi t_i} e_i.$$

- Commonly used for nonuniform data.

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Then estimates power at ξ from $f(t)$.

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Then estimates power at ξ from $f(t)$.

BNSE:

+ Probabilistic

Lomb–Scargle:

– Deterministic

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Then estimates power at ξ from $f(t)$.

BNSE:

+ Probabilistic

+ Nonparametric

Lomb–Scargle:

– Deterministic

– Parametric

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Then estimates power at ξ from $f(t)$.

BNSE:

- + Probabilistic
- + Nonparametric
- + Closed-form estimate

Lomb–Scargle:

- Deterministic
- Parametric
- Optimisation per ξ

- Commonly used for nonuniform data.
- For each ξ , LS fits a sine using least squares:

$$f(t) = A \sin(2\pi\xi t) + B \cos(2\pi\xi t).$$

- Then estimates power at ξ from $f(t)$.

BNSE:

+ Probabilistic

+ Nonparametric

+ Closed-form estimate

Lomb–Scargle:

– Deterministic

– Parametric


– Optimisation per ξ

- Gaussian prior on A and B : LS recovers BNSE in the limit.

- $k = \text{SMK}$.

- $k = \text{SMK}$.
- Trained on first, tested on second.


- $k = \text{SMK}$.
- Trained on first, tested on second.




(Figure taken from Tobar (2018).)

- BNSE (Tobar, 2018) novel nonparametric model for SE.

- BNSE (Tobar, 2018) novel nonparametric model for SE.
- Bayesian in nature: handles uncertainty automatically.




- BNSE (Tobar, 2018) novel nonparametric model for SE.
- Bayesian in nature: handles uncertainty automatically.



- Closed-form estimate of PSD.

- BNSE (Tobar, 2018) novel nonparametric model for SE.
- Bayesian in nature: handles uncertainty automatically.



- Closed-form estimate of PSD.

⇒ Can optimise to find periodicities.