Japanese Journal of Radiology (2022) 40:1006–1008
https://doi.org/10.1007/s11604-022-01342-6

CORRECTION

Correction to: Stereotactic body radiation therapy for metastatic lung metastases

Tomoki Kimura[1] · Toshiki Fujiwara[1] · Tsubasa Kameoka[1] · Yoshinori Adachi[1,2] · Shinji Kariya[1]

Published online: 26 September 2022
© The Author(s) 2022

Correction to: Japanese Journal of Radiology
https://doi.org/10.1007/s11604-022-01323-9

In the original publication, Tables 1 and 2 were published incorrectly.

The correct version of Tables 1 and 2 is given in this Correction.

The original publication has been corrected.

The original article can be found online at https://doi.org/10.1007/s11604-022-01323-9.

✉ Tomoki Kimura
tkkimura@kochi-u.ac.jp

[1] Department of Radiation Oncology, Kochi Medical School, Kochi University, Oko-cho, Nangoku-shi, KohasuKochi 783-8505, Japan

[2] Department of Radiation Oncology, Hiroshima Red Cross Hospital and Atomic-Bomb Survivors Hospital, 1-9-6 Sendamachi, Naka-ku, Hiroshima 730-8619, Japan

Table 1 SBRT for pulmonary oligometastatic disease (OMD)

Author/year	Study design	Patients	Lesions	Primary cancer (% CRC*)	Dose/fraction (Gy/fr)	Prescription	Local control	Overall survival	Toxicity grade ≧ 3
Norihisa, 2005, Japan [10]	Retrospective	34	43	26.5%	42–60 Gy/3 fr	Isocenter	90% (2y)	84.3% (2y)	3%
Rusthoven, 2009, USA [11] (USA, IN)	Phase I/II	38	63	23.7%	48- 60 Gy/3 fr	80–90% isodose	96% (2y)	39% (2y)	7.9%
Takeda, 2011, Japan [12]	Retrospective	34	44	CRC: 15 pts	50 Gy/5 fr	80% isodose	CRC 72% (2y)	N.A	3%
				Non-CRC: 19 pts			Non-CRC 94% (2y)	N.A	
Widder, 2013, Nertherlands [13]	Retrospective	42	N.A.**	73.8%	60 Gy/3 fr	N.A	94% (2y)	62% (3y)	2.4%
Comito, 2014, Italy [14]	Retrospective	41	60	100%	48–75 Gy/3-4fr	PTV D95%#	70% (3y)	58% (3y)	0%
Jung, 2015, Korea [15]	Retrospective	50	79	100%	48 Gy/4 fr (median)	85–90% isodose	70.6% (3y)	64% (3y)	0%
Rieber, 2016, Germany [16]	Retrospective	700	N.A	21.9%	3–33 Gy × 1–13 fr	88.7% isodose (median)	81.2% (2y)	54.4% (2y)	6.5%#
Agolli, 2016, Germany [17]	Retrospective	44	69	100%	23–45 Gy/1–3 fr	95% isodose	60.2% (2y)	50.8% (3y)	0%
Jingu, 2017, Japan [18]	Retrospective	93	104	100%	40–65 Gy/3–15 fr	Isocenter (83%)	65% (3y)	56% (3y)	2%
Helou, 2017, UK [19]	Prospective cohort	120	184	CRC: 101 pts	56–60 Gy/4 fr	PTV D95%	76.4% (2y)	N.A	1.7%
				Non-CRC: 83 pts	48–52 Gy/ 4 or 5 fr	PTV D95%	91.7% (2y)	N.A	
Osti, 2018, Italy [20]	Retrospective	129	166	31.7%	30 Gy/1 fr	95% isodose	80.1% (3y)	34% (3y)	7.4%
Sharma, 2018, Netherland [21]	Retrospective	206	327	57.3%	30–60 Gy/1–8 fr BED	70–90% isodose	85% (2y)	36% (2y)	2%
Berkovic, 2020, Belgium [22]	Retrospective	104	132	33.7%	20–60 Gy/3 or 5 fr	80% isodose	77.8% (3y)	72% (3y)	2%
Yamamoto, 2020, Japan [23]	Retrospective	1378	1547	25.3%	48 Gy/4 fr (median)	Isocenter (71.3%)	81.3% (3y)	60.3% (3y)	2.5%
Siva, 2021, Australia [24]	Randomized Phase II	45	69	46.7%	28 Gy/1 fr	PTV D99%#	64% (3y)	81% (3y)	5%
		45	69	46.7%	48 Gy/4 fr	PTV D99%	80% (3y)	67% (3y)	3%

*CRC: Colorectal cancer, **N.A.: not available, #Grade ≧ 2, # PTV D95%/99%: the dose covering 95%/99% of the planning target volume (PTV)

Table 2 SBRT for centrally located pulmonary oligometastatic disease (OMD)

Author/year	Study design	Patients	Lesions	Lesionsof OMD (%)	Dose/fraction (Gy/fr)	Prescription	Local control	Overall survival	Toxicity grade ≧ 3 (Grade 5)
Milano, 2009, USA [45]	Retrospective	53	63	34 (54%)	30–60 Gy/4–18 fr	80% isodose	73% (2y)	44% (2y)	N.A. (19%)
Rowe, 2012, USA [46] (USA, IN)	Retrospective	47	51	21 (41%)	50 Gy/4 fr	70–90% isodose	94% (2y)	N.A.*	13% (2%)
Davis, 2015, USA [47]	Retrospective	64	66	66 (100%)	37.5 Gy/3 fr (median)	N.A	69.8% (2y)	49.6% (2y)	0%
Lischalk, 2016, USA [48]	Retrospective	20	20	20 (100%)	35 or 40 Gy/5 fr	PTV D95%**	57.4% (2y)	40% (2y)	10% (0%)
Figlia, 2018, Italy [49]	Retrospective	39	39	13 (33%)	40–70 Gy/8–10 fr	PTV D95%	92.9% (2y)	83.9% (2y)	0%
Chang, 2018, Australia [50]	Retrospective	107	107	107 (100%)	30–50 Gy/1–3 fr	83% isodose	96.6%/95.7% (2y)#	55.1% (2y)	5.6% (2.8%)
Sharma, 2018, Netherland [21]	Retrospective	N.A	83	83 (100%)	45–60 Gy/5 fr	70–90% isodose	82% (3y)	N.A	2% (0%)
Lindberg, 2021, Sweden [51]	Phase II	65	68	14 (22%)	56 Gy/8 fr	67% isodose	83% (3y)	50% (3y)	34% (15%)

*N.A.: not available, ** PTV D95%: the dose covering 95% of the planning target volume (PTV), # 96.6% in central tumors and 95.7% in ultra-central tumors

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.