PSALTer results panel

$$S := \iiiint \Big(\tfrac{1}{4}\big(2\,a_2\,\mathscr{R}^{\alpha}{}_{\alpha}{}^{\beta}\,\mathscr{R}^{\chi}{}_{\beta\chi} + \mathscr{R}^{\alpha\beta\chi}\big(-2\,a_0\,\mathscr{R}_{\beta\chi\alpha} + 4\,\mathscr{W}_{\alpha\beta\chi}\big) + 4\,\mathscr{T}^{\alpha\beta}\,\mathscr{H}_{\alpha\beta} - a_0\,h^{\chi}{}_{\chi}\,\partial_{\beta}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\beta} + a_0\,h^{\chi}{}_{\chi}\,\partial_{\beta}\mathscr{R}^{\alpha\beta}{}_{\alpha} - 2\,a_0\,h_{\alpha\chi}\,\partial_{\beta}\mathscr{R}^{\alpha\beta\chi} + 2\,a_1\,h_{\beta\chi}\,\partial^{\chi}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\beta} - c_1\,\partial_{\beta}\mathscr{R}^{\delta}\,\partial^{\chi}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\delta} - c_1\,\partial_{\beta}\mathscr{R}^{\delta}{}_{\delta\beta}\,\partial^{\chi}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\delta} - c_1\,\partial_{\beta}\mathscr{R}^{\alpha\beta\chi}\,\partial_{\delta}\mathscr{R}_{\alpha}{}^{\delta}{}_{\chi} + 2\,c_1\,\partial^{\chi}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\beta}\,\partial_{\delta}\mathscr{R}_{\beta\chi}{}^{\delta} - c_1\,\partial_{\beta}\mathscr{R}^{\alpha\beta\chi}\,\partial_{\delta}\mathscr{R}_{\chi}{}^{\alpha}{}_{\alpha}{}^{\delta} + 2\,c_1\,\partial^{\chi}\mathscr{R}^{\alpha}{}_{\alpha}{}^{\beta}\,\partial_{\delta}\mathscr{R}_{\chi}{}^{\delta}{}_{\beta}\big)\Big)\,dz\,dy\,dx\,dt$$

Wave operator

[Matrix of wave operator components — too dense to transcribe reliably]

Saturated propagator

[Matrix of saturated propagator components — too dense to transcribe reliably]

Source constraints

Spin-parity form	Covariant form	Multiplicities
$k\,{}^{0^+}\!\mathcal{W}_{S}{}^{\parallel} + 2\,k\,{}^{0^+}\!\mathcal{W}_{S}{}^{\text{th}} - 6\,i\,{}^{0^+}\!\mathcal{T}^{\perp} \coloneqq 0$	$2\,\partial_{\beta}\partial_{\chi}\mathcal{T}^{\alpha\beta} + \partial_{\chi}\partial^{\delta}\partial_{\delta}\mathcal{W}^{\alpha\beta} \coloneqq \partial_{\chi}\partial_{\beta}\partial_{\alpha}\mathcal{W}^{\alpha\beta\chi}$	1
$k\,{}^{0^+}\!\mathcal{W}_{S}{}^{\text{t}} + 2\,i\,{}^{0^+}\!\mathcal{T}^{\perp} \coloneqq 0$	$2\,\partial_{\beta}\partial_{\chi}\mathcal{T}^{\alpha\beta} \coloneqq \partial_{\chi}\partial_{\beta}\partial_{\alpha}\mathcal{W}^{\alpha\beta\chi}$	1
$6\,k\,{}^{1^-}\!\mathcal{W}_{a}{}^{\perp} + 2\,k\,{}^{1^-}\!\mathcal{W}_{S}{}^{\text{th}\,\alpha} + k\,{}^{1^-}\!\mathcal{W}_{S}{}^{\text{t}\,\alpha} + 3\,k\,{}^{1^-}\!\mathcal{W}_{S}{}^{\text{t}\,\alpha} + 12\,i\,{}^{1^-}\!\mathcal{T}^{\perp\,\alpha} \coloneqq 0$	$4\,\partial_{\chi}\partial_{\beta}\partial^{\alpha}\mathcal{T}^{\beta\chi} + 2\,\partial_{\beta}\partial_{\delta}\partial_{\chi}\mathcal{W}^{\beta\alpha\chi} + 2\,\partial_{\delta}\partial^{\delta}\partial_{\chi}\mathcal{W}^{\alpha\beta}{}_{\beta} \coloneqq 4\,\partial_{\alpha}\partial^{\chi}\partial_{\beta}\mathcal{T}^{\alpha\beta} + 2\,\partial_{\beta}\partial_{\beta}\partial^{\alpha}\mathcal{W}^{\beta\chi\delta} + \partial_{\delta}\partial^{\delta}\partial^{\alpha}\mathcal{W}^{\beta\chi}{}_{\chi}$	3
$k\,{}^{1^-}\!\mathcal{W}_{S}{}^{\text{th}\,\alpha} - 6\,i\,{}^{1^-}\!\mathcal{T}^{\perp\,\alpha} \coloneqq k\,\big(3\,{}^{1^-}\!\mathcal{W}_{a}{}^{\perp\,\alpha} + {}^{1^-}\!\mathcal{W}_{S}{}^{\text{t}\,\alpha}\big)$	$2\,\partial_{\chi}\partial_{\beta}\partial^{\alpha}\mathcal{T}^{\beta\chi} + \partial_{\delta}\partial^{\delta}\partial_{\chi}\mathcal{W}^{\beta\alpha\chi} \coloneqq 2\,\partial_{\chi}\partial^{\chi}\partial_{\beta}\mathcal{T}^{\alpha\beta} + \partial_{\delta}\partial_{\beta}\partial^{\alpha}\mathcal{W}^{\beta\chi\delta}{}_{\chi}$	3
Total expected gauge generators:		8

Massive spectrum

Massless spectrum

Massless particle

Pole residue: $-\dfrac{p^2}{a_0} > 0$
Polarisations: 2

Unitarity conditions

$a_0 < 0$