Lagrangian density

$$\tfrac{1}{2}\,\alpha\,\partial_\beta h^{\chi}{}_{\chi}\,\partial^\beta h^{\alpha}{}_{\alpha} + \alpha\,\partial_\alpha h^{\alpha\beta}\,\partial_\chi h_\beta{}^{\chi} - \alpha\,\partial^\beta h^{\alpha}{}_{\alpha}\,\partial_\chi h_\beta{}^{\chi} - \tfrac{1}{2}\,\alpha\,\partial_\chi h_{\alpha\beta}\,\partial^\chi h^{\alpha\beta}$$

Added source term: $h^{\alpha\beta}\,\mathcal{T}_{\alpha\beta}$

Source constraints

SO(3) irreps	#
$\mathcal{T}^{\#2}_{0^+} == 0$	1
$\mathcal{T}^{\#1\alpha}_{1^-} == 0$	3
Total #:	4

	$\mathcal{T}^{\#1}_{0^+}$	$\mathcal{T}^{\#2}_{0^+}$
$\mathcal{T}^{\#1}_{0^+}$ †	$\frac{1}{\alpha k^2}$	0
$\mathcal{T}^{\#2}_{0^+}$ †	0	0

	$\mathcal{T}^{\#1}_{1^-\alpha}$
$\mathcal{T}^{\#1}_{1^-\alpha}$ †	0

	$h^{\#1}_{1^-\alpha}$
$h^{\#1}_{1^-\alpha}$ †	0

	$\mathcal{T}^{\#1}_{2^+\alpha\beta}$
$\mathcal{T}^{\#1}_{2^+}{}^{\alpha\beta}$ †	$-\frac{2}{\alpha k^2}$

	$h^{\#1}_{2^+\alpha\beta}$
$h^{\#1}_{2^+}{}^{\alpha\beta}$ †	$-\frac{\alpha k^2}{2}$

	$h^{\#1}_{0^+}$	$h^{\#2}_{0^+}$
$h^{\#1}_{0^+}$ †	αk^2	0
$h^{\#2}_{0^+}$ †	0	0



Quadratic pole

Pole residue:	$-\frac{1}{\alpha} > 0$
Polarisations:	2

Unitarity conditions

$\alpha < 0$

(No massive particles)