

Particle spectrograph

Wave operator and propagator

Source constraints		
SO(3) irreps	Fundamental fields	Multiplicities
$\tau_{0+}^{\#2} == 0$	$\partial_\beta \partial_\alpha \tau^{\alpha\beta} == 0$	1
$\tau_{0+}^{\#1} - 2 \, i \, k \, \sigma_{0+}^{\#1} == 0$	$\partial_\beta \partial_\alpha \tau^{\alpha\beta} == \partial_\beta \partial^\beta \tau^\alpha_\alpha + 2 \, \partial_\chi \partial^\chi \partial_\beta \sigma^{\alpha\beta}_\alpha$	1
$\tau_{1+}^{\#2\alpha} + 2 \, i \, k \, \sigma_{1+}^{\#2\alpha} == 0$	$\partial_\chi \partial_\beta \partial^\alpha \tau^\beta \chi == \partial_\chi \partial^\chi \partial_\beta \tau^{\alpha\beta} + 2 \, \partial_\delta \partial^\delta \partial_\chi \partial_\beta \sigma^{\alpha\beta} \chi$	3
$\tau_{1-}^{\#1\alpha} == 0$	$\partial_\chi \partial_\beta \partial^\alpha \tau^\beta \chi == \partial_\chi \partial^\chi \partial_\beta \tau^{\beta\alpha}$	3
$\tau_{1+}^{\#1\alpha\beta} + i \, k \, \sigma_{1+}^{\#2\alpha\beta} == 0$	$\partial_\chi \partial^\alpha \tau^{\beta\chi} + \partial_\chi \partial^\beta \tau^{\chi\alpha} + \partial_\chi \partial^\chi \tau^{\alpha\beta} +$ $2 \, \partial_\delta \partial_\chi \partial^\alpha \sigma^{\beta\chi\delta} + 2 \, \partial_\delta \partial^\delta \partial_\chi \sigma^{\alpha\beta} \chi ==$ $\partial_\chi \partial^\alpha \tau^{\chi\beta} + \partial_\chi \partial^\beta \tau^{\alpha\chi} +$ $\partial_\chi \partial^\chi \tau^{\beta\alpha} + 2 \, \partial_\delta \partial_\chi \partial^\beta \sigma^{\alpha\chi\delta}$	3
$\tau_{2+}^{\#1\alpha\beta} - 2 \, i \, k \, \sigma_{2+}^{\#1\alpha\beta} == 0$	$-i \, (4 \, \partial_\delta \partial_\chi \partial_\beta \partial^\alpha \tau^{\chi\delta} + 2 \, \partial_\delta \partial^\delta \partial_\beta \partial^\alpha \tau^{\chi\chi} -$ $3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\alpha \tau^{\beta\chi} - 3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\alpha \tau^{\chi\beta} -$ $3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\beta \tau^{\alpha\chi} - 3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\beta \tau^{\chi\alpha} +$ $3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\chi \tau^{\alpha\beta} + 3 \, \partial_\delta \partial^\delta \partial_\chi \partial^\chi \tau^{\beta\alpha} +$ $4 \, i \, k^\chi \, \partial_\epsilon \partial_\chi \partial_\beta \partial^\alpha \sigma^{\delta\epsilon}_\delta -$ $6 \, i \, k^\chi \, \partial_\epsilon \partial_\delta \partial_\chi \partial^\alpha \sigma^{\delta\epsilon}_\delta -$ $6 \, i \, k^\chi \, \partial_\epsilon \partial_\delta \partial_\chi \partial^\beta \sigma^{\alpha\delta\epsilon} +$ $2 \, \eta^{\alpha\beta} \, \partial_\epsilon \partial^\epsilon \partial_\delta \partial_\chi \tau^{\chi\delta} +$ $6 \, i \, k^\chi \, \partial_\epsilon \partial^\epsilon \partial_\delta \partial_\chi \sigma^{\alpha\delta\beta} +$ $6 \, i \, k^\chi \, \partial_\epsilon \partial^\epsilon \partial_\delta \partial_\chi \sigma^{\beta\delta\alpha} -$ $2 \, \eta^{\alpha\beta} \, \partial_\epsilon \partial^\epsilon \partial_\delta \partial^\chi \tau^{\chi\chi} -$ $4 \, i \, \eta^{\alpha\beta} \, k^\chi \, \partial_\phi \partial_\epsilon \partial_\chi \sigma^{\delta\epsilon}_\delta) == 0$	5
Total constraints/gauge generators:		16

Quadratic (free) action
$S == \iiint \iiint (f^{\alpha\beta} \, \tau_{\alpha\beta} + \omega^{\alpha\beta\chi} \, \sigma_{\alpha\beta\chi} +$ $\frac{1}{2} \, t_1 \, (2 \, \omega^{\alpha\iota}_\alpha \, \omega^{\theta\theta}_\theta - 4 \, \omega^{\theta\theta}_\alpha \partial_\iota f^{\alpha\iota} + 4 \, \omega^{\theta\theta}_\theta \partial^\iota f^\alpha_\alpha -$ $2 \, \partial_\iota f^\theta_\theta \partial^\iota f^\alpha_\alpha - 2 \, \partial_\iota f^{\alpha\iota} \partial_\theta f^\theta_\alpha + 4 \, \partial^\iota f^\alpha_\alpha \partial_\theta f^\theta_\iota - 2 \, \partial_\omega f^\theta_\iota \partial^\theta f^{\alpha\iota} - \partial_\omega f^\alpha_\iota \partial^\theta f^{\alpha\iota} + \partial_\theta f^\alpha_\iota \partial^\theta f^{\alpha\iota} + \partial_\theta f^\theta_\iota \partial^\theta f^{\alpha\iota} + 2 \, \omega_{\alpha\theta\iota} \, (\omega^{\alpha\iota\theta} + 2 \, \partial^\theta f^{\alpha\iota})) -$ $\frac{1}{3} \, r_1 \, (3 \, \partial_\beta \omega^{\theta\theta}_\iota \partial^\iota \omega^{\alpha\beta}_\alpha - 3 \, \partial_\iota \omega^{\theta\theta}_\beta \partial^\iota \omega^{\alpha\beta}_\alpha - 3 \, \partial_\alpha \omega^{\alpha\beta\iota} \partial_\theta \omega^{\theta\theta}_\iota +$ $6 \, \partial^\iota \omega^{\alpha\beta}_\alpha \partial_\theta \omega^{\theta\theta}_\beta + 3 \, \partial_\alpha \omega^{\alpha\beta\iota} \partial_\theta \omega^{\theta\theta}_\iota - 6 \, \partial^\iota \omega^{\alpha\beta}_\alpha \partial_\theta \omega^{\theta\theta}_\iota + 4 \, \partial_\beta \omega_{\alpha\iota\theta} \partial^\theta \omega^{\alpha\beta\iota} - 2 \, \partial_\beta \omega_{\alpha\theta\iota} \partial^\theta \omega^{\alpha\beta\iota} +$ $8 \, \partial_\beta \omega_{\iota\theta\alpha} \partial^\theta \omega^{\alpha\beta\iota} + 2 \, \partial_\iota \omega_{\alpha\beta\theta} \partial^\theta \omega^{\alpha\beta\iota} - 2 \, \partial_\theta \omega_{\alpha\beta\iota} \partial^\theta \omega^{\alpha\beta\iota} -$ $2 \, \partial_\theta \omega_{\alpha\iota\beta} \partial^\theta \omega^{\alpha\beta\iota})) [t, x, y, z] d\iota z d\iota y d\iota x d\iota t$

$\sigma_{1+}^{\#1} \dagger^{\alpha\beta}$	$\sigma_{1+}^{\#2}$	$\tau_{1+}^{\#1}$	$\sigma_{1-}^{\#1}$	$\sigma_{1-}^{\#2}$	$\tau_{1-}^{\#1}$	$\tau_{1-}^{\#2}$
0	$-\frac{\sqrt{2}}{t_1+k^2}t_1$	$-\frac{i\sqrt{2}k}{t_1+k^2}t_1$	0	0	0	0
$\sigma_{1+}^{\#2} \dagger^{\alpha\beta}$	$-\frac{\sqrt{2}}{t_1+k^2}t_1$	$-\frac{i(2k^3r_1-k t_1)}{(1+k^2)^2t_1^2}$	0	0	0	0
$\tau_{1+}^{\#1} \dagger^{\alpha\beta}$	$\frac{i\sqrt{2}k}{t_1+k^2}t_1$	$\frac{-2k^4r_1+k^2t_1}{(1+k^2)^2t_1^2}$	0	0	0	0
$\sigma_{1-}^{\#1} \dagger^\alpha$	0	0	0	$\frac{\sqrt{2}}{t_1+2k^2}t_1$	$\frac{\sqrt{2}}{t_1+2k^2}t_1$	$\frac{2ik}{t_1+2k^2}t_1$
$\sigma_{1-}^{\#2} \dagger^\alpha$	0	0	0	$\frac{\sqrt{2}}{t_1+2k^2}t_1$	0	$\frac{i\sqrt{2}k}{(1+2k^2)^2}t_1$
$\tau_{1-}^{\#1} \dagger^\alpha$	0	0	0	0	0	0
$\tau_{1-}^{\#2} \dagger^\alpha$	0	0	0	$-\frac{2ik}{t_1+2k^2}t_1$	$-\frac{i\sqrt{2}k}{(1+2k^2)^2}t_1$	$\frac{2k^2}{(1+2k^2)^2}t_1$


$\omega_{1+}^{\#1} \dagger^{\alpha\beta}$	$\omega_{1+}^{\#2}$	$f_{1+}^{\#1}$	$f_{1+}^{\#1\alpha\beta}$	$\omega_{1-}^{\#1}$	$\omega_{1-}^{\#2}$	$f_{1-}^{\#1}$	$f_{1-}^{\#2}$
$k^2r_1-\frac{t_1}{2}$	$-\frac{t_1}{\sqrt{2}}$	$-\frac{ikt_1}{\sqrt{2}}$	0	0	0	0	0
$\omega_{1+}^{\#2} \dagger^{\alpha\beta}$	$-\frac{t_1}{\sqrt{2}}$	0	0	0	0	0	0
$f_{1+}^{\#1} \dagger^{\alpha\beta}$	$\frac{ikt_1}{\sqrt{2}}$	0	0	0	0	0	0
$\omega_{1-}^{\#1} \dagger^\alpha$	0	0	0	$-\frac{t_1}{2}$	$\frac{t_1}{\sqrt{2}}$	0	$ik t_1$
$\omega_{1-}^{\#2} \dagger^\alpha$	0	0	0	$\frac{t_1}{\sqrt{2}}$	0	0	0
$f_{1-}^{\#1} \dagger^\alpha$	0	0	0	0	0	0	0
$f_{1-}^{\#2} \dagger^\alpha$	0	0	0	$-ik t_1$	0	0	0

$\sigma_{2+}^{\#1} \dagger^{\alpha\beta}$	$\tau_{2+}^{\#1}$	$\sigma_{2-}^{\#1}$
$\alpha\beta$	$\alpha\beta$	$\alpha\beta\chi$
$\frac{2}{(1+2k^2)^2}t_1$	$-\frac{2i\sqrt{2}k}{(1+2k^2)^2}t_1$	0
$\frac{2i\sqrt{2}k}{(1+2k^2)^2}t_1$	$\frac{4k^2}{(1+2k^2)^2}t_1$	0
$\alpha\beta\chi$	0	$\frac{2}{2k^2r_1+t_1}$

$\omega_{0+}^{\#1} \dagger$	$\omega_{0+}^{\#1}$	$f_{0+}^{\#1}$	$f_{0+}^{\#2}$
$\omega_{0+}^{\#1}$	$\omega_{0+}^{\#1}$	$f_{0+}^{\#1}$	$f_{0+}^{\#2}$
$-t_1$	$i\sqrt{2}kt_1$	0	0
$-i\sqrt{2}kt_1$	$-2k^2t_1$	0	0
0	0	0	0
0	0	0	$-t_1$

$\frac{t_1}{2}$	$-\frac{ikt_1}{\sqrt{2}}$	0
$\frac{ikt_1}{\sqrt{2}}$	k^2t_1	0
0	0	$k^2r_1+\frac{t_1}{2}$

Massive and massless spectra



Massive particle	
Pole residue:	$-\frac{1}{r_1} > 0$
Polarisations:	5
Square mass:	$-\frac{t_1}{2r_1} > 0$
Spin:	2
Parity:	Odd

(no massless particles)

Unitarity conditions

$r_1 < 0 \ \&\& \ t_1 > 0$