Particle spectrograph

Wave operator and propagator

Quadratic (free) action

$$S == \int\int\int\int (\beta\,(h_{\alpha\beta}\,h^{\alpha\beta} - h^{\alpha}{}_{\alpha}\,h^{\beta}{}_{\beta}) + h^{\alpha\beta}\,\mathcal{T}_{\alpha\beta} + \frac{1}{2}\,\alpha\,(\partial_{\beta}h^{\chi}{}_{\chi}\,\partial^{\beta}h^{\alpha}{}_{\alpha} + 2\,\partial_{\alpha}h^{\alpha\beta}\,\partial_{\chi}h_{\beta}{}^{\chi} - 2\,\partial^{\beta}h^{\alpha}{}_{\alpha}\,\partial_{\chi}h_{\beta}{}^{\chi} - \partial_{\chi}h_{\alpha\beta}\,\partial^{\chi}h^{\alpha\beta}))[t, x, y, z]\,dz\,dy\,dx\,dt$$

	$h^{\#1}_{0^+}$	$h^{\#2}_{0^+}$
$h^{\#1}_{0^+}\,\dagger$	$-2\,\beta + \alpha k^2$	$-\sqrt{3}\,\beta$
$h^{\#2}_{0^+}\,\dagger$	$-\sqrt{3}\,\beta$	0

(No source constraints)

	$\mathcal{T}^{\#1}_{2^+\,\alpha\beta}$
$\mathcal{T}^{\#1}_{2^+}\,\dagger^{\alpha\beta}$	$\frac{1}{\beta - \frac{\alpha k^2}{2}}$

	$h^{\#1}_{2^+\,\alpha\beta}$
$h^{\#1}_{2^+}\,\dagger^{\alpha\beta}$	$\beta - \frac{\alpha k^2}{2}$

	$\mathcal{T}^{\#1}_{1^-\,\alpha}$
$\mathcal{T}^{\#1}_{1^-}\,\dagger^{\alpha}$	$\frac{1}{\beta}$

	$h^{\#1}_{1^-\,\alpha}$
$h^{\#1}_{1^-}\,\dagger^{\alpha}$	β

	$\mathcal{T}^{\#2}_{0^+}\,\dagger$	$\mathcal{T}^{\#1}_{0^+}\,\dagger$
$\mathcal{T}^{\#1}_{0^+}$	$-\frac{1}{\sqrt{3}\,\beta}$	0
$\mathcal{T}^{\#2}_{0^+}$	$\frac{2\beta - \alpha k^2}{3\beta^2}$	$-\frac{1}{\sqrt{3}\,\beta}$

Massive and massless spectra



Massive particle	
Pole residue:	$-\frac{2}{\alpha} > 0$
Polarisations:	5
Square mass:	$\frac{2\,\beta}{\alpha} > 0$
Spin:	2
Parity:	Even

(No massless particles)

Unitarity conditions

$\alpha < 0$ && $\beta < 0$