
PSALTer results panel

$$S = \iiint (h^{\alpha\beta} \mathcal{T}_{\alpha\beta} + \frac{1}{2} \alpha_2 \partial_\beta h^\chi_{\chi} \partial^\beta h^\alpha_{\alpha} + \alpha_1 (\partial_\alpha h^{\alpha\beta} \partial_\chi h^\chi_{\beta} - \partial^\beta h^\alpha_{\alpha} \partial_\chi h^\chi_{\beta} - \frac{1}{2} \partial_\chi h_{\alpha\beta} \partial^\chi h^{\alpha\beta}))[t, \, x, \, y, \, z] \, dz \, dy \, dx \, dt$$

Saturated propagator




Source constraints

Spin-parity form	Covariant form	Multiplicities
${}^{1}\mathcal{T}^{\perp}{}^{\alpha}=0$	$\partial_{\chi}\partial_{\beta}\partial^{\alpha}\mathcal{T}^{\beta\chi} == \partial_{\chi}\partial^{\chi}\partial_{\beta}\mathcal{T}^{\alpha\beta}$	3
Total expected gauge generators:		3

Massive spectrum

(No particles)

Massless spectrum



Massless particle

Massless particle

Pole residue:
$$\frac{p^2}{\frac{-\alpha_1 + \alpha_2}{1}} > 0$$
Polarisations: 1

Pole residue:	$-\frac{p^2}{\alpha_1} > 0$
Polarisations:	2

Unitarity conditions

$$\alpha_{.} < 0 \&\& \alpha_{.} > \alpha_{.}$$