

# Particle spectrograph

## Wave operator and propagator

Quadratic (free) action

$S_F ==$

$$\iiint (\beta h_{\alpha\beta} h^{\alpha\beta} - \gamma h^\alpha_\alpha h^\beta_\beta + h^{\alpha\beta} \mathcal{T}_{\alpha\beta} + \frac{1}{2} \alpha \partial_\beta h^\chi_\chi \partial^\beta h^\alpha_\alpha + \alpha \partial_\alpha h^{\alpha\beta} \partial_\chi h^\chi_\beta - \alpha \partial^\beta h^\alpha_\alpha \partial_\chi h^\chi_\beta - \frac{1}{2} \alpha \partial_\chi h_{\alpha\beta} \partial^\chi h^{\alpha\beta}) [t, x, y, z] dz dy dx dt$$

	$\mathcal{T}_{0+}^{\#1}$	$\mathcal{T}_{0+}^{\#2}$
$\mathcal{T}_{0+}^{\#1} \dagger$	$\frac{1}{\frac{\beta(\beta-4)\gamma}{\beta-\gamma} + \alpha k^2}$	$\frac{\sqrt{3}\gamma}{\beta(\beta-4\gamma) + \alpha(\beta-\gamma)k^2}$
$\mathcal{T}_{0+}^{\#2} \dagger$	$\frac{\sqrt{3}\gamma}{\beta(\beta-4\gamma) + \alpha(\beta-\gamma)k^2}$	$\frac{1}{\beta + \gamma(-1 - \frac{3\gamma}{\beta-3\gamma + \alpha k^2})}$

	$h_{0+}^{\#1}$	$h_{0+}^{\#2}$
$h_{0+}^{\#1} \dagger$	$\beta - 3\gamma + \alpha k^2$	$-\sqrt{3}\gamma$
$h_{0+}^{\#2} \dagger$	$-\sqrt{3}\gamma$	$\beta - \gamma$

(No source constraints)


	$h_{1-}^{\#1} \alpha$
$h_{1-}^{\#1} \dagger \alpha$	$\beta$

	$\mathcal{T}_{2+}^{\#1} \alpha \beta$
$\mathcal{T}_{2+}^{\#1} \dagger \alpha \beta$	$\frac{1}{\beta - \frac{\alpha k^2}{2}}$

	$h_{2+}^{\#1} \alpha \beta$
$h_{2+}^{\#1} \dagger \alpha \beta$	$\beta - \frac{\alpha k^2}{2}$

	$\mathcal{T}_{1-}^{\#1} \alpha$
$\mathcal{T}_{1-}^{\#1} \dagger \alpha$	$\frac{1}{\beta}$


## Massive and massless spectra



Massive particle

Pole residue:	$\frac{\beta^2 - 2\beta\gamma + 4\gamma^2}{\alpha(\beta-\gamma)^2} > 0$
Polarisations:	1
Square mass:	$-\frac{\beta(\beta-4\gamma)}{\alpha(\beta-\gamma)} > 0$
Spin:	0
Parity:	Even

(No massless particles)



Massive particle	
Pole residue:	$-\frac{2}{\alpha} > 0$
Polarisations:	5
Square mass:	$\frac{2\beta}{\alpha} > 0$
Spin:	2
Parity:	Even

## Unitarity conditions

(Unitarity is demonstrably impossible)