$$
\begin{array}{c|cc|cc|c|c}
 & \mathcal{K}^{\#1}_{0^+} & \mathcal{K}^{\#2}_{0^+} & \mathcal{K}^{\#1}_{1^-\,\alpha} & \mathcal{K}^{\#2}_{1^-\,\alpha} & \mathcal{K}^{\#1}_{2^+\,\alpha\beta} & \mathcal{K}^{\#1}_{3^-\,\alpha\beta\chi} \\
\hline
\mathcal{K}^{\#1}_{0^+}\dagger & \Upsilon_1 k^2 & \frac{\Upsilon_2 k^2}{2} & & & & \\
\mathcal{K}^{\#2}_{0^+}\dagger & \frac{\Upsilon_2 k^2}{2} & \Upsilon_1 k^2 & & & & \\
\mathcal{K}^{\#1}_{1^-}\dagger^{\alpha} & & & \frac{5 k^2\, {}^{(4)}\kappa_2}{6} & -\frac{1}{6}\sqrt{5}\, k^2\, {}^{(4)}\kappa_2 & & \\
\mathcal{K}^{\#2}_{1^-}\dagger^{\alpha} & & & -\frac{1}{6}\sqrt{5}\, k^2\, {}^{(4)}\kappa_2 & \frac{k^2\, {}^{(4)}\kappa_2}{6} & & \\
\mathcal{K}^{\#1}_{2^+}\dagger^{\alpha\beta} & & & & & 0 & \\
\mathcal{K}^{\#1}_{3^-}\dagger^{\alpha\beta\chi} & & & & & & -\frac{3 k^2\, {}^{(4)}\kappa_2}{2}
\end{array}
$$

$$
\begin{array}{c|cc|cc|c|c}
 & \mathcal{J}^{\#1}_{0^+} & \mathcal{J}^{\#2}_{0^+} & \mathcal{J}^{\#1}_{1^-\,\alpha} & \mathcal{J}^{\#2}_{1^-\,\alpha} & \mathcal{J}^{\#1}_{2^+\,\alpha\beta} & \mathcal{J}^{\#1}_{3^-\,\alpha\beta\chi} \\
\hline
\mathcal{J}^{\#1}_{0^+}\dagger & \frac{\Upsilon_1 k^2}{\mathrm{Det}(0^+)} & -\frac{\Upsilon_2 k^2}{2\,\mathrm{Det}(0^+)} & & & & \\
\mathcal{J}^{\#2}_{0^+}\dagger & -\frac{\Upsilon_2 k^2}{2\,\mathrm{Det}(0^+)} & \frac{\Upsilon_1 k^2}{\mathrm{Det}(0^+)} & & & & \\
\mathcal{J}^{\#1}_{1^-}\dagger^{\alpha} & & & \frac{5}{6 k^2\, {}^{(4)}\kappa_2} & -\frac{\sqrt{5}}{6 k^2\, {}^{(4)}\kappa_2} & & \\
\mathcal{J}^{\#2}_{1^-}\dagger^{\alpha} & & & -\frac{\sqrt{5}}{6 k^2\, {}^{(4)}\kappa_2} & \frac{1}{6 k^2\, {}^{(4)}\kappa_2} & & \\
\mathcal{J}^{\#1}_{2^+}\dagger^{\alpha\beta} & & & & & 0 & \\
\mathcal{J}^{\#1}_{3^-}\dagger^{\alpha\beta\chi} & & & & & & -\frac{2}{3 k^2\, {}^{(4)}\kappa_2}
\end{array}
$$

Abbreviations used in matrices

$\Upsilon_1 == {}^{(4)}\kappa_1 + {}^{(4)}\kappa_2$ && $\Upsilon_2 == 2\,{}^{(4)}\kappa_1 - {}^{(4)}\kappa_2$ && $\mathrm{Det}(0^+) == \frac{3}{4} k^4\, {}^{(4)}\kappa_2 (4\,{}^{(4)}\kappa_1 + {}^{(4)}\kappa_2)$

Lagrangian

$${}^{(4)}\kappa_1 \partial_\beta \mathcal{K}_\chi{}^\delta{}_\delta \partial^\chi \mathcal{K}^\alpha{}_\alpha{}^\beta + {}^{(4)}\kappa_2 \partial_\chi \mathcal{K}_\beta{}^\delta{}_\delta \partial^\chi \mathcal{K}^\alpha{}_\alpha{}^\beta + \frac{9}{2}\, {}^{(4)}\kappa_2 \partial_\alpha \mathcal{K}^{\alpha\beta\chi} \partial_\delta \mathcal{K}_{\beta\chi}{}^\delta - 3\, {}^{(4)}\kappa_2 \partial^\chi \mathcal{K}^\alpha{}_\alpha{}^\beta \partial_\delta \mathcal{K}_{\beta\chi}{}^\delta - \frac{3}{2}\, {}^{(4)}\kappa_2 \partial_\delta \mathcal{K}_{\alpha\beta\chi} \partial^\delta \mathcal{K}^{\alpha\beta\chi}$$

Added source term(s):	$\mathcal{K}^{\alpha\beta\chi}\ \mathcal{J}_{\alpha\beta\chi}$		
Source constraint(s)	# constraint(s)	Covariant form	
$\mathcal{J}^{\#2\,\alpha}_{1^-} + \mathcal{J}^{\#1\,\alpha}_{1^-} == 0$	3	$\partial_\chi \partial^\alpha \mathcal{J}^\beta{}_\beta{}^\chi == \partial_\chi \partial^\chi \mathcal{J}^{\alpha\beta}{}_\beta$	
$\mathcal{J}^{\#1\,\alpha\beta}_{2^+} == 0$	5	$k^\chi\, (\partial_\epsilon \partial_\chi \partial^\beta \partial^\alpha \mathcal{J}^\delta{}_\delta{}^\epsilon + 2\, \partial_\epsilon \partial_\delta \partial^\beta \partial^\alpha \mathcal{J}_\chi{}^{\delta\epsilon} - 3\, \partial_\epsilon \partial_\delta \partial_\chi \partial^\alpha \mathcal{J}^{\beta\delta\epsilon} - 3\, \partial_\epsilon \partial_\delta \partial_\chi \partial^\beta \mathcal{J}^{\alpha\delta\epsilon} + 3\, \partial_\epsilon \partial^\epsilon \partial_\delta \partial_\chi \mathcal{J}^{\alpha\beta\delta} + \eta^{\alpha\beta}\, \partial_\phi \partial_\epsilon \partial_\delta \partial_\chi \mathcal{J}^{\delta\epsilon\phi} - \eta^{\alpha\beta}\, \partial_\phi \partial^\phi \partial_\epsilon \partial_\chi \mathcal{J}^\delta{}_\delta{}^\epsilon) == 0$	
Total # constraint(s):	8		
Resolved pole(s)	# polarization(s)	Square mass	Residue
	2	0	$\frac{1}{{}^{(4)}\kappa_2}$
Resolved unitarity condition(s):		${}^{(4)}\kappa_2 > 0$	