



$$\begin{array}{cc}
 & \mathcal{T}_{1^+}^{\#1}{}_{\alpha\beta} \quad \mathcal{T}_{1^-}^{\#1}{}_{\alpha} \\
 \mathcal{T}_{1^+}^{\#1}{}_{\dagger^{\alpha\beta}} & \begin{array}{|c|c|} \hline \alpha k^2 & 0 \\ \hline \end{array} \\
 \mathcal{T}_{1^-}^{\#1}{}_{\dagger^{\alpha}} & \begin{array}{|c|c|} \hline 0 & -\alpha k^2 \\ \hline \end{array}
 \end{array}$$

$$\begin{array}{cc}
 & \sigma_{1^+}^{\#1}{}_{\alpha\beta} \quad \sigma_{1^-}^{\#1}{}_{\alpha} \\
 \sigma_{1^+}^{\#1}{}_{\dagger^{\alpha\beta}} & \begin{array}{|c|c|} \hline \frac{1}{\alpha k^2} & 0 \\ \hline \end{array} \\
 \sigma_{1^-}^{\#1}{}_{\dagger^{\alpha}} & \begin{array}{|c|c|} \hline 0 & -\frac{1}{\alpha k^2} \\ \hline \end{array}
 \end{array}$$

Lagrangian			
$\alpha \partial_\rho \mathcal{T}_{\mu\nu} \partial^\rho \mathcal{T}^{\mu\nu} - 4 \alpha \partial_\rho \mathcal{T}^{\rho\nu} \partial_\sigma \mathcal{T}^\sigma{}_\nu$			
Added source term(s):	$\mathcal{T}^{\alpha\beta} \sigma_{\alpha\beta}$		
Resolved pole(s)	# polarization(s)	Square mass	Residue
	2	0	$\frac{1}{\alpha}$
	2	0	$\frac{p^2}{\alpha}$
Resolved unitarity condition(s):		$\alpha > 0$	