





$\mathcal{K}_{0^+}^{\#1} \uparrow$	$\mathcal{K}_{0^+}^{\#1} \uparrow^{\alpha\beta\chi}$	$\mathcal{K}_{1^+}^{\#1} \alpha\beta$	$\mathcal{K}_{1^+}^{\#2} \alpha\beta$	$\mathcal{K}_{1^+}^{\#1} \alpha$	$\mathcal{K}_{1^+}^{\#2} \alpha$	$\mathcal{K}_{2^+}^{\#1} \alpha\beta$	$\mathcal{K}_{2^+}^{\#1} \alpha\beta\chi$
$\mathcal{K}_{0^+}^{\#1} \uparrow$	0	0					
$\mathcal{K}_{0^+}^{\#1} \uparrow^{\alpha\beta\chi}$	0	$\Upsilon_1 k^2$					
$\mathcal{K}_{1^+}^{\#1} \uparrow^{\alpha\beta}$	$\frac{1}{18} (-\Upsilon_2 k^2 + 9 \binom{(2)}{K_1})$	$\frac{\Upsilon_2 k^2 - 9 \binom{(2)}{K_1}}{9 \sqrt{2}}$	0	0			
$\mathcal{K}_{1^+}^{\#2} \uparrow^{\alpha\beta}$	$\frac{\Upsilon_2 k^2 - 9 \binom{(2)}{K_1}}{9 \sqrt{2}}$	$\frac{1}{9} (-\Upsilon_2 k^2 + 9 \binom{(2)}{K_1})$	0	0			
$\mathcal{K}_{1^+}^{\#1} \uparrow^\alpha$	0	0	$\frac{1}{2} (\Upsilon_3 k^2 + \binom{(2)}{K_1})$	$\frac{\Upsilon_3 k^2 + \binom{(2)}{K_1}}{\sqrt{2}}$			
$\mathcal{K}_{1^+}^{\#2} \uparrow^\alpha$	0	0	$\frac{\Upsilon_3 k^2 + \binom{(2)}{K_1}}{\sqrt{2}}$	$\Upsilon_3 k^2 + \binom{(2)}{K_1}$			
			$\mathcal{K}_{2^+}^{\#1} \uparrow^{\alpha\beta}$	$\frac{1}{2} (3 \Upsilon_4 k^2 + 3 \binom{(2)}{K_1})$	0		
			$\mathcal{K}_{2^+}^{\#1} \uparrow^{\alpha\beta\chi}$	0	$\frac{1}{2} (\Upsilon_5 k^2 + 3 \binom{(2)}{K_1})$		

$\mathcal{J}_{0^+}^{\#1} \uparrow$	$\mathcal{J}_{0^+}^{\#1} \uparrow^{\alpha\beta\chi}$	$\mathcal{J}_{1^+}^{\#1} \alpha\beta$	$\mathcal{J}_{1^+}^{\#2} \alpha\beta$	$\mathcal{J}_{1^+}^{\#1} \alpha$	$\mathcal{J}_{1^+}^{\#2} \alpha$	$\mathcal{J}_{2^+}^{\#1} \alpha\beta$	$\mathcal{J}_{2^+}^{\#1} \alpha\beta\chi$
$\mathcal{J}_{0^+}^{\#1} \uparrow$	0	0					
$\mathcal{J}_{0^+}^{\#1} \uparrow^{\alpha\beta\chi}$	0	$\frac{1}{k^2 \binom{(4)}{K_{15}} - \binom{(4)}{K_{16}}}$					
$\mathcal{J}_{1^+}^{\#1} \uparrow^{\alpha\beta}$	$\frac{1}{3 \text{Det}(1^+)}$	$-\frac{\sqrt{2}}{3 \text{Det}(1^+)}$	0	0			
$\mathcal{J}_{1^+}^{\#2} \uparrow^{\alpha\beta}$	$-\frac{\sqrt{2}}{3 \text{Det}(1^+)}$	$\frac{2}{3 \text{Det}(1^+)}$	0	0			
$\mathcal{J}_{1^+}^{\#1} \uparrow^\alpha$	0	0	$\frac{1}{3 \text{Det}(1^+)}$	$\frac{\sqrt{2}}{3 \text{Det}(1^+)}$			
$\mathcal{J}_{1^+}^{\#2} \uparrow^\alpha$	0	0	$\frac{\sqrt{2}}{3 \text{Det}(1^+)}$	$\frac{2}{3 \text{Det}(1^+)}$			
			$\mathcal{J}_{2^+}^{\#1} \uparrow^{\alpha\beta}$	$\frac{1}{\text{Det}(2^+)}$	0		
			$\mathcal{J}_{2^+}^{\#1} \uparrow^{\alpha\beta\chi}$	0	$\frac{1}{\text{Det}(2^+)}$		

Abbreviations used in matrices

$\Upsilon_1 = \binom{(4)}{K_{15}} - \binom{(4)}{K_{16}}$ & $\Upsilon_2 = 3 \binom{(4)}{K_1} - 10 \binom{(4)}{2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}}} + 21 \binom{(4)}{K_2}$ & $\Upsilon_3 = 2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}} - 2 \binom{(4)}{K_2}$ & $\Upsilon_4 = \binom{(4)}{K_1} + \binom{(4)}{K_2}$ & $\Upsilon_5 = 2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}}$ & $\text{Det}(1^+) = \frac{1}{6} (k^2 (-3 \binom{(4)}{K_1} + 20 \binom{(4)}{K_{15}} + 10 \binom{(4)}{K_{16}} - 21 \binom{(4)}{K_2}) + 9 \binom{(2)}{K_1})$ & $\text{Det}(2^+) = \frac{3}{2} (k^2 (\binom{(4)}{K_1} + \binom{(4)}{K_2}) + \binom{(2)}{K_1})$

Added source term(s):	$\mathcal{K}^{\alpha\beta\chi} \mathcal{J}_{\alpha\beta\chi}$		
Source constraint(s)	# constraint(s)	Covariant form	
$\mathcal{J}_{0^+}^{\#1} = 0$	1	$\partial_\beta \mathcal{J}^{\alpha\beta} = 0$	
$\mathcal{J}_{1^+}^{\#1\alpha} = \mathcal{J}_{1^+}^{\#2\alpha}$	3	$\partial_\chi \partial^\alpha \mathcal{J}^{\beta\chi} + \partial_\chi \partial^\chi \mathcal{J}^{\beta\alpha} = 0$	
$2 \mathcal{J}_{1^+}^{\#1\alpha\beta} + \mathcal{J}_{1^+}^{\#2\alpha\beta} = 0$	3	$\partial_\chi \mathcal{J}^{\alpha\beta\chi} + \partial_\chi \mathcal{J}^{\chi\alpha\beta} = \partial_\chi \mathcal{J}^{\beta\alpha\chi}$	
Total # constraint(s):	7		
Resolved pole(s)	# polarization(s)	Square mass	Residue
	3	$\frac{9 \binom{(2)}{K_1}}{3 \binom{(4)}{K_1} - 10 \binom{(4)}{2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}}} + 21 \binom{(4)}{K_2}}$	$-\frac{6}{3 \binom{(4)}{K_1} - 20 \binom{(4)}{K_{15}} - 10 \binom{(4)}{K_{16}} + 21 \binom{(4)}{K_2}}$
	3	$-\frac{\binom{(2)}{K_1}}{2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}} - 2 \binom{(4)}{K_2}}$	$-\frac{2}{3 (2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}} - 2 \binom{(4)}{K_2})}$
	5	$-\frac{\binom{(2)}{K_1}}{\binom{(4)}{K_1} + \binom{(4)}{K_2}}$	$\frac{2}{3 (\binom{(4)}{K_1} + \binom{(4)}{K_2})}$
	5	$-\frac{3 \binom{(2)}{K_1}}{2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}}}$	$-\frac{2}{2 \binom{(4)}{K_{15}} + \binom{(4)}{K_{16}}}$
Resolved unitarity condition(s):		(Demonstrably impossible)	