


Color-blindness of direction-selective units in the zebrafish optic tectum


Alexander Wendt, Patrick Weygoldt

Supervisor: Aristides Arrenberg, Tim Hladnik, David Burkardt



Introduction


Chromaticity has a big influence on motion vision in zebrafish. Orger and Baier (2004) used the opto-motor response to show that combinations of green and red can be used to null the motion perception.



But little is known about the midbrain structures conveying the „color-motion“ perception. We investigated the activity of direction-selective units in the optic tectum of zebrafish in response to gratings of various color contrasts using a combination of two-photon microscopy and calcium imaging.


Preprocessing:

1. Region of Interests (ROI): corresponds to neurons with genetically encoded calcium indicators. The fluorescence F of the calcium imaging is calculated from the change of luminance normalized to the average luminance $F = \frac{\Delta F}{\langle F \rangle}$.




2. Active ROIs:






Strongly autocorrelated ROIs across stimulus repeats are „responding“.



3. Direction-selective ROIs: ROIs that correlated with a direction regressor (clockwise cw or counter-clockwise ccw).




2-photon calcium imaging



- ROIs responded to moving gratings of red and green
- Both directions cw and ccw responded equally
- If both red and green had the same contrast the response was suppressed indicating color blindness
- A slight shift in the troughs of activity could be explained by a higher intensity of green compared to red.

Behavior



- The behavioral response (OKR) reflect the pattern shown in calcium activity.

Conclusion

- We observed that the optic tectum lack the response to red-green merged out-of-phase stimuli and is therefore color-blind