

Understanding diabetes heterogeneity: key steps towards precision medicine in diabetes

Richard David Leslie^{1*}, FAoP. Professor of Diabetes and Autoimmunity

Ronald Ching Wan Ma^{2,3,4*}, FRCP. Professor

Paul W Franks^{5,6,7,8}, PhD. Professor

Kristen J Nadeau⁹, MD. Professor of Paediatrics

Ewan R Pearson¹⁰, FRCP. Professor of Diabetic Medicine

Maria Jose Redondo¹¹, MD. Professor of Paediatrics

*These authors should be considered joint first authors

¹Blizard Institute, Barts and Royal London Medical School, London, UK

²Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Prince of Wales Hospital, Hong Kong SAR, China;

³Chinese University of Hong Kong-Shanghai Jiao Tong University Joint Research Centre in Diabetes Genomics and Precision Medicine, Hong Kong Institute of Diabetes and Obesity, The Chinese University of Hong Kong, Hong Kong SAR, China;

⁴Laboratory for Molecular Epidemiology in Diabetes, Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong SAR, China

⁵Novo Nordisk Foundation, Tuborg Havnevej 19, 2900 Hellerup, Denmark

⁶Lund University Diabetes Centre, Department of Clinical Sciences, Lund University, Malmo, Sweden.

⁷Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine, ⁸University of Oxford, Oxford, UK.

⁸Harvard T.H. Chan School of Public Health, Boston, MA, USA

⁹University of Colorado Denver - Anschutz Medical Campus

¹⁰Population Health & Genomics, School of Medicine, University of Dundee

¹¹Texas Children's Hospital, Baylor College of Medicine

Correspondence:

Richard David Leslie (r.d.g.leslie@qmul.ac.uk)

32 Word count: 5516

33 Tables: 1

34 Figures: 5

35 Panel: 1

36 References: 107

37

38

39 **Summary**

40 Diabetes is highly heterogeneous yet diagnosed by measuring a single blood-borne metabolite,
41 glucose, irrespective of aetiology. Whilst pragmatically helpful, disease classification can become a
42 burden, limiting advances in research and medical care. Here, we describe diabetes heterogeneity,
43 highlighting recent approaches that could facilitate management by integrating three disease
44 models across all forms of diabetes, namely, the 'Palette Model', 'Threshold Model' and 'Gradient
45 Model'. Since progression to diabetes, further worsening of established diabetes and the
46 subsequent emergence of diabetes complications are kept in check by multiple processes designed
47 to prevent or circumvent metabolic dysfunction, the impact of any given disease risk factor will vary
48 from person to person depending on their background diabetes-related propensity and
49 environmental exposures. Defining the consequent heterogeneity within diabetes, both in terms
50 of diabetes risk and risk of complications, should improve health outcomes today, and shine a light
51 on avenues for novel therapy in the future.

Search strategy and selection criteria

We searched PubMed for published articles from Jan 1, 2010 to December 7, 2022, with the search terms including: “heterogeneity AND diabetes”, “classification AND diabetes”, “autoantibodies AND diabetes”, “phenotyping AND diabetes”, “diabetes subtypes”. We also searched the reference lists of articles identified by this search strategy and selected those we judged to be relevant. Review articles are cited in several instances to provide readers with more details and more references than we had space to cite.

Introduction

Diabetes carries a substantial burden, affecting approximately >500 million people worldwide (1), at a global cost estimated at nearly 1 trillion USD (2). That burden is expected to increase in the coming years, particularly in Africa, the Middle East, South Asia, and Latin America (1).

The pathophysiology and clinical phenotype of diabetes is highly heterogeneous. It is a disease diagnosed solely by an elevation in blood glucose, often with little regard for the aetiological processes that caused it. Some types of diabetes, such as type 1 diabetes (T1D) or monogenic diabetes, are often adequately defined by their distinct aetiologies; others, most notably type 2 diabetes (T2D), are not. Type 1 diabetes is defined by the autoimmune destruction of the beta-cells and accounts for ~2-10% of all diabetes cases, with its prevalence varying from one population to the next. The most common form of diabetes, T2D, accounts for 90-95% of all cases. Unlike other forms of diabetes, most cases of T2D are diagnosed through a process of exclusion, with other known explanations for chronic hyperglycaemia having been considered and rejected. Thus, the precise pathophysiology of T2D in any given case is usually unknown, and there is substantial variation in clinical presentation, treatment requirements and prognosis. When the aperture is widened to the global scale, such uncertainty is even more striking, given different diabetes phenotypes across ethnicities and geographic locations (3, 4).

Challenges of applying classification to a heterogeneous disease

In clinical medicine, disease classification is often applied to categorise different diseases to facilitate making decisions on clinical care, and to stimulate epidemiological research. The rationale for this approach is that medical practice is structured around establishing categorical diagnoses to guide management. When therapeutic options are limited, such an approach is reasonable as was the case with insulin therapy in 1922. However, when there is a range of options, as is now the case, disease classification can become an obstacle in recognizing heterogeneity, even within a given category (Table 1) (5). For example, individuals with a monogenic mutation associated with maturity onset of diabetes of the young (MODY) can show different phenotypes when carrying that mutation on different polygenic backgrounds, so that someone with Glucokinase (GCK)-MODY (usually not requiring medical intervention) can show progressive hyperglycaemia when carrying additional genetic variants for T2D or with the development of adiposity-induced insulin resistance. Here we will focus on the heterogeneity of diabetes in general, not monogenic diabetes, but this example

highlights how it may be inappropriate to simplify complex interactions that contribute to common diseases like diabetes.

Despite these limitations, disease classification does facilitate clinical diagnosis, prescribing practices, as well as global comparisons of health-related data. Growing appreciation of diabetes heterogeneity prompted an update in 2019 of the previous World Health Organization (WHO) classification of diabetes. Table 1 summarizes the current classification (6), as well as some important considerations when considering the diagnosis. For example, monogenic forms of diabetes would need to be considered in someone presenting with youth-onset diabetes, especially if there is strong family history of diabetes. Conversely, among individuals with weight gain, other endocrinopathies such as Cushing's syndrome may need to be considered.

Beta-cell destruction or dysfunction is the key pathophysiological defect underpinning most cases and forms of diabetes (7, 8). By inference, it may not be possible or necessary to group people with diabetes into specific categories. However, recognizing the 'risk-continuum' of pathophysiological processes that contribute to hyperglycaemia can provide a better guide to our understanding of diabetes, and with that, better prediction and treatment. Future approaches to disease classification should be based on defining categories with shared pathophysiology, so that the numbers of "subtypes" would be variable, and dependent on data quality and the statistical methodology being used for the clustering.

Several emerging approaches to diabetes classification have been adopted, including the integration of the "Palette Model" and the "Threshold Model" (9). Here, we will use these models, and add a third, the "Gradient Model", to add the dimension of severity of common chronic diseases, including diabetes (Figure 1). The Palette Model identifies a wide range of disease-risk factors (as with different colours on an artist's palette), contributing to disease in different ways in different individuals, such as islet autoimmunity and insulin resistance in the same person. The Threshold Model, on the other hand, highlights how a combination of diabetogenic factors, which may or may not cross the threshold for clinical diabetes (9) (10)(Figure 2). For example, an individual with mild islet autoimmunity may only develop clinical diabetes in the presence of T2D-associated genes (11) or insulin resistance (12). As a result, the disease phenotype could reflect a combination of diabetes-related mechanisms (13) which interact to modify the disease's natural course (14) (15). This dynamic is captured in the Gradient Model, allied to the Threshold Model, given that disease risk factors operate within time-specific windows, so that the presence and/or severity of diabetes, after disease-onset, may change, e.g., individuals with T2D can remit when insulin sensitivity improves and individuals with T1D can have a 'honeymoon' off insulin therapy. Moreover, not all individuals at high genetic risk of either T1D or T2D will develop islet autoimmunity or metabolic dysfunction respectively, while not everyone with islet autoimmunity or metabolic dysfunction develops diabetes. By implication, these three Models highlight the importance of allostasis through regulatory pathways, be they immune or metabolic, to reduce disease progression. Whilst none of these conceptual models alone can fully account for the complexity of diabetes, each model does offer a better appreciation of that heterogeneity, which underlies diabetes.

Heterogeneity in diabetes – biological variation or error?

Understanding diabetes heterogeneity should enhance our ability to predict disease incidence, treatment response, and prognosis. Ideally, a physician would want precisely defined, long-term (sometimes lifelong) measures of those environmental and biological features causing the disease and its subsequent trajectory. Instead, current data is sparse and imperfectly characterised i.e., often with considerable systematic [bias] and non-systematic error. Clinically, heterogeneity in diabetes is a consequence of variations either in 'biological' processes underlying the disease phenotype (signal) or error in the 'quantification' of that phenotype (noise). Decoding heterogeneity requires that signal to be successfully parsed from noise.

Many established environmental features drive diabetogenic exposures including sedentary behaviour, certain diets, stress, sleep disturbance, and smoking. Accurate and precise quantification of these exposures may help resolve some heterogeneity within diabetes, yet doing so is challenging, as such exposures are often complex, difficult to assess without systematic and non-systematic error, and conveyed over extended periods from birth through adulthood (16-18). The impact of biological variants and diabetogenic exposures on disease-related cellular networks and phenotypes have only recently been explored. These 'biological variants' include variations in DNA, epigenetic marks, RNA, proteins, hormones, and metabolites. The process of quantification of these variants is subject to technical variation and this may mislead interpretation of their impact on diabetes heterogeneity. Variations in the assessment of blood glucose also exacerbate diabetes heterogeneity: diabetes is typically diagnosed using random, fasting and post-challenge blood glucose measurements, or with HbA1c, each reflecting different features of glycaemic regulation. Fasting glucose, for example, is influenced mainly by gluconeogenesis and glycogenolysis, whereas post-challenge glycaemia is mainly determined by peripheral insulin resistance and the beta-cell response (19) (20). Whilst insulin resistance is a key pathophysiological defect in T2D (21), recovery of first phase insulin secretion identifies subjects who have remission of T2D following weight loss with reduction in hepatic and pancreatic fat (22), highlighting the key role of beta-cell function in the development of T2D. Using few point-estimates of time-varying biological features, such as glycaemia and/or autoantibodies, are also error prone owing to a 'regression dilution effect' (23). There are additional contextual factors (e.g. stress, ambient temperature, recent exercise) that limit test reproducibility and variability in sample handling and storage that can introduce error into assessments and obstruct the study of diabetes heterogeneity.

Heterogeneity within Autoimmune Diabetes

The discovery of insulin in 1922 readily divided diabetes into insulin-dependent cases, who required it for their very survival, and those who did not. Since children often presented with ketoacidosis, which is life-threatening without insulin treatment, their disease was seen as a juvenile-onset insulin-dependent form of diabetes, later called type 1 diabetes (T1D). Appreciation of the heterogeneity within diabetes also comes with uncertainty about the disease nosology, especially since current classification relies on clinical features and biomarkers which are not pathognomonic of underlying disease aetiology (24). Furthermore, epidemiological studies suggest that the majority of clinically defined cases of T1D are diagnosed in adulthood (22), whether in China (65% of total T1D cases) or USA (59%) (24) (25); indeed, in the UK Biobank a significant proportion of those diagnosed with diabetes over the age of 30 years had an elevated GRS for T1D (26).

The Palette Model proposes that variable elements can cluster in any given individual at a given time. Genetics and environment combine and lead to diabetes in any given individual by engaging diverse pathways at a given time. For example, genetic determinants of islet autoimmunity (e.g., HLA and non-HLA alleles associated with T1D) and processes typically associated with T2D, such as impaired beta-cell function or insulin resistance, may interact with microorganisms (viruses and the gut microbiome), dietary factors (vitamin D and gluten exposure), and factors associated with industrialization i.e., heavier BMI, hygiene, pollution and overcrowding. Pathophysiological processes that are highly prevalent in the population, such as today are obesity-related factors, can appear in combination other mechanisms, including islet autoimmunity.

The Threshold Model advances the concept that the combined effect of various diabetogenic influences must reach a critical threshold to induce diabetes. Genetic susceptibility alone may never result in clinical diabetes, and even identical twins who share a common environment may remain discordant for islet autoimmunity and T1D (27). Moreover, islet autoimmunity, when mild, may never lead to diabetes, as demonstrated by individuals who have persistent single autoantibody positivity without progression to diabetes. Even, those relatives of T1D cases carrying the protective genotype *HLA DQA1*0102/DQB1*0602*, whilst not developing diabetes, can have abnormal glucose tolerance and reduced insulin secretory capacity (28). However, such individuals can progress to diabetes as other factors accumulate, such as insulin resistance due to obesity (12) (29, 30) or concurrent T2D-associated genes (11, 13). In the Threshold Model, the number and combination of factors (e.g., genetics, environment) may vary. At one extreme, a single gene mutation e.g. *AIRE* or *FOXP3*, can cause insulin-dependent diabetes, at the other extreme, pancreatectomy or pancreatitis can do the same (31). Similarly, aggressive islet autoimmunity can lead to diabetes without requiring additional factors. The appearance of diabetes-associated autoantibodies in early life seems to reveal two distinct endotypes, one associated with insulin (proinsulin) autoantibodies and HLA DR4 (DR4PRO) (also called T1D endotype 1 or T1DE1), and the other associated with glutamic acid decarboxylase autoantibodies (GADA) and HLA DR3 (DR3GAD) (also called T1D endotype 2 or T1DE2), (25). The former identifies a cohort with an earlier peak incidence (typically diagnosed under age 7 years), insulin autoantibody as the first islet autoantibody to appear, rapid progression to multiple autoantibodies and more severe insulin secretory loss (low serum C-peptide as a proxy for insulin), allied with more marked islet inflammation (high levels of CD20 and CD8 cells infiltrating islets). On the other hand, DR3GAD endotype is usually diagnosed over age 12 years with GADA as the first and often only autoantibody to appear, less severe islet inflammation, as well as insulin secretory loss (32). T1DE1 is almost exclusive to very early childhood and T1DE2 to older individuals. At the individual level these endotypes are distinct and do not co-exist, yet at the group level they are not clearly delineated, with intermediate ages at diagnosis reflecting a continuum extending to the prevalence of multiple autoantibodies and the severity of both islet inflammation and C-peptide loss (33) (34).

The Gradient Model captures the notion that combinations of genetic and environmental factors can determine the velocity and severity of the disease process. High genetic risk with multiple serum autoantibodies at a young age are each associated with earlier progression to T1D and lower fasting C-peptide at diagnosis. Conversely, individuals can undergo remission when beta-cell function and/or insulin sensitivity improve, as with the 'honeymoon' period and following pregnancy. Both Threshold and Gradient Models account for that variable severity of insulin secretory loss once

diabetes develops, while the gradient effect extends to disease risk before clinical onset with greater insulin secretory loss in the DR4PRO endotype compared with the later onset DR3GAD endotype (Figure 2). T1D genetic risk, including the association with HLA Class 1 haplotypes, is also less striking in older cases, (35). While histologically, heterogeneity in disease severity (13) can even be detected within islets, given that not all insulin-containing cells are destroyed, while whole lobules of the pancreas can remain unaffected, so-called pancreatic 'vitiligo' (32, 36). Although a full description is beyond the scope of this article, race, ethnicity and ancestry are additional important modifiers of disease development and course. For example, elevated BMI increases progression to T1D in non-Hispanic White (by 36%), but in Hispanic children by 400% (37). Studies are underway to extend research beyond European and Chinese ancestry cohorts (38-40). Finally, and importantly, there could be utility in defining endotypes within T1D if they impact management e.g., studies suggest that immune therapy with teplizumab and GAD-alum might be more beneficial in DR4PRO and DR3GAD cases respectively, though these proposals need validation (41) (42). In T1D cases aged 7 years or more, a non-immune based agent, verapamil, known to reduce endoplasmic reticular stress in the beta cell, also appears to limit progressive decline of C-peptide post-diagnosis (43, 44).

Identifying T1D in the adult population can be challenging, as T2D becomes increasingly prevalent with age and cases can masquerade as T2D when they do not require insulin therapy, at least initially. Especially since, in cases presenting with GADA, many do not require insulin at diagnosis and have variable rates of progression to insulin therapy. Some, likely have false positive GADA, that is, the limited assay specificity provides a false signal, and have T2D (45) (46). The remainder may have an attenuated T1D phenotype or reflect the pleiotropic effects of variable disease-risk loci (24, 47) (48), the former, now called slowly evolving autoimmune diabetes of adulthood, and previously also referred to as latent autoimmune diabetes of adults (LADA), can initially masquerades as T2D (Table 1)(see Case1). However, combining clinical features, autoantibodies, C-peptide and genetic risk score (GRS) can help differentiate T1D (49) (50). Increasingly, clinics measure GADA in adult-onset cases, in combination with random blood C-peptide and glucose, to determine risk of progressing to insulin dependence (Figure 5). Identifying adults with T1D should also improve management, whether by informing prognosis relating to insulin dependence, risk of other autoimmune diseases including disease risks in relatives, or avoiding potentially less suitable adjunctive therapy (51).

Even cases with established T1D show that the severity of insulin deficiency correlates with clinical risk. In a recent study of 6,000 cases with poor glycaemic control (HbA1c 8.0-8.5%, 63-68 mmol/mol), over an average period of 5.2 years, those with fasting C-peptide >0.20 nmol/L had lower insulin requirement, HbA1c, and risk for both ketoacidosis and hypoglycaemia compared with cases with undetectable C-peptide (34). By inference, even minimal endogenous insulin secretion has a buffering activity on glucose homeostasis detectable many years post-diagnosis (33, 52). Efforts to limit the progression of the disease process using novel therapies, whether before clinical diagnosis following the appearance of immunologic and metabolic abnormalities, as found with teplizumab, or at clinical diagnosis, are worthwhile, given that there is a gradient effect so that the severity of insulin deficiency correlates with prognosis.

Heterogeneity within Type 2 diabetes (T2D)

Diabetes aetiology is highly heterogenous reflecting the many routes whereby defects can result in raised blood glucose. In the context of this underlying heterogeneity, the fact that T2D is a diagnosis of exclusion, where all diabetes that cannot be otherwise classified is grouped, makes the T2D heterogenous and poorly defined. In people labelled as T2D, there are varying contributions from factors that reduce beta-cell function and/or insulin sensitivity, such as genetics, aging, obesity, poor diet, physical inactivity, inadequate sleep, circadian disruption, stress and illness. Such non-genetic effects are striking when considering differences according to age at onset. Ideally, clinical recognition of this heterogeneity should improve our treatment and ability to predict progression and complications. Current management and guidelines of T2D (53) do little to incorporate disease heterogeneity into their recommendations.

Several recent studies have utilised a Palette Model to quantify T2D heterogeneity applying machine learning methods to two key types of data: 1) simple clinical measures or deeper phenotyping at diagnosis, or 2) grouping genetic risk variants based upon their aetiological processes to generate partitioned polygenic scores (pPS) and explore how T2D phenotypes differ in people with the genetic burden.

Utilising a “hard clustering” approach based on defined traits (age at diagnosis, presence of diabetes-associated autoantibodies, BMI, HbA1c, beta-cell function and indices of insulin resistance), diabetes heterogeneity was distilled using selected sets into five subtypes (54). These subtypes include a cluster of severe autoimmune diabetes (SAID), as well as four other forms, encompassing: severe insulin deficient diabetes (SIDD), severe insulin resistant diabetes (SIRD), mild obesity-related diabetes and mild age-related diabetes (54). In line with the Gradient Model, subtypes were associated with different disease trajectories e.g. severe insulin deficient diabetes had more rapid progression to insulin requirement, and severe insulin resistant diabetes had a worse renal prognosis (54) as well as being more likely to develop hepatic fibrosis (55). Subsequent studies showed that the T2D risk variants differed between subtypes, for example, the severe insulin resistant subtype was not associated with the classic T2D risk variant at *TCF7L2*, in contrast to the other non-autoimmune subtypes (56). In addition, subjects within the SIRD cluster were more likely carriers of the rs738409(G) variant in *PNPLA3*, which has been linked with NAFLD (57).

Although clinical features associated with these five adult diabetes subtypes tend to be consistent across different ethnicities (58) (59), “pigeonholing” individuals has inherent limitations, especially at the boundaries of hard clusters where the data points have similar distance from two different centroid. In contrast, utilizing “soft clustering”, or “fuzzy clustering”, allows individuals to be defined by more than one characteristic, given overlapping defects can contribute to diabetes in any one individual as proposed by the Palette Model (60). Such “fuzzy” or “soft” clustering has two main advantages over “hard” clustering methods. Firstly, memberships can be combined with other information. In particular, when membership of a cluster reflects probabilities, results can be combined from different sources using Bayes’ theorem. Secondly, that membership for any given subject enables membership of a ‘second best’ cluster that is almost as good as the ‘best’ cluster, a phenomenon often not apparent when using other clustering techniques (61).

One such soft clustering approach, “archetypes clustering”, applied to 726 deeply phenotyped adults with T2D from the IMI-DIRECT study (62), was consistent with the Palette Model: many people in the

middle of the palette show diabetes onset driven by many small defects, so they cannot be reliably allocated to a discrete subgroup, whereas in line with a Threshold Model, those at extremes of the palette can be allocated to discrete subgroups (Figure 3). Using this approach, while 65% of adults could not be reliably allocated to a discrete subgroup, the remainder could be allocated, with a >60% probability, to one of four clusters characterised by distinct pathophysiological mechanisms, and rates of glycaemic deterioration over 3-years. Finally, an alternative approach did not attempt to allocate people to discrete clusters, instead recognising the phenotypic distribution as continuous within adults with T2D (63); a non-linear dimensionality reduction approach using reverse graph embedding (DDRtree) was applied to nine routine clinical measures in 23,000 adults at diagnosis. The resulting tree structures visually represent phenotypic heterogeneity in this population, but also shows how that heterogeneity maps to risk of macro- and micro-vascular disease, progression to insulin requirement, and differences in drug response (Figure 4).

Another approach to clustering by phenotype employs genetic aetiology. By partitioning polygenic scores of different pathophysiological defects (e.g. reduced beta-cell function, obesity, lipodystrophy, altered liver metabolism), or other traits that capture these defects, five partitioned polygenic scores (pPS) were generated in adults (64) (65) using a fuzzy clustering approach. Every adult with T2D was allocated a score for each of these pPS; consistent with the Palette Model, most were in the middle of the 'palette'. Nevertheless, some adults with T2D had a high genetic burden for a particular pPS and consistent phenotypes (e.g., lipodystrophy, beta-cell deficiency, obesity with insulin resistance, differences in cardiovascular risk and renal disease) (65). Recent analysis showed that subjects with diabetes with different pPS have different rates of progression to diabetes-related outcomes. For example, those with higher obesity pPS have higher risks of developing end-stage renal disease and coronary heart disease, whereas those with elevated beta-cell dysfunction pPS have higher risk of proliferative diabetic retinopathy (66).

It is now clear that environmental, phenotypic and genetic heterogeneity are associated with heterogeneity in diabetes outcomes and drug response, an effect that underpins the Threshold Model of diabetes. That heterogeneity also results in variation in the severity of disease as in the Gradient Model, for example, in relation to progression of diabetes towards need for insulin (67) (68) (69). Although this concept of diabetes heterogeneity has been appreciated for some time, it has yet to be translated systematically into clinical diabetes care. However, a number of recent studies have progressed the field of precision medicine to the point of clinical implementation using simple clinical phenotypes. By recognising differences in sex, BMI and renal function, several studies could predict differences in drug responses to common treatments, i.e. sulphonylureas, DPP4 inhibitors, thiazolidinediones and SGLT2 inhibitors (70, 71). Importantly, observational adult primary care data and post-hoc analysis of randomised controlled trials found that reduced renal function can be used to preferentially select DPP4 inhibitor over an SGLT2 inhibitor and, from the Trimaster 3-way cross-over study, to select pioglitazone over DPP4 inhibitors when individuals have a BMI>30 kg/m² (72). These studies pave the way for incorporation of phenotypic heterogeneity into routine clinical care, guiding treatment choice. More work is required before genetic information is used for clinical management in T2D – although there are robust genetic determinants of drug response (reviewed in (73)), likely implementable when genetic data are routinely embedded into medical record systems.

Heterogeneity in non-autoimmune diabetes in youth

Historically, T2D was thought to occur almost exclusively in adults, as a consequence of factors noted above, most notably those linked with ageing. However, that is changing and while the overall incidence of youth-onset T1D still outnumbers youth-onset T2D, there has been a marked increase in the latter in multiple populations from Canada, the United States, through the Middle-East, to China (74, 75),(76, 77) (78). As with adult-onset T2D, youth-onset diabetes is also heterogeneous (see Case 2). The Treatment Options for Diabetes in Adolescents and Youth (TODAY) study in the United States (79) showed that most youths with T2D have obesity and are pubertal, with a very sedentary lifestyle, strong family history of T2D and/or *in utero* exposure to T2D or gestational diabetes, while in the United States they tend to be of underrepresented race/ethnicity (80). However, in line with a Threshold Model, only the minority of youth with these risk factors develop T2D, therefore other factors, both genetic and non-genetic, are likely needed to reach the threshold to develop diabetes.

Using the Palette Model, genetic factors (e.g., HLA and non-HLA alleles), including those typically associated with T2D, such as lower beta-cell function, insulin resistance and environmental factors noted above, but extending to adverse childhood experiences, can lead to diabetes in any given youth by engaging diverse pathways at a given time. Employing the Threshold Model, that interaction between genes and environment likely must reach a critical threshold to induce diabetes. Genetic abnormalities, such as those that determine insulin resistance, lower beta-cell function, or T2D-associated genes (11) may never result in clinical diabetes unless additional factors e.g., obesity or puberty exacerbate the condition. As with the Gradient Model, any combination of these factors once diabetes develops can determine disease severity. Conversely, individuals can undergo remission when insulin sensitivity improves, as when exiting puberty, or following bariatric surgery. The association between obesity and T2D is stronger in youth with younger age at presentation, while prepubertal T2D, albeit rare, appears almost exclusively in children with elevated BMI (81).

As with adult-onset T2D, phenotype clustering occurs in youth. The SEARCH study classified 2,291 recently diagnosed youth aged <20-years using diabetes autoantibodies (DAA) and insulin sensitivity (IS), the latter estimated using HbA1c, waist circumference and triglycerides, validated against hyperinsulinemic-euglycaemic clamps (82). Four categories were described based on presence or not of autoimmunity and insulin resistance. The largest categories were DAA+/IS+ve (54.5%) or DAA-/IS-ve (15.9%), which aligned with provider clinical diagnosis in 90% of cases. The DAA-/IS-ve group also had a greater urinary albumin: creatinine ratio than other groups (83). Since DAA+/IS-ve group (19.5%) had similar prevalence and levels of DAA and similar distribution of HLA risk genotypes as the DAA+/IS+ve group, thus may represent T1D in youth with obesity. The remainder (10.1%) without DAA but insulin sensitive may represent a combination of undetected autoimmunity and monogenic diabetes requiring further investigation. Using these same four categories SEARCH also found that the T1D GRS was highest in DAA+/IS+ve and lowest in the DAA-/IS-ve groups, while the converse held true for the T2D GRS. The genetic probability of T1D identified those without DAA most likely to progress to insulin requirement (39).

Just as defining T1D in adulthood is difficult, it is also difficult to determine T2D in youth given its increasing incidence, coupled with rising rates of obesity. The multi-centre SEARCH for Diabetes in Youth study in the US showed that *HNF1A*, *HNF4A*, and *GCK* mutations causing MODY accounted for 1.2% of diabetes cases and 7–15% of all youth cases without diabetes-associated autoantibodies (84). MODY should be considered in autoantibody-negative diabetes cases, even when obesity is present. Supporting that conclusion, the TODAY study, found that 4.5% of youth clinically diagnosed

with T2D and obesity had genetic variants consistent with MODY (85). Both insulin secretory loss and insulin sensitivity vary greatly in youth onset T2D. Recent data from both adolescents and adults with T1D and T2D found that whole-body insulin sensitivity in normal-weight youth with T1D is roughly equivalent to insulin sensitivity in nondiabetic youth with obesity, while youth with obesity plus T1D have reduced insulin sensitivity similar to youth with T2D (86). Further, adjunctive treatment with metformin (87) (88) and bromocriptine quick-release (89) have been shown in youth with T1D to lower insulin dose and improve insulin resistance, BMI, blood pressure, carotid intimal media thickness (90), and aortic stiffness (with metformin), and blood pressure and aortic stiffness (with bromocriptine). Interestingly, the metabolic syndrome phenotype characteristic of insulin resistance in youth T2D differs from that in T1D; low HDL cholesterol and adiponectin concentrations and elevated triglycerides and ectopic fat (liver, visceral and muscle fat) typical of the former are less common in the latter. Therefore, a metabolic syndrome phenotype can support the diagnosis of T2D. As noted above, in youths as in adults, those with a T2D phenotype, but single GADA, present a diagnostic challenge, and the phenotype of youth should be assessed to personalize treatment and address co-morbidities (Fig 4) (91).

Some of the conclusions about the differences between adult- and youth-onset diabetes are likely to be confounded by ethnicity, socio-demography, and temporal factors, as these tend to differ by age-group. For example, the phenotype of youth-onset T2D differs from adults, including a strong female predominance (79) (92), although some features are confounded by other factors including ethnicity, e.g., female preponderance is not apparent in Asia (93). The clinical course of youth-onset T2D is more aggressive than adult-onset T2D, with more rapid insulin secretory loss, poorer response to oral medications, and earlier onset of diabetes-related complications (see Case 4). Glycaemic failure rates and decline in insulin secretory function were more than twice higher in the TODAY study (comparing metformin alone or in combination with rosiglitazone or intensive lifestyle therapy) than in similar adult trials, despite youth having a shorter diabetes duration (94), e.g. therapy failure rate with metformin alone was 51.7% in TODAY vs. 12% in the adult ADOPT study (95). Similarly, the Restoring Insulin Secretion (RISE) Consortium, comparing 3 months of insulin glargine then 9 months metformin with 12 months of metformin, it was found that youth with T2D had lower insulin sensitivity and higher insulin secretion than adults, yet insulin secretion declined more rapidly in youth than adults in whom it was stable. Multiple studies also showed the much earlier onset of diabetes-associated complications despite shorter diabetes duration (96) (97) (98) (99) (100). These observations argue for careful categorising of youth with diabetes and aggressive treatment of their glycemia, co-morbidities and diabetes complications.

Looking to the future

To understand disease heterogeneity, one should capture the nature of that variability and define how, in any given individual, it is related to the disease pathogenesis, risk and outcome. Here we integrated three models: starting with multiple risk factors on a palette, their combination may attain a given threshold to affect a given individual to cause diabetes, while the impact of those risk factors once diabetes has developed, as well as leading up to the diagnosis, will create a gradient of disease severity, both for T1D and T2D.

We envisage integrating genetic, demographic and phenotypic data to define disease risk and disease-outcome in a given individual (51, 101) (102), recognising that it may not always be possible to assign individuals to a distinct diagnostic category. Recognition of the different pathophysiological pathways contributing to hyperglycaemia, including beta-cell dysfunction, adipose tissue dysfunction, altered incretin response etc. has highlighted the opportunity to better tailor treatment

according to underlying pathophysiology (103). Interestingly, some of the pathways leading to diabetes converge at the insulin-secreting cell endoplasmic reticulum whose stress response is regulated by gene variants common to both major types of diabetes (8); by addressing that stress response it should be possible to protect the insulin-secreting cell, indicating that early therapy could better prevent diabetes, irrespective of its cause (43) (44). Proteomic or metabolite studies should aid identification of at-risk cases e.g., using machine learning to proteomic profiles from a single fasted sample found that only three proteins (RTN4R, CBPM and GHR) could identify individuals at risk of T2D (104).

Since lifestyle and metformin can both limit progression from impaired glucose tolerance to T2D, identifying that risk at an even earlier stage should be valuable. Not least when we can exploit an arsenal of new drugs to reduce risk. We already define risk for microvascular and macrovascular disease with BMI, hypertension and dyslipidaemia. Furthermore, in line with BMI impacting progression through preclinical stages (14, 29, 30), it is already incorporated in a predictive model for T1D (102). When GRS is eventually used clinically, its utility could be expanded to diagnose diabetes type, guide treatment selections and predict treatment success, as well as define risk of vascular complications and other conditions such as coeliac disease. Such biomarkers and risk scores need to be developed into easy-to-use tools to communicate risk to clinicians and patients to guide clinical decision. Screening programmes must be made available, both locally and globally, and tailored to different ethnic groups. For example, despite GRS for T1D being different between China and Europe (105, 106), it should be possible to define a “cosmopolitan” GRS that can be used in ethnically -diverse populations, which is problematic in admixed populations, in addition to ethnicity-specific GRS (reviewed in (38)). Likewise, a multi-ethnic GRS for T2D has also been generated (66).

Defining heterogeneity of disease risk or complications risk within types of diabetes should improve the outcome of interventions, and enable us to identify factors which account for that heterogeneity. For example, endotypes identified in T1D may define the optimum immune therapy and that approach offers a signpost towards targeted therapy to be exploited in the future in all forms of diabetes. There is an ongoing international effort, led by the American Diabetes Association and European Association for the Study of Diabetes, to review the current evidence on diabetes diagnosis, classification and prognostication, with the ultimate aim to improve precision medicine in diabetes (5).

Finally, there are two important elements to population screening. First, it should not be limited to GRS, a relatively crude tool when taken in isolation, but should be integrated into an individualised package which considers demographics, family history and a broad range of clinical features as well as laboratory-based features, including proteomics (24, 107). Second, as with all screening programmes, it will be essential that the benefits and the limitations of screening are fully explained and that a robust system is in place to deal with uncertainties and with queries as they arise. As more user-friendly tools such as risk calculators or pathway-specific scores are developed, they should be incorporated into clinical service to improve diabetes care. It will require widespread engagement across different sectors to fully realize the potential of precision medicine to improve diabetes care.

Conflict of Interest:

RDL is on the Advisory Boards of Diamyd and Provention and has received educational grants from Novo-Nordisk, Eli Lilly, Sanofi, Astra Zeneca and MSD. RCWM has received research funding from AstraZeneca, Bayer, Novo Nordisk, Pfizer, Roche Diagnostics (HK) Limited and Tricida Inc. for carrying

out clinical trials, and has received speaker honorarium or consultancy in advisory boards from AstraZeneca, Bayer, Boehringer Ingelheim, Medtronic, Merck and Kyowa Kirin. All proceeds have been donated to the Chinese University of Hong Kong to support diabetes research. RCWM is a cofounder of GemVCare, a diabetes technology start-up, established through support from the Hong Kong Government Technology Start-up Support Scheme for Universities (TSSSU). ERP has received honoraria for speaking from Lilly, Novo and Illumina. The other authors have no conflicts to disclose.

Funding acknowledgement:

RDL is funded by the British Diabetic Twin Trust, Barts Charity and EU (FP7) 282510. RCWM acknowledges support from the Research Grants Council of the Hong Kong Special Administrative Region (CU R4012-18), a Croucher Foundation Senior Medical Research Fellowship, University Grants Committee Research Grants Matching Scheme and the Health and Medical Research Fund (CFS-CUHK2). PWF acknowledges support from the European Research Council (CoG-2015_681742_NASCENT) and the Swedish Research Council (Distinguished Young Researchers Award in Medicine). KJN acknowledges support from NIDDK grant # P30-DK116073, and NIH NHLBI HL145076-01. MJR is funded by NIH NIDDK through R01 DK124395, R01 DK121843, R01 DK053456 and U54 DK118638.

Contributions statement:

RDL, RCWM, PWF, KJN, ERP, MJR all contributed to the conceptualisation, data curation, formal analysis, visualization, writing original draft, and writing-review and editing of the revised manuscript. RDL and RCWM contributed equally and should be considered joint first authors.

Box- Illustrative cases

Case 1- A lean fit man presents at age 33 with diabetes diagnosed on the basis of both oral glucose tolerance and HbA1c. He has a family history of T1D and his monozygotic twin has T1D diagnosed at age 14 years with HLA-DR4 and GADA alone. Multiple testing for diabetes-associated autoantibodies were initially positive at an earlier age but subsequently negative and his random serum C-peptide was >600 pM. He had thyroid autoantibodies but was euthyroid. Started on metformin he was intolerant to it and put on a DPP-4 inhibitor. He remains on a DPP-4 inhibitor 15 years later with HbA1c <53 mmol/mol/ 7.0%.

Case 2- A 43 year-old Chinese man was found to have hypertension at health check, and was also found to have elevated blood glucose of 13mmol/l, with preceding weight loss of 20 lbs. BMI 21.7 kg/m². HbA1c 119 mmol/mol /13% at presentation. Family history of diabetes present. Started metformin and sitagliptin initially by primary care physician and referred for specialist evaluation. HbA1c improved and condition then stable with HbA1c 44 mmol/mol /6.2% on metformin alone. Further evaluation showed adequate fasting C-peptide of 1026pmol/l, and GADA and anti-IA2 antibodies were negative. Gene screening panel for monogenic forms of diabetes including Lamin A/C gene were all negative. GRS showed high polygenic risk for T2D.

538 Case 3- A 10 year-old girl presented with polyuria and polydipsia (HbA1c was 108 mmol/mol /12%
539 and fasting blood glucose 250 mg/dL / 14 mmol/l), marked ketonuria, and autoimmune primary
540 hypothyroidism (initial TSH 15.8, free T4 1.0 ng/dl [0.7-1.9] and positive thyroid peroxidase
541 autoantibodies). She had acanthosis nigricans, was in puberty with height at the 75th centile and
542 weight BMI both above 97th centile (i.e. obesity). She also had high triglycerides, low HDL and
543 elevated liver enzymes. Antibodies against GAD, insulin, islet cell and zinc transporter were all
544 negative. Her parents were overweight with dyslipidaemia and hypothyroidism. She was initially
545 treated with levothyroxine and insulin infusion. The latter was then replaced with long-acting insulin
546 and metformin. Subsequently, insulin was stopped and metformin continued.

547

548 Case 4- A 12 year-old Asian (from Indian subcontinent) boy was diagnosed one year earlier with
549 diabetes ketoacidosis and presumed T1D with single GADA positive (negative for IA-2, insulin and
550 ZnT8 autoantibodies). He did not have HLA DR3 or DR4. A screen for MODY genes was negative. His
551 father had T2D. The patient was started on insulin, but he was able to come off insulin 3 weeks after
552 onset, and 8 months post-diagnosis had a HbA1c of 38 mmol/mol /5.6% with random C-peptide 773
553 pM, though he remains GADA positive.

554 Learning Points from the cases:

- 555
- Diabetes with GADA is not invariably and initially insulin-requiring (case 1, case 4)
 - 556 • Asian and African cases may not fit into phenotypes as designated by European and
 - 557 American studies (case 2, case 4) e.g. without high risk HLA variants or obesity.
 - 558 • C-peptide can help decide whether insulin treatment is imperative (case 1, case 2)
 - 559 • Even ketoacidosis in children is not inevitably associated with early insulin-dependence (case
 - 560 3, case 4)

561

Figure legends.

Figure 1. Conceptual diagram illustrating three different models to illustrate heterogeneity of common diseases including the Palette Model, the Threshold Model and the Gradient Model. Upper panel highlights the general concept of the models, whilst the lower panel provides examples when the models are applied in the context of diabetes heterogeneity. The Palette Model illustrates how multiple risk-factors (genetic and non-genetic) can contribute to a given disease. The Threshold Model illustrates how, in any given individual, some but not necessarily all, risk factors will be of sufficient severity that the altered tissue function crosses a threshold leading to clinical disease. The Gradient Model illustrates how the same disease-risk factors leading to the clinical disease will impact the severity of that clinical disease post-diagnosis, which will vary between individuals. In the top figure, a circular dot represents risk-factors associated with diabetes, be they genetic, non-genetic or disease-associated pathways and networks. In the Threshold Model, each colour represents those same processes in given individuals of sufficient intensity, operating in tandem, to lead to clinical disease. Once the disease develops, it will be of variable severity as shown in the Gradient Model, and that severity will reflect the intensity of those factors leading to the disease in the first place.

In the figure depicting the palette model, each colour represents example of an underlying aetiological process, such as:

Green = lipodystrophy

Blue = obesity and insulin resistance

Red = beta-cell deficiency

Purple = disorders of incretin action

Yellow = beta-cell autoimmunity

Most people with T2D are 'grey' and represent a mixed aetiology.

In the threshold model, increasing severity from yellow to red indicate varying severity of the defect in for a particular tissue dysfunction, e.g. insulin secretory dysfunction, and hence increasing risk of diabetes once a threshold is reached for the diagnosis of diabetes

In the gradient model, individuals can be viewed as being on a continuum of risk post-diagnosis of "T1D" or "T2D", in terms of different prognosis, including progression to insulin.

Figure 2. Conceptual diagram illustrating the heterogeneity of diabetes in terms of underlying pathophysiological defects, relative contribution of genetic factors, and its impact on progression to hyperglycaemia. Figure adapted from Redondo et al, Diabetologia 2020 (9).

Figure 3. A visualisation of the heterogeneity of diabetes, based upon an extension of the Palette Model (60) and reproduced from reference (62). Each colour represents an aetiological process, and each dot, a person with diabetes, with increasing depth of colour representing a greater contribution to the diabetes aetiology. For example:

- Green = lipodystrophy
- Blue = obesity and insulin resistance
- Red = beta-cell deficiency
- Purple = disorders of incretin action
- Yellow = Beta-cell autoimmunity

Most people with T2D are 'grey' and represent a mixed aetiology; some with T2D are deeper in colour as they have diabetes due predominantly to a defect in that pathway. The aetiological drivers of T1D and monogenic diabetes are largely distinct but there are people who have overlap in aetiology for their diabetes. PC1 and PC2 represent the principal components 1 and 2, respectively.

Figure 4. Schematic diagram highlighting the phenotypic heterogeneity of diabetes. People with newly diagnosed T2D can be represented on the tree based upon nine phenotypic characteristics. The phenotypic values are overlaid on a tree structure to visualize the distribution of the nine phenotypic traits over the tree structure. Magenta colour indicates high value whereas green colour indicates low value of the phenotype. Figure reproduced from Nair et al, Nature Medicine 2022 (63).


Figure 5. Suggested algorithm for the evaluation of diabetes at presentation according to age and disease duration exploiting diabetes-associated autoantibodies, C-peptide and T1D GRS.

Key: T1D GRS: T1D genetic risk score
AABBCC (see text): Age, Autoimmunity, BMI, Background, Control (HbA1c), Complications.

625
626
627

Figure 1.

Common Chronic Diseases




628

Diabetes: Different Models

Pathophysiology:
Beta-cell autoimmunity; beta-cell
deficiency, obesity, lipodystrophy,
impaired incretin action

Insulin Secretory
Dysfunction

Type 1 diabetes
and
Type 2 diabetes



629
630
631
632
633
634
635
636

Figure 2.




Figure 3.




Figure 4.





Figure 5.



677

678 Table 1. Types of diabetes according to current 2019 WHO classification (table adapted from (6))

679

Types of diabetes	Brief description	Clinical implications	Updates from previous classification
Type 1 diabetes (T1D)	Immune-mediated beta-cell destruction and absolute insulin deficiency. Onset most commonly in childhood or early adulthood	Requires insulin	T1D subclasses now removed
Type 2 diabetes (T2D)	Most common type, various degrees of beta-cell dysfunction and insulin resistance. Commonly, but not invariably, associated with overweight, obesity, inactivity and metabolic syndrome.	Individuals assumed to have “T2D” when not fitting apparent T1D or other forms. Variable demographics and disease course. Different treatments now for different pathophysiologies	T2D subclasses now removed
Hybrid forms of diabetes			
Slowly evolving, immune-mediated diabetes of adults	Similar to slowly evolving T1D in adults, with a single GAD autoantibody, initial preservation of beta-cell function, often may have features of metabolic syndrome	Variable course but ultimately often require exogenous insulin.	Change in nomenclature- previously referred to as Latent Autoimmune Diabetes of Adults (LADA)
Ketosis-prone diabetes	Presents with ketosis and insulin deficiency, which resolves, at least initially. Variable pathophysiology includes T1D and T2D.	May require insulin initially, but with variable disease cause and course	
Other specific types			

<p>Monogenic diabetes</p> <ul style="list-style-type: none"> - Monogenic defects of beta-cell function - Monogenic defects in insulin action 	<p>Caused by specific genetic mutations, with presentation ranging between the neonatal period and early adulthood.</p> <p>Caused by specific mutations leading to severe insulin resistance in absence of obesity</p>	<p>Different pancreatic and extra-pancreatic clinical manifestations with specific treatment implications.</p>	<p>Updated nomenclature to highlight specific genetic defects</p>
<p>Type 3 diabetes (Diseases of exocrine pancreas)</p>	<p>Caused by various injury to the pancreas</p>	<p>Exocrine pancreas now implicated in T1D and T2D diabetes</p>	
<p>Endocrine disorders</p>	<p>Occurs in diseases with excessive production of hormones that antagonize insulin action</p>	<p>Treatment of underlying condition may improve glycaemia</p>	
<p>Drug or chemical-induced</p>	<p>Secondary to chemicals that affect insulin secretion or action</p>	<p>May be reversible; consider switching responsible medication</p>	
<p>Infection-related diabetes</p>	<p>May be due to direct virus-related beta-cell destruction; systemic inflammation; stress-related insulin resistance; etc</p>	<p>May or may not be reversible</p>	
<p>Uncommon specific forms of immune-mediated diabetes</p>	<p>Associated with rare immune-mediated diseases</p>		
<p>Other genetic syndromes sometimes associated with diabetes</p>	<p>Rare genetic or chromosomal disorders linked to diabetes through various mechanisms</p>	<p>Variable implications on treatment, with some requiring early insulin treatment</p>	
<p>Unclassified diabetes</p>	<p>For cases that do not fit</p>		

	into any of the categories		
Hyperglycaemia first detected during pregnancy			
Diabetes mellitus in pregnancy	T1D or T2D first diagnosed during pregnancy	May be confused with gestational diabetes mellitus (GDM). Associated with increased risk of pregnancy complications.	
Gestational diabetes mellitus (GDM)	Hyperglycaemia first presenting only in pregnancy, with high-risk of later progression to T2D	Affects significant proportion of pregnancies using current thresholds of diagnosis.	Defined by the WHO 2013 criteria for GDM

680

681

References

1. Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB, Stein C, Basit A, Chan JCN, Mbanya JC, Pavkov ME, Ramachandaran A, Wild SH, James S, Herman WH, Zhang P, Bommer C, Kuo S, Boyko EJ, Magliano DJ. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes research and clinical practice*. 2022;183:109119.
2. Magliano DJ, Boyko EJ. IDF DIABETES ATLAS. IDF Diabetes Atlas. 10th ed. Brussels2021.
3. Varghese JS, Narayan KMV. Ethnic differences between Asians and non-Asians in clustering-based phenotype classification of adult-onset diabetes mellitus: A systematic narrative review. *Prim Care Diabetes*. 2022;16(6):853-6.
4. Ma RC, Chan JC. Type 2 diabetes in East Asians: similarities and differences with populations in Europe and the United States. *Ann N Y Acad Sci*. 2013;1281:64-91.
5. Chung WK, Erion K, Florez JC, Hattersley AT, Hivert MF, Lee CG, McCarthy MI, Nolan JJ, Norris JM, Pearson ER, Philipson L, McElvaine AT, Cefalu WT, Rich SS, Franks PW. Precision Medicine in Diabetes: A Consensus Report From the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes care*. 2020;43(7):1617-35.
6. World Health Organization. Classification of diabetes mellitus. 2019 21 April 2019. Report No.: ISBN 978-92-4-151570-2.
7. Deutsch AJ, Ahlqvist E, Udler MS. Phenotypic and genetic classification of diabetes. *Diabetologia*. 2022;65(11):1758-69.
8. Charles MA, Leslie RD. Diabetes: Concepts of beta-Cell Organ Dysfunction and Failure Would Lead to Earlier Diagnoses and Prevention. *Diabetes*. 2021;70(11):2444-56.
9. Redondo MJ, Hagopian WA, Oram R, Steck AK, Vehik K, Weedon M, Balasubramanyam A, Dabelea D. The clinical consequences of heterogeneity within and between different diabetes types. *Diabetologia*. 2020;63(10):2040-8.
10. Redondo MJ, Evans-Molina C, Steck AK, Atkinson MA, Sosenko J. The Influence of Type 2 Diabetes-Associated Factors on Type 1 Diabetes. *Diabetes care*. 2019;42(8):1357-64.
11. Redondo MJ, Geyer S, Steck AK, Sosenko J, Anderson M, Antinozzi P, Michels A, Wentworth J, Xu P, Pugliese A, Type 1 Diabetes TrialNet Study G. TCF7L2 Genetic Variants Contribute to Phenotypic Heterogeneity of Type 1 Diabetes. *Diabetes care*. 2018;41(2):311-7.
12. Redondo MJ, Sosenko J, Libman I, McVean JF, Tosur M, Atkinson MA, Becker D, Geyer S. Single Islet Autoantibody at Diagnosis of Clinical Type 1 Diabetes is Associated With Older Age and Insulin Resistance. *The Journal of clinical endocrinology and metabolism*. 2020;105(5):1629-40.
13. Redondo MJ, Richardson SJ, Perry D, Minard CG, Carr ALJ, Brusko T, Kusmartseva I, Pugliese A, Atkinson MA. Milder loss of insulin-containing islets in individuals with type 1 diabetes and type 2 diabetes-associated TCF7L2 genetic variants. *Diabetologia*. 2023;66(1):127-31.
14. Ferrara-Cook C, Geyer SM, Evans-Molina C, Libman IM, Becker DJ, Gitelman SE, Redondo MJ, Type 1 Diabetes TrialNet Study G. Excess BMI Accelerates Islet Autoimmunity in Older Children and Adolescents. *Diabetes care*. 2020;43(3):580-7.
15. Redondo MJ, Steck AK, Sosenko J, Anderson M, Antinozzi P, Michels A, Wentworth JM, Atkinson MA, Pugliese A, Geyer S, Type 1 Diabetes TrialNet Study G. Transcription Factor 7-Like 2 (TCF7L2) Gene Polymorphism and Progression From Single to Multiple Autoantibody Positivity in Individuals at Risk for Type 1 Diabetes. *Diabetes care*. 2018;41(12):2480-6.
16. Dabelea D, Mayer-Davis EJ, Lamichhane AP, D'Agostino RB, Jr., Liese AD, Vehik KS, Narayan KM, Zeitler P, Hamman RF. Association of intrauterine exposure to maternal diabetes and obesity with type 2 diabetes in youth: the SEARCH Case-Control Study. *Diabetes care*. 2008;31(7):1422-6.
17. Lowe WL, Jr., Scholtens DM, Lowe LP, Kuang A, Nodzenski M, Talbot O, Catalano PM, Linder B, Brickman WJ, Clayton P, Deerochanawong C, Hamilton J, Josefson JL, Lashley M, Lawrence JM, Lebenthal Y, Ma R, Maresh M, McCance D, Tam WH, Sacks DA, Dyer AR, Metzger BE. Association of

732 Gestational Diabetes With Maternal Disorders of Glucose Metabolism and Childhood Adiposity.
733 *Jama*. 2018;320(10):1005-16.

734 18. Armengaud JB, Ma RCW, Siddeek B, Visser GHA, Simeoni U. Offspring of mothers with
735 hyperglycaemia in pregnancy: The short term and long-term impact. What is new? *Diabetes research*
736 *and clinical practice*. 2018;145:155-66.

737 19. Kahn SE. The relative contributions of insulin resistance and beta-cell dysfunction to the
738 pathophysiology of Type 2 diabetes. *Diabetologia*. 2003;46(1):3-19.

739 20. DeFronzo RA, Banerji MA, Bray GA, Buchanan TA, Clement S, Henry RR, Kitabchi AE,
740 Mudaliar S, Musi N, Ratner R, Reaven P, Schwenke DC, Stentz FD, Tripathy D, Group ANS.
741 Determinants of glucose tolerance in impaired glucose tolerance at baseline in the Actos Now for
742 Prevention of Diabetes (ACT NOW) study. *Diabetologia*. 2010;53(3):435-45.

743 21. Roden M, Shulman GI. The integrative biology of type 2 diabetes. *Nature*.
744 2019;576(7785):51-60.

745 22. Taylor R, Al-Mrabeh A, Zhyzhneuskaya S, Peters C, Barnes AC, Aribisala BS, Hollingsworth KG,
746 Mathers JC, Sattar N, Lean MEJ. Remission of Human Type 2 Diabetes Requires Decrease in Liver and
747 Pancreas Fat Content but Is Dependent upon Capacity for beta Cell Recovery. *Cell metabolism*.
748 2018;28(4):547-56 e3.

749 23. Self S, McMahan C, Mokalled S. Capturing the pool dilution effect in group testing
750 regression: A Bayesian approach. *Stat Med*. 2022;41(23):4682-96.

751 24. Leslie RD, Evans-Molina C, Freund-Brown J, Buzzetti R, Dabelea D, Gillespie KM, Goland R,
752 Jones AG, Kacher M, Phillips LS, Rolandsson O, Wardian JL, Dunne JL. Adult-Onset Type 1 Diabetes:
753 Current Understanding and Challenges. *Diabetes care*. 2021;44(11):2449-56.

754 25. Weng J, Zhou Z, Guo L, Zhu D, Ji L, Luo X, Mu Y, Jia W, Group TDCS. Incidence of type 1
755 diabetes in China, 2010-13: population based study. *BMJ (Clinical research ed)*. 2018;360:j5295.

756 26. Thomas NJ, Jones SE, Weedon MN, Shields BM, Oram RA, Hattersley AT. Frequency and
757 phenotype of type 1 diabetes in the first six decades of life: a cross-sectional, genetically stratified
758 survival analysis from UK Biobank. *Lancet Diabetes Endocrinol*. 2018;6(2):122-9.

759 27. Redondo MJ, Yu L, Hawa M, Mackenzie T, Pyke DA, Eisenbarth GS, Leslie RD. Heterogeneity
760 of type I diabetes: analysis of monozygotic twins in Great Britain and the United States. *Diabetologia*.
761 2001;44(3):354-62.

762 28. Greenbaum CJ, Eisenbarth G, Atkinson M, Yu L, Babu S, Schatz D, Zeidler A, Orban T,
763 Wasserfall C, Cuthbertson D, Krischer J, group DPTs. High frequency of abnormal glucose tolerance
764 in DQA1*0102/DQB1*0602 relatives identified as part of the Diabetes Prevention Trial--Type 1
765 Diabetes. *Diabetologia*. 2005;48(1):68-74.

766 29. Ferrara CT, Geyer SM, Evans-Molina C, Libman IM, Becker DJ, Wentworth JM, Moran A,
767 Gitelman SE, Redondo MJ, Type 1 Diabetes TrialNet Study G. The Role of Age and Excess Body Mass
768 Index in Progression to Type 1 Diabetes in At-Risk Adults. *The Journal of clinical endocrinology and*
769 *metabolism*. 2017;102(12):4596-603.

770 30. Ferrara CT, Geyer SM, Liu YF, Evans-Molina C, Libman IM, Besser R, Becker DJ, Rodriguez H,
771 Moran A, Gitelman SE, Redondo MJ, Type 1 Diabetes TrialNet Study G. Excess BMI in Childhood: A
772 Modifiable Risk Factor for Type 1 Diabetes Development? *Diabetes care*. 2017;40(5):698-701.

773 31. Wasserfall C, Nead K, Mathews C, Atkinson MA. The threshold hypothesis: solving the
774 equation of nurture vs nature in type 1 diabetes. *Diabetologia*. 2011;54(9):2232-6.

775 32. Leete P, Willcox A, Krogvold L, Dahl-Jorgensen K, Foulis AK, Richardson SJ, Morgan NG.
776 Differential Insulinitic Profiles Determine the Extent of beta-Cell Destruction and the Age at Onset of
777 Type 1 Diabetes. *Diabetes*. 2016;65(5):1362-9.

778 33. Jeyam A, Colhoun H, McGurnaghan S, Blackbourn L, McDonald TJ, Palmer CNA, McKnight JA,
779 Strachan MWJ, Patrick AW, Chalmers J, Lindsay RS, Petrie JR, Thekkepat S, Collier A, MacRury S,
780 McKeigue PM. Clinical Impact of Residual C-Peptide Secretion in Type 1 Diabetes on Glycemia and
781 Microvascular Complications. *Diabetes care*. 2021;44(2):390-8.

34. Rickels MR, Evans-Molina C, Bahnson HT, Ylescupidez A, Nadeau KJ, Hao W, Clements MA, Sherr JL, Pratley RE, Hannon TS, Shah VN, Miller KM, Greenbaum CJ. High residual C-peptide likely contributes to glycemic control in type 1 diabetes. *The Journal of clinical investigation*. 2020;130(4):1850-62.
35. Mishra R, Akerlund M, Cousminer DL, Ahlqvist E, Bradfield JP, Chesi A, Hodge KM, Guy VC, Brillon DJ, Pratley RE, Rickels MR, Vella A, Ovalle F, Harris RI, Melander O, Varvel S, Hakonarson H, Froguel P, Lonsdale JT, Mauricio D, Schloot NC, Khunti K, Greenbaum CJ, Yderstraede KB, Tuomi T, Voight BF, Schwartz S, Boehm BO, Groop L, Leslie RD, Grant SFA. Genetic Discrimination Between LADA and Childhood-Onset Type 1 Diabetes Within the MHC. *Diabetes care*. 2020;43(2):418-25.
36. Battaglia M, Ahmed S, Anderson MS, Atkinson MA, Becker D, Bingley PJ, Bosi E, Brusko TM, DiMeglio LA, Evans-Molina C, Gitelman SE, Greenbaum CJ, Gottlieb PA, Herold KC, Hessner MJ, Knip M, Jacobsen L, Krischer JP, Long SA, Lundgren M, McKinney EF, Morgan NG, Oram RA, Pastinen T, Peters MC, Petrelli A, Qian X, Redondo MJ, Roep BO, Schatz D, Skibinski D, Peakman M. Introducing the Endotype Concept to Address the Challenge of Disease Heterogeneity in Type 1 Diabetes. *Diabetes care*. 2020;43(1):5-12.
37. Tosur M, Geyer SM, Rodriguez H, Libman I, Baidal DA, Redondo MJ, Type 1 Diabetes TrialNet Study G. Ethnic differences in progression of islet autoimmunity and type 1 diabetes in relatives at risk. *Diabetologia*. 2018;61(9):2043-53.
38. Redondo MJ, Gignoux CR, Dabelea D, Hagopian WA, Onengut-Gumuscu S, Oram RA, Rich SS. Type 1 diabetes in diverse ancestries and the use of genetic risk scores. *Lancet Diabetes Endocrinol*. 2022;10(8):597-608.
39. Oram RA, Sharp SA, Pihoker C, Ferrat L, Imperatore G, Williams A, Redondo MJ, Wagenknecht L, Dolan LM, Lawrence JM, Weedon MN, D'Agostino R, Hagopian WA, Divers J, Dabelea D. Utility of Diabetes Type-Specific Genetic Risk Scores for the Classification of Diabetes Type Among Multiethnic Youth. *Diabetes care*. 2022;45(5):1124-31.
40. Balcha SA, Demisse AG, Mishra R, Vartak T, Cousminer DL, Hodge KM, Voight BF, Lorenz K, Schwartz S, Jerram ST, Gamper A, Holmes A, Wilson HF, Williams AJK, Grant SFA, Leslie RD, Phillips DIW, Trimble ER. Type 1 diabetes in Africa: an immunogenetic study in the Amhara of North-West Ethiopia. *Diabetologia*. 2020;63(10):2158-68.
41. Herold KC, Bundy BN, Long SA, Bluestone JA, DiMeglio LA, Dufort MJ, Gitelman SE, Gottlieb PA, Krischer JP, Linsley PS, Marks JB, Moore W, Moran A, Rodriguez H, Russell WE, Schatz D, Skyler JS, Tsalikian E, Wherrett DK, Ziegler AG, Greenbaum CJ, Type 1 Diabetes TrialNet Study G. An Anti-CD3 Antibody, Teplizumab, in Relatives at Risk for Type 1 Diabetes. *The New England journal of medicine*. 2019;381(7):603-13.
42. Nowak C, Lind M, Sumnik Z, Pelikanova T, Nattero-Chavez L, Lundberg E, Rica I, Martinez-Brocca MA, Ruiz de Adana M, Wahlberg J, Hanas R, Hernandez C, Clemente-Leon M, Gomez-Gila A, Ferrer Lozano M, Sas T, Pruhova S, Dietrich F, Puente-Marin S, Hannelius U, Casas R, Ludvigsson J. Intralymphatic GAD-Alum (Diamyd(R)) Improves Glycemic Control in Type 1 Diabetes With HLA DR3-DQ2. *The Journal of clinical endocrinology and metabolism*. 2022;107(9):2644-51.
43. Ovalle F, Grimes T, Xu G, Patel AJ, Grayson TB, Thielen LA, Li P, Shalev A. Verapamil and beta cell function in adults with recent-onset type 1 diabetes. *Nat Med*. 2018;24(8):1108-12.
44. Forlenza GP, McVean J, Beck RW, Bauza C, Bailey R, Buckingham B, DiMeglio LA, Sherr JL, Clements M, Neyman A, Evans-Molina C, Sims EK, Messer LH, Ekhlaspour L, McDonough R, Van Name M, Rojas D, Beasley S, DuBose S, Kollman C, Moran A, Group CLS. Effect of Verapamil on Pancreatic Beta Cell Function in Newly Diagnosed Pediatric Type 1 Diabetes: A Randomized Clinical Trial. *JAMA*. 2023;329(12):990-9.
45. Gu Y, Jia X, Vartak T, Miao D, Dong F, Jerram ST, Rewers M, Ferrara A, Lawrence JM, Yu L, Leslie RD, Action Lc, the Diabetes in Young Adults Study G. Improving clinical utility of GAD65 autoantibodies by electrochemiluminescence assay and clinical phenotype when identifying autoimmune adult-onset diabetes. *Diabetologia*. 2021;64(9):2052-60.

832 46. Jones AG, McDonald TJ, Shields BM, Hagopian W, Hattersley AT. Latent Autoimmune
833 Diabetes of Adults (LADA) Is Likely to Represent a Mixed Population of Autoimmune (Type 1) and
834 Nonautoimmune (Type 2) Diabetes. *Diabetes care*. 2021.

835 47. Balasubramanyam A, Garza G, Rodriguez L, Hampe CS, Gaur L, Lernmark A, Maldonado MR.
836 Accuracy and predictive value of classification schemes for ketosis-prone diabetes. *Diabetes care*.
837 2006;29(12):2575-9.

838 48. Li X, Chen Y, Xie Y, Xiang Y, Yan X, Huang G, Zhou Z. Decline Pattern of Beta-cell Function in
839 Adult-onset Latent Autoimmune Diabetes: an 8-year Prospective Study. *The Journal of clinical*
840 *endocrinology and metabolism*. 2020;105(7).

841 49. Eason RJ, Thomas NJ, Hill AV, Knight BA, Carr A, Hattersley AT, McDonald TJ, Shields BM,
842 Jones AG, StartRight Study G, StartRight Study G. Routine Islet Autoantibody Testing in Clinically
843 Diagnosed Adult-Onset Type 1 Diabetes Can Help Identify Misclassification and the Possibility of
844 Successful Insulin Cessation. *Diabetes care*. 2022.

845 50. Hannelius U, Beam CA, Ludvigsson J. Efficacy of GAD-alum immunotherapy associated with
846 HLA-DR3-DQ2 in recently diagnosed type 1 diabetes. *Diabetologia*. 2020;63(10):2177-81.

847 51. Buzzetti R, Tuomi T, Mauricio D, Pietropaolo M, Zhou Z, Pozzilli P, Leslie RD. Management of
848 Latent Autoimmune Diabetes in Adults: A Consensus Statement From an International Expert Panel.
849 *Diabetes*. 2020;69(10):2037-47.

850 52. Sims EK, Bahnson HT, Nyalwidhe J, Haataja L, Davis AK, Speake C, DiMeglio LA, Blum J,
851 Morris MA, Mirmira RG, Nadler J, Mastracci TL, Marcovina S, Qian WJ, Yi L, Swensen AC, Yip-
852 Schneider M, Schmidt CM, Considine RV, Arvan P, Greenbaum CJ, Evans-Molina C. Proinsulin
853 Secretion Is a Persistent Feature of Type 1 Diabetes. *Diabetes care*. 2019;42(2):258-64.

854 53. Davies MJ, Aroda VR, Collins BS, Gabbay RA, Green J, Maruthur NM, Rosas SE, Del Prato S,
855 Mathieu C, Mingrone G, Rossing P, Tankova T, Tsapas A, Buse JB. Management of hyperglycaemia in
856 type 2 diabetes, 2022. A consensus report by the American Diabetes Association (ADA) and the
857 European Association for the Study of Diabetes (EASD). *Diabetologia*. 2022;65(12):1925-66.

858 54. Ahlqvist E, Storm P, Karajamaki A, Martinell M, Dorkhan M, Carlsson A, Vikman P, Prasad RB,
859 Aly DM, Almgren P, Wessman Y, Shaat N, Spegel P, Mulder H, Lindholm E, Melander O, Hansson O,
860 Malmqvist U, Lernmark A, Lahti K, Forsen T, Tuomi T, Rosengren AH, Groop L. Novel subgroups of
861 adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six
862 variables. *Lancet Diabetes Endocrinol*. 2018;6(5):361-9.

863 55. Zaharia OP, Strassburger K, Strom A, Bonhof GJ, Karusheva Y, Antoniou S, Bodis K, Markgraf
864 DF, Burkart V, Mussig K, Hwang JH, Asplund O, Groop L, Ahlqvist E, Seissler J, Nawroth P, Kopf S,
865 Schmid SM, Stumvoll M, Pfeiffer AFH, Kabisch S, Tselmin S, Haring HU, Ziegler D, Kuss O, Szendroedi
866 J, Roden M, German Diabetes Study G. Risk of diabetes-associated diseases in subgroups of patients
867 with recent-onset diabetes: a 5-year follow-up study. *Lancet Diabetes Endocrinol*. 2019;7(9):684-94.

868 56. Mansour Aly D, Dwivedi OP, Prasad RB, Karajamaki A, Hjort R, Thangam M, Akerlund M,
869 Mahajan A, Udler MS, Florez JC, McCarthy MI, Regeneron Genetics C, Brosnan J, Melander O,
870 Carlsson S, Hansson O, Tuomi T, Groop L, Ahlqvist E. Genome-wide association analyses highlight
871 etiological differences underlying newly defined subtypes of diabetes. *Nature genetics*.
872 2021;53(11):1534-42.

873 57. Zaharia OP, Strassburger K, Knebel B, Kupriyanova Y, Karusheva Y, Wolkersdorfer M, Bodis K,
874 Markgraf DF, Burkart V, Hwang JH, Kotzka J, Al-Hasani H, Szendroedi J, Roden M, Group GDS. Role of
875 Patatin-Like Phospholipase Domain-Containing 3 Gene for Hepatic Lipid Content and Insulin
876 Resistance in Diabetes. *Diabetes care*. 2020;43(9):2161-8.

877 58. Zou X, Zhou X, Zhu Z, Ji L. Novel subgroups of patients with adult-onset diabetes in Chinese
878 and US populations. *Lancet Diabetes Endocrinol*. 2019;7(1):9-11.

879 59. Anjana RM, Baskar V, Nair ATN, Jebarani S, Siddiqui MK, Pradeepa R, Unnikrishnan R, Palmer
880 C, Pearson E, Mohan V. Novel subgroups of type 2 diabetes and their association with microvascular
881 outcomes in an Asian Indian population: a data-driven cluster analysis: the INSPIRED study. *BMJ*
882 *Open Diabetes Res Care*. 2020;8(1).

60. McCarthy MI. Painting a new picture of personalised medicine for diabetes. *Diabetologia*. 2017;60(5):793-9.

61. Everitt B, Landau S, Leese M, Stahl D. Cluster analysis. Chichester: Wiley; 2011.

62. Wesolowska-Andersen A, Brorsson CA, Bizzotto R, Mari A, Tura A, Koivula R, Mahajan A, Vinuela A, Tajes JF, Sharma S, Haid M, Prehn C, Artati A, Hong MG, Musholt PB, Kurbasic A, De Masi F, Tsigos K, Pedersen HK, Gudmundsdottir V, Thomas CE, Banasik K, Jennison C, Jones A, Kennedy G, Bell J, Thomas L, Frost G, Thomsen H, Allin K, Hansen TH, Vestergaard H, Hansen T, Rutters F, Elders P, t'Hart L, Bonnefond A, Canouil M, Brage S, Kokkola T, Heggie A, McEvoy D, Hattersley A, McDonald T, Teare H, Ridderstrale M, Walker M, Forgie I, Giordano GN, Froguel P, Pavo I, Ruetten H, Pedersen O, Dermitzakis E, Franks PW, Schwenk JM, Adamski J, Pearson E, McCarthy MI, Brunak S, Consortium ID. Four groups of type 2 diabetes contribute to the etiological and clinical heterogeneity in newly diagnosed individuals: An IMI DIRECT study. *Cell Rep Med*. 2022;3(1):100477.

63. Nair ATN, Wesolowska-Andersen A, Brorsson C, Rajendrakumar AL, Hapca S, Gan S, Dawed AY, Donnelly LA, McCrimmon R, Doney ASF, Palmer CNA, Mohan V, Anjana RM, Hattersley AT, Dennis JM, Pearson ER. Heterogeneity in phenotype, disease progression and drug response in type 2 diabetes. *Nat Med*. 2022;28(5):982-8.

64. Udler MS, Kim J, von Grotthuss M, Bonas-Guarch S, Cole JB, Chiou J, Boehnke M, Laakso M, Atzmon G, Glaser B, Mercader JM, Gaulton K, Flannick J, Getz G, Florez JC. Type 2 diabetes genetic loci informed by multi-trait associations point to disease mechanisms and subtypes: A soft clustering analysis. *PLoS medicine*. 2018;15(9):e1002654.

65. DiCorpo D, LeClair J, Cole JB, Sarnowski C, Ahmadizar F, Bielak LF, Blokstra A, Bottinger EP, Chaker L, Chen YI, Chen Y, de Vries PS, Faquih T, Ghanbari M, Gudmundsdottir V, Guo X, Hasbani NR, Ibi D, Ikram MA, Kavousi M, Leonard HL, Leong A, Mercader JM, Morrison AC, Nadkarni GN, Nalls MA, Noordam R, Preuss M, Smith JA, Trompet S, Vissink P, Yao J, Zhao W, Boerwinkle E, Goodarzi MO, Gudnason V, Jukema JW, Kardina SLR, Loos RJF, Liu CT, Manning AK, Mook-Kanamori D, Pankow JS, Picavet HJ, Sattar N, Simonsick EM, Verschuren WMM, Willems van Dijk K, Florez JC, Rotter JI, Meigs JB, Dupuis J, Udler MS. Type 2 Diabetes Partitioned Polygenic Scores Associate With Disease Outcomes in 454,193 Individuals Across 13 Cohorts. *Diabetes care*. 2022;45(3):674-83.

66. Suzuki K, Hatzikotoulas K, Southam L, Taylor HJ, Yin X, Lorenz KM, Mandla R, Huerta-Chagoya A, Rayner NW, Bocher O, Ana Luiza de SVA, Sonehara K, Namba S, Lee SSK, Preuss MH, Petty LE, Schroeder P, Vanderwerff B, Kals M, Bragg F, Lin K, Guo X, Zhang W, Yao J, Kim YJ, Graff M, Takeuchi F, Nano J, Lamri A, Nakatochi M, Moon S, Scott RA, Cook JP, Lee JJ, Pan I, Taliun D, Parra EJ, Chai JF, Bielak LF, Tabara Y, Hai Y, Thorleifsson G, Grarup N, Sofer T, Wuttke M, Sarnowski C, Gieger C, Nounsou D, Trompet S, Kwak SH, Long J, Sun M, Tong L, Chen WM, Nongmaithem SS, Noordam R, Lim VJY, Tam CHT, Joo YY, Chen CH, Raffield LM, Prins BP, Nicolas A, Yanek LR, Chen G, Brody JA, Kabagambe E, An P, Xiang AH, Choi HS, Cade BE, Tan J, Elaine Broadway K, Williamson A, Kamali Z, Cui J, Adair LS, Adeyemo A, Aguilar-Salinas CA, Ahluwalia TS, Anand SS, Bertoni A, Bork-Jensen J, Brandslund I, Buchanan TA, Burant CF, Butterworth AS, Canouil M, Chan JCN, Chang LC, Chee ML, Chen J, Chen SH, Chen YT, Chen Z, Chuang LM, Cushman M, Danesh J, Das SK, Janaka de Silva H, Dedoussis G, Dimitrov L, Doumatey AP, Du S, Duan Q, Eckardt KU, Emery LS, Evans DS, Evans MK, Fischer K, Floyd JS, Ford I, Franco OH, Frayling TM, Freedman BI, Genter P, Gerstein HC, Giedraitis V, Gonzalez-Villalpando C, Gonzalez-Villalpando ME, Gordon-Larsen P, Gross M, Guare LA, Hackinger S, Han S, Hattersley AT, Herder C, Horikoshi M, Howard AG, Hsueh W, Huang M, Huang W, Hung YJ, Hwang MY, Hwu CM, Ichihara S, Ikram MA, Ingelsson M, Islam MT, Isono M, Jang HM, Jasmine F, Jiang G, Jonas JB, Jorgensen T, Kandeel FR, Kasturiratne A, Katsuya T, Kaur V, Kawaguchi T, Keaton JM, Kho AN, Khor CC, Kibriya MG, Kim DH, Kronenberg F, Kuusisto J, Lall K, Lange LA, Lee KM, Lee MS, Lee NR, Leong A, Li L, Li Y, Li-Gao R, Lithgart S, Lindgren CM, Linneberg A, Liu CT, Liu J, Locke AE, Louie T, Luan J, Luk AO, Luo X, Lv J, Lynch JA, Lyssenko V, Maeda S, Mamakou V, Mansuri SR, Matsuda K, Meitinger T, Metspalu A, Mo H, Morris AD, Nadler JL, Nalls MA, Nayak U, Ntalla I, Okada Y, Orozco L, Patel SR, Patil S, Pei P, Pereira MA, Peters A, Pirie FJ, Polikowsky HG, Porneala B, Prasad G, Rasmussen-Torvik LJ, Reiner AP, Roden M, Rohde R, Roll K, Sabanayagam C, Sandow K,

934 Sankareswaran A, Sattar N, Schonherr S, Shahriar M, Shen B, Shi J, Shin DM, Shojima N, Smith JA, So
 935 WY, Stancakova A, Steinthorsdottir V, Stilp AM, Strauch K, Taylor KD, Thorand B, Thorsteinsdottir U,
 936 Tomlinson B, Tran TC, Tsai FJ, Tuomilehto J, Tusie-Luna T, Udler MS, Valladares-Salgado A, van Dam
 937 RM, van Klinken JB, Varma R, Wachter-Rodarte N, Wheeler E, Wickremasinghe AR, van Dijk KW, Witte
 938 DR, Yajnik CS, Yamamoto K, Yamamoto K, Yoon K, Yu C, Yuan JM, Yusuf S, Zawistowski M, Zhang L,
 939 Zheng W, Va Million Veteran Program AGDIJ, Project BJ, BioBank PM, Center RG, Consortium E,
 940 International Consortium for Blood P, Meta-Analyses of G, Insulin-Related Traits C, Raffel LJ, Igase M,
 941 Ipp E, Redline S, Cho YS, Lind L, Province MA, Fornage M, Hanis CL, Ingelsson E, Zonderman AB, Psaty
 942 BM, Wang YX, Rotimi CN, Becker DM, Matsuda F, Liu Y, Yokota M, Kardia SLR, Peyser PA, Pankow JS,
 943 Engert JC, Bonnefond A, Froguel P, Wilson JG, Sheu WHH, Wu JY, Geoffrey Hayes M, Ma RCW, Wong
 944 TY, Mook-Kanamori DO, Tuomi T, Chandak GR, Collins FS, Bharadwaj D, Pare G, Sale MM, Ahsan H,
 945 Motala AA, Shu XO, Park KS, Jukema JW, Cruz M, Chen YI, Rich SS, McKean-Cowdin R, Grallert H,
 946 Cheng CY, Ghanbari M, Tai ES, Dupuis J, Kato N, Laakso M, Kottgen A, Koh WP, Bowden DW, Palmer
 947 CNA, Kooner JS, Kooperberg C, Liu S, North KE, Saleheen D, Hansen T, Pedersen O, Wareham NJ, Lee
 948 J, Kim BJ, Millwood IY, Walters RG, Stefansson K, Goodarzi MO, Mohlke KL, Langenberg C, Haiman
 949 CA, Loos RJF, Florez JC, Rader DJ, Ritchie MD, Zollner S, Magi R, Denny JC, Yamauchi T, Kadowaki T,
 950 Chambers JC, Ng MCY, Sim X, Below JE, Tsao PS, Chang KM, McCarthy MI, Meigs JB, Mahajan A,
 951 Spracklen CN, Mercader JM, Boehnke M, Rotter JI, Vujkovic M, Voight BF, Morris AP, Zeggini E.
 952 Multi-ancestry genome-wide study in >2.5 million individuals reveals heterogeneity in mechanistic
 953 pathways of type 2 diabetes and complications. medRxiv. 2023.
 954 67. Zhou K, Donnelly LA, Morris AD, Franks PW, Jennison C, Palmer CN, Pearson ER. Clinical and
 955 genetic determinants of progression of type 2 diabetes: a DIRECT study. Diabetes care.
 956 2014;37(3):718-24.
 957 68. Jiang G, Luk AO, Tam CHT, Lau ES, Ozaki R, Chow EYK, Kong APS, Lim CKP, Lee KF, Siu SC, Hui
 958 G, Tsang CC, Lau KP, Leung JYY, Tsang MW, Kam G, Lau IT, Li JK, Yeung VT, Lau E, Lo S, Fung SKS,
 959 Cheng YL, Chow CC, Pearson ER, So WY, Chan JCN, Ma RCW. Obesity, clinical, and genetic predictors
 960 for glycemic progression in Chinese patients with type 2 diabetes: A cohort study using the Hong
 961 Kong Diabetes Register and Hong Kong Diabetes Biobank. PLoS medicine. 2020;17(7):e1003209.
 962 69. Fan B, Lim CKP, Poon EWM, Lau ESH, Wu H, Yang A, Shi M, Tam CHT, Wong SYS, Lee EK,
 963 Wang MHT, Chu NHS, Ozaki R, Kong APS, Chow E, Ma RCW, Luk AOY, Chan JCN. Differential
 964 Associations of GAD Antibodies (GADA) and C-Peptide With Insulin Initiation, Glycemic Responses,
 965 and Severe Hypoglycemia in Patients Diagnosed With Type 2 Diabetes. Diabetes care. 2023.
 966 70. Dennis JM, Henley WE, Weedon MN, Lonergan M, Rodgers LR, Jones AG, Hamilton WT,
 967 Sattar N, Janmohamed S, Holman RR, Pearson ER, Shields BM, Hattersley AT. Sex and BMI Alter the
 968 Benefits and Risks of Sulfonylureas and Thiazolidinediones in Type 2 Diabetes: A Framework for
 969 Evaluating Stratification Using Routine Clinical and Individual Trial Data. Diabetes care.
 970 2018;41(9):1844-53.
 971 71. Dennis JM, Young KG, McGovern AP, Mateen BA, Vollmer SJ, Simpson MD, Henley WE,
 972 Holman RR, Sattar N, Pearson ER, Hattersley AT, Jones AG, Shields BM, consortium M. Development
 973 of a treatment selection algorithm for SGLT2 and DPP-4 inhibitor therapies in people with type 2
 974 diabetes: a retrospective cohort study. Lancet Digit Health. 2022;4(12):e873-e83.
 975 72. Shields BM, Dennis JM, Angwin CD, Warren F, Henley WE, Farmer AJ, Sattar N, Holman RR,
 976 Jones AG, Pearson ER, Hattersley AT, TriMaster Study g. Patient stratification for determining
 977 optimal second-line and third-line therapy for type 2 diabetes: the TriMaster study. Nat Med. 2022.
 978 73. Dawed AY, Zhou K, Pearson ER. Pharmacogenetics in type 2 diabetes: influence on response
 979 to oral hypoglycemic agents. Pharmgenomics Pers Med. 2016;9:17-29.
 980 74. Alyafei F, Soliman A, Alkhalaf F, Sabt A, De Sanctis V, Waseef R, Elsayed N. Incidence of type
 981 1 and type 2 diabetes, between 2012-2016, among children and adolescents in Qatar. Acta Biomed.
 982 2018;89(S5):7-10.
 983 75. Mayer-Davis EJ, Dabelea D, Lawrence JM. Incidence Trends of Type 1 and Type 2 Diabetes
 984 among Youths, 2002-2012. The New England journal of medicine. 2017;377(3):301.

76. Candler TP, Mahmoud O, Lynn RM, Majbar AA, Barrett TG, Shield JPH. Continuing rise of Type 2 diabetes incidence in children and young people in the UK. *Diabet Med*. 2018;35(6):737-44.
77. Amed S, Islam N, Sutherland J, Reimer K. Incidence and prevalence trends of youth-onset type 2 diabetes in a cohort of Canadian youth: 2002-2013. *Pediatric diabetes*. 2018;19(4):630-6.
78. Luk AOY, Ke C, Lau ESH, Wu H, Goggins W, Ma RCW, Chow E, Kong APS, So WY, Chan JCN. Secular trends in incidence of type 1 and type 2 diabetes in Hong Kong: A retrospective cohort study. *PLoS medicine*. 2020;17(2):e1003052.
79. Zeitler P, Hirst K, Pyle L, Linder B, Copeland K, Arslanian S, Cuttler L, Nathan DM, Tollefsen S, Wilfley D, Kaufman F. A clinical trial to maintain glycemic control in youth with type 2 diabetes. *The New England journal of medicine*. 2012;366(24):2247-56.
80. Shah AS, Nadeau KJ. The changing face of paediatric diabetes. *Diabetologia*. 2020;63(4):683-91.
81. Astudillo M, Tosur M, Castillo B, Rafaey A, Siller AF, Nieto J, Sisley S, McKay S, Nella AA, Balasubramanyam A, Bacha F, Redondo MJ. Type 2 diabetes in prepubertal children. *Pediatric diabetes*. 2021;22(7):946-50.
82. Dabelea D, D'Agostino RB, Jr., Mason CC, West N, Hamman RF, Mayer-Davis EJ, Maahs D, Klingensmith G, Knowler WC, Nadeau K. Development, validation and use of an insulin sensitivity score in youths with diabetes: the SEARCH for Diabetes in Youth study. *Diabetologia*. 2011;54(1):78-86.
83. Mottl AK, Lauer A, Dabelea D, Maahs DM, D'Agostino RB, Jr., Dolan LM, Gilliam LK, Lawrence JM, Rodriguez B, Marcovina SM, Imperatore G, Kanakatti Shankar R, Afkarian M, Reynolds K, Liese AD, Mauer M, Mayer-Davis EJ, Group SfdiYS. Albuminuria according to status of autoimmunity and insulin sensitivity among youth with type 1 and type 2 diabetes. *Diabetes care*. 2013;36(11):3633-8.
84. Pihoker C, Gilliam LK, Ellard S, Dabelea D, Davis C, Dolan LM, Greenbaum CJ, Imperatore G, Lawrence JM, Marcovina SM, Mayer-Davis E, Rodriguez BL, Steck AK, Williams DE, Hattersley AT, Group SfdiYS. Prevalence, characteristics and clinical diagnosis of maturity onset diabetes of the young due to mutations in HNF1A, HNF4A, and glucokinase: results from the SEARCH for Diabetes in Youth. *The Journal of clinical endocrinology and metabolism*. 2013;98(10):4055-62.
85. Arslanian S, El Ghormli L, Haymond MH, Chan CL, Chernaused SD, Gandica RG, Gubitosi-Klug R, Levitsky LL, Siska M, Willi SM, Group TS. Beta cell function and insulin sensitivity in obese youth with maturity onset diabetes of youth mutations vs type 2 diabetes in TODAY: Longitudinal observations and glycemic failure. *Pediatric diabetes*. 2020;21(4):575-85.
86. Nadeau KJ, Regensteiner JG, Bauer TA, Brown MS, Dorosz JL, Hull A, Zeitler P, Draznin B, Reusch JE. Insulin resistance in adolescents with type 1 diabetes and its relationship to cardiovascular function. *The Journal of clinical endocrinology and metabolism*. 2010;95(2):513-21.
87. Bjornstad P, Schafer M, Truong U, Cree-Green M, Pyle L, Baumgartner A, Garcia Reyes Y, Maniatis A, Nayak S, Wadwa RP, Browne LP, Reusch JEB, Nadeau KJ. Metformin Improves Insulin Sensitivity and Vascular Health in Youth With Type 1 Diabetes Mellitus. *Circulation*. 2018;138(25):2895-907.
88. Cree-Green M, Bergman BC, Cengiz E, Fox LA, Hannon TS, Miller K, Nathan B, Pyle L, Kahn D, Tansey M, Tichy E, Tsalikian E, Libman I, Nadeau KJ. Metformin Improves Peripheral Insulin Sensitivity in Youth With Type 1 Diabetes. *The Journal of clinical endocrinology and metabolism*. 2019;104(8):3265-78.
89. Tell SS, Schafer M, Vigers T, Baumgartner AD, Lyon E, Gross S, Polsky S, Snell-Bergeon JK, Schauer IE, Nadeau KJ. Bromocriptine quick-release as adjunct therapy in youth and adults with type 1 diabetes: A randomized, placebo-controlled crossover study. *Diabetes, obesity & metabolism*. 2022;24(11):2148-58.
90. Dabelea D, Kinney G, Snell-Bergeon JK, Hokanson JE, Eckel RH, Ehrlich J, Garg S, Hamman RF, Rewers M, Coronary Artery Calcification in Type 1 Diabetes S. Effect of type 1 diabetes on the gender difference in coronary artery calcification: a role for insulin resistance? The Coronary Artery Calcification in Type 1 Diabetes (CACTI) Study. *Diabetes*. 2003;52(11):2833-9.

91. Ginnard OZB, Redondo MJ, Lyons SK. Diabetes of Unclear Type in an Adolescent Boy With Multiple Islet-cell Autoantibody Positivity Successfully Managed With Glucagon-like Peptide-1 Receptor Agonist Alone: A Case Report. *Can J Diabetes*. 2022.
92. Rise Consortium. Impact of Insulin and Metformin Versus Metformin Alone on beta-Cell Function in Youth With Impaired Glucose Tolerance or Recently Diagnosed Type 2 Diabetes. *Diabetes care*. 2018;41(8):1717-25.
93. Fazeli Farsani S, van der Aa MP, van der Vorst MM, Knibbe CA, de Boer A. Global trends in the incidence and prevalence of type 2 diabetes in children and adolescents: a systematic review and evaluation of methodological approaches. *Diabetologia*. 2013;56(7):1471-88.
94. Nadeau KJ, Anderson BJ, Berg EG, Chiang JL, Chou H, Copeland KC, Hannon TS, Huang TT, Lynch JL, Powell J, Sellers E, Tamborlane WV, Zeitler P. Youth-Onset Type 2 Diabetes Consensus Report: Current Status, Challenges, and Priorities. *Diabetes care*. 2016;39(9):1635-42.
95. Kahn SE, Haffner SM, Heise MA, Herman WH, Holman RR, Jones NP, Kravitz BG, Lachin JM, O'Neill MC, Zinman B, Viberti G. Glycemic durability of rosiglitazone, metformin, or glyburide monotherapy. *The New England journal of medicine*. 2006;355(23):2427-43.
96. Dabelea D, Stafford JM, Mayer-Davis EJ, D'Agostino R, Jr., Dolan L, Imperatore G, Linder B, Lawrence JM, Marcovina SM, Mottl AK, Black MH, Pop-Busui R, Saydah S, Hamman RF, Pihoker C. Association of Type 1 Diabetes vs Type 2 Diabetes Diagnosed During Childhood and Adolescence With Complications During Teenage Years and Young Adulthood. *Jama*. 2017;317(8):825-35.
97. TODAY Study Group. Rapid rise in hypertension and nephropathy in youth with type 2 diabetes: the TODAY clinical trial. *Diabetes care*. 2013;36(6):1735-41.
98. Kahkoska AR, Isom S, Divers J, Mayer-Davis EJ, Dolan L, Shah AS, Afkarian M, Pettitt DJ, Lawrence JM, Marcovina S, Saydah SH, Dabelea D, Maahs DM, Mottl AK. The early natural history of albuminuria in young adults with youth-onset type 1 and type 2 diabetes. *Journal of diabetes and its complications*. 2018;32(12):1160-8.
99. Kershner AK, Daniels SR, Imperatore G, Palla SL, Pettitt DB, Pettitt DJ, Marcovina S, Dolan LM, Hamman RF, Liese AD, Pihoker C, Rodriguez BL. Lipid abnormalities are prevalent in youth with type 1 and type 2 diabetes: the SEARCH for Diabetes in Youth Study. *The Journal of pediatrics*. 2006;149(3):314-9.
100. Levitt Katz L, Gidding SS, Bacha F, Hirst K, McKay S, Pyle L, Lima JA. Alterations in left ventricular, left atrial, and right ventricular structure and function to cardiovascular risk factors in adolescents with type 2 diabetes participating in the TODAY clinical trial. *Pediatric diabetes*. 2015;16(1):39-47.
101. Ferrat LA, Vehik K, Sharp SA, Lernmark A, Rewers MJ, She JX, Ziegler AG, Toppari J, Akolkar B, Krischer JP, Weedon MN, Oram RA, Hagopian WA, Group TS, Committees. A combined risk score enhances prediction of type 1 diabetes among susceptible children. *Nat Med*. 2020;26(8):1247-55.
102. Redondo MJ, Geyer S, Steck AK, Sharp S, Wentworth JM, Weedon MN, Antinozzi P, Sosenko J, Atkinson M, Pugliese A, Oram RA, Type 1 Diabetes TrialNet Study G. A Type 1 Diabetes Genetic Risk Score Predicts Progression of Islet Autoimmunity and Development of Type 1 Diabetes in Individuals at Risk. *Diabetes care*. 2018;41(9):1887-94.
103. DeFronzo RA. Banting Lecture. From the triumvirate to the ominous octet: a new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes*. 2009;58(4):773-95.
104. Carrasco-Zanini J, Pietzner M, Lindbohm JV, Wheeler E, Oerton E, Kerrison N, Simpson M, Westacott M, Drolet D, Kivimaki M, Ostroff R, Williams SA, Wareham NJ, Langenberg C. Proteomic signatures for identification of impaired glucose tolerance. *Nat Med*. 2022;28(11):2293-300.
105. Zhu M, Xu K, Chen Y, Gu Y, Zhang M, Luo F, Liu Y, Gu W, Hu J, Xu H, Xie Z, Sun C, Li Y, Sun M, Xu X, Hsu HT, Chen H, Fu Q, Shi Y, Xu J, Ji L, Liu J, Bian L, Zhu J, Chen S, Xiao L, Li X, Jiang H, Shen M, Huang Q, Fang C, Li X, Huang G, Fan J, Jiang Z, Jiang Y, Dai J, Ma H, Zheng S, Cai Y, Dai H, Zheng X, Zhou H, Ni S, Jin G, She JX, Yu L, Polychronakos C, Hu Z, Zhou Z, Weng J, Shen H, Yang T. Identification of Novel T1D Risk Loci and Their Association With Age and Islet Function at Diagnosis in

1086 Autoantibody-Positive T1D Individuals: Based on a Two-Stage Genome-Wide Association Study.
1087 Diabetes care. 2019;42(8):1414-21.

1088 106. Cousminer DL, Ahlqvist E, Mishra R, Andersen MK, Chesi A, Hawa MI, Davis A, Hodge KM,
1089 Bradfield JP, Zhou K, Guy VC, Akerlund M, Wod M, Fritsche LG, Vestergaard H, Snyder J, Hojlund K,
1090 Linneberg A, Karajamaki A, Brandslund I, Kim CE, Witte D, Sorgjerd EP, Brillon DJ, Pedersen O, Beck-
1091 Nielsen H, Grarup N, Pratley RE, Rickels MR, Vella A, Ovalle F, Melander O, Harris RI, Varvel S, Grill
1092 VER, Hakonarson H, Froguel P, Lonsdale JT, Mauricio D, Schloot NC, Khunti K, Greenbaum CJ, Asvold
1093 BO, Yderstraede KB, Pearson ER, Schwartz S, Voight BF, Hansen T, Tuomi T, Boehm BO, Groop L,
1094 Leslie RD, Grant SFA. First Genome-Wide Association Study of Latent Autoimmune Diabetes in Adults
1095 Reveals Novel Insights Linking Immune and Metabolic Diabetes. Diabetes care. 2018;41(11):2396-
1096 403.

1097 107. Lynam A, McDonald T, Hill A, Dennis J, Oram R, Pearson E, Weedon M, Hattersley A, Owen K,
1098 Shields B, Jones A. Development and validation of multivariable clinical diagnostic models to identify
1099 type 1 diabetes requiring rapid insulin therapy in adults aged 18-50 years. BMJ Open.
1100 2019;9(9):e031586.

1101