

# "PRINCIPLES OF PHYLOGENETICS: ECOLOGY AND EVOLUTION"

*Integrative Biology 200B*

University of California, Berkeley

Spring 2016

D.D. Ackerly

April 13, 2016. **Adaptive radiation**

## Reading:

Hodges, S. 1995. Spurring plant diversification: Are floral nectar spurs a key innovation? *Proc. Roy. Soc. London Ser. B* 262:343-348.

Harmon, L. J., J. A. Schulte, II, A. Larson, and J. B. Losos. 2003. Tempo and mode of evolutionary radiation in Iguanian lizards. *Science* 301:961-964.

Ackerly, D. D. 2009. Conservatism and diversification of plant functional traits: Evolutionary rates versus phylogenetic signal. *PNAS* 106:19699–19706.

### 1. Central questions of adaptive radiation research:

- What is an adaptive radiation? Is it something special. Is it a category or a continuum (i.e. some clades are adaptive radiations and some aren't).
- What is the link between trait and lineage diversification? One can imagine a 3x2 matrix with 6 different scenarios/phenomena. Fill in the blanks!

	Rate of lineage diversification	
Trait evolution	LOW	HIGH
Ancestral trait innovation		
Low rate of trait diversification		
High rate of trait diversification		

### 2. Key innovation and diversification

The case of nectar spurs in *Aquilegia* (S. Hodges)

Floral symmetry and sister clade diversification in angiosperms (Sargent)

BISSE – trait dependent diversification (Maddison)

### 3. Adaptive radiation: evolution of disparity and diversification

*Anolis* lizards:

Ecomorphs and phylogeny (Losos)


Macrohabitat radiations within ecomorphs (Glor)

Disparity vs speciation (Harmon)


#### 4. Testing for rate variation in continuous trait evolution

O'Meara – brownie: ML test for 2 vs. 1 rate parameter

Martin and Wainwright (2011) application to divergence in feeding morphology



Hodges and Arnold 1995 Proc Roy Soc



Sargent 2004 Proc Roy Soc Lond B






FIG. 1. An example tree and character distribution, simulated with biased speciation. Rate of speciation with state 1 (black) five times higher than rate of speciation with state 0 (white). States at internal nodes are parsimony reconstructions, not the actual ancestral states as simulated.

## Maddison 2006 Evolution




## Maddison et al. 2007 Evolution



**Fig. 1.** The posterior probabilities that diploids exhibit higher rates of (A) diversification, (B) speciation, and (C) extinction than polyploids.

Mayrose et al. 2011



Losos et al. 1998 Science




Fig. 1. Lineage accumulation curves for four lizard taxa compared with expectations under the null pure-birth model. Time is expressed as a proportion of the total time since the first cladogenetic event inferred for the taxon. Only the first two-thirds of each phylogeny is shown. Expected curves are obtained using an exponential model with the first branching in a clade set to time = 0 and the number of extant species (Table 1) set to time = 1. Solid line, actual number of reconstructed lineages for the clade; dashed line, expected number of lineages under the null pure-birth model.







Fig. 3. Relationship between LDI and MDI. Data show a highly significant negative correlation (19) ( $r = -0.998$ ,  $P = 0.004$ ). A variety of analyses indicate that the among-clade relationship between LDI and MDI is well-supported and robust to relaxation of assumptions concerning mode of character evolution, sampling, and extinction (19) (figs. S6 and S7; 95% confidence limits were established by calculating MDI separately for each of the 1000 simulations of character evolution that were used to calculate MDI for each clade). This result is also significant if  $\gamma$  (corrected for incomplete sampling by subtracting the mean of the  $\gamma$  values in the MCCR test simulations and dividing by the standard deviation of these simulation  $\gamma$  values) and  $\delta$  values are used for each clade (standardized  $\gamma$  versus  $\delta_{MLE}$ ;  $r = 0.97$ ,  $P = 0.03$ ; in this analysis, the correlation is positive because more negative values of  $\gamma$  reflect a slowdown in diversification, corresponding to larger values of our LDI statistic).





## O'Meara et al. 2006 Evolution



**Figure 4.** Relative rates of morphological diversification (mean  $\pm$  SE) on the (A) first and (B) second principal component axes relative to clade age. (C) The morphological diversification rate-space (mean  $\pm$  SE) on the first and second principal component axes. Observed rates are indicated for the Chichancanab (○) and San Salvador (●) clades, and all other two-species (●), three-species (●), and four-species (●) *Cyprinodon* clades measured; note that points are not statistically independent due to nested clade structure and shared histories. Means and standard errors were calculated by repeating rate analyses for 100 trees sampled from the stationary MCMC distribution. (A, B) Simulated relative rates of morphological diversification (mean  $\pm$  2SD) under a Brownian motion model for 1000 iterations are shown in gray (●). (C) Each simulation is indicated by one point (●) in the morphological diversification rate-space.

## Martin and Wainwright 2011 Evolution

**Reading list:**

- Ackerly, D. D., D. W. Schilck, and C. O. Webb. 2006. Niche evolution and adaptive radiation: testing the order of trait divergence. *Ecology* 87:S50-S61.
- Ackerly, D. D. 2009. Conservatism and diversification of plant functional traits: Evolutionary rates versus phylogenetic signal. *Proceedings of the National Academy of Sciences USA* 106:19699-19706. doi: 10.1073/pnas.0901635106.
- Becerra, J. X. 2005. Timing the origin and expansion of the Mexican tropical dry forest. *Proceedings of the National Academy of Sciences of the United States of America* 102:10919-10923.
- FitzJohn, R. G., Maddison, W. P., & Otto, S. P. (2009). Estimating trait-dependent speciation and extinction rates from incompletely resolved phylogenies. *Systematic Biology*, 58(6), 595-611.
- FitzJohn, R. G. (2010). Quantitative traits and diversification. *Systematic Biology*, 59(6), 619-633.
- Glor, R. E., J. J. Kolbe, R. Powell, A. Larson, and J. B. Losos. 2003. Phylogenetic analysis of ecological and morphological diversification in Hispaniolan trunk-ground anoles (anolis cybotes group). *Evolution* 57:2383-2397.
- Harmon, L. J., J. A. Schulte, II, A. Larson, and J. B. Losos. 2003. Tempo and mode of evolutionary radiation in Iguanian lizards. *Science* 301:961-964.
- Harvey, P. H., and A. Rambaut. 2000. Comparative analyses for adaptive radiations. *PhilTransRoySocLondSerB* 355:1599-1605.
- Hodges, S. 1995. Spurring plant diversification: Are floral nectar spurs a key innovation? *Proc. Roy. Soc. London Ser. B* 262:343-348.
- Hodges, S. 1997. Floral nectar spurs and diversification. *Int. J. Plant Sci.* 158:S81-S88.
- Losos, J. 1992. The evolution of convergent structure in Caribbean *Anolis* communities. *SystBiol.* 41:403-420.
- Losos, J. B., and D. B. Miles. 2002. Testing the hypothesis that a clade has adaptively radiated: iguanid lizard clades as a case study. *Amer. Nat.* 160:147-157.
- Losos, J. B., M. Leal, R. E. Glor, K. de Queiroz, P. E. Hertz, L. R. Schettino, A. C. Lara, T. R. Jackman, and A. Larson. 2003. Niche lability in the evolution of a Caribbean lizard community. *Nature* 424:542-545.
- Losos, J., T. Jackman, A. Larson, K. de Queiroz, and L. Rodriguez-Schettino. 1998. Contingency and determinism in replicated adaptive radiations of island lizards. *Science* 279:2115-2118.
- Maddison, W.P. (2006). Confounding asymmetries in evolutionary diversification and character change. *Evolution*, 60, 1743-1746.
- Maddison, W.P., Midford, P.E. & Otto, S.P. (2007). Estimating a binary character's effect on speciation and extinction. *Syst. Biol.*, 56, 701-710.
- Marks, C. O., and M. J. Lechowicz. 2006. Alternative designs and the evolution of functional diversity. *American Naturalist* 167:55-66.
- Martin, C. H., and P. C. Wainwright. 2011. Trophic novelty is linked to exceptional rates of morphological diversification in two adaptive radiations of Cyprinodon pupfish. *Evolution* 65:2197-212. doi: 10.1111/j.1558-5646.2011.01294.x.
- O'Meara, B. C., C. M. Ane, M. J. Sanderson, and P. C. Wainwright. 2006. Testing for different rates of continuous trait evolution in different groups using likelihood. *Evolution* 60:922-933.
- Richardson, J. E., R. T. Pennington, T. D. Pennington, and P. M. Hollingsworth. 2001. Rapid diversification of a species-rich genus of Neotropical rain forest trees. *Science* 293:2242-2245.
- Richardson, J., F. Weitz, M. Fay, Q. Cronk, H. Linder, G. Reeves, and M. Chase. 2001. Rapid and recent origin of species richness in the Cape flora of South Africa. *Nature* 412:181-183.
- Sanderson, M. 1998. Reappraising adaptive radiation. *AmerJBot.* 85:1650-1655.

- Sanderson, M., and M. Donoghue. 1994. Shifts in diversification rate with the origin of angiosperms. *Science* 264:1590-1593.
- Sanderson, M. J., and M. J. Donoghue. 1996. Reconstructing shifts in diversification rates on phylogenetic trees. *TREE* 11:15-20.
- Sargent, R.D. (2004) Floral symmetry affect speciation rates in angiosperms. *Proc. Roy. Soc. London Ser. B*, **271**, 603-608.
- Schluter, D. 2000. *The ecology of adaptive radiations*. Oxford University Press, Oxford.
- SILVERTOWN, J., J. FRANCISCO-ORTEGA, and M. CARINE. 2005. The monophyly of island radiations: an evaluation of niche pre-emption and some alternative explanations. *Journal of Ecology* 93:653-657.