

Patterns of floral and chromosome evolution in Onagraceae: a fossil-calibrated supermatrix analysis

William A. Freyman^{a,*}

^a*Jepson Herbarium and Department of Integrative Biology, University of California,
Berkeley*

Abstract

Onagraceae is a cosmopolitan plant family in the order Myrtales comprised of 22 genera and 657 species. I used a GenBank data-mining approach to construct an 11 gene supermatrix of 521 taxa in Onagraceae and Lythraceae. Maximum likelihood and Bayesian inference phylogenetic analyses were performed, and divergence time estimates were calibrated using 5 fossils. Chromosome number evolution was inferred using probabilistic models, and ancestral states of petal color and petal number were reconstructed using maximum likelihood. Correlated evolution between petal color and petal number was tested using Bayesian stochastic character mapping. The monophyly of all major clades in Onagraceae was supported, and Onagraceae was estimated to have diverged from Lythraceae 109 Mya. The base chromosome number of Onagraceae was inferred to be $x = 5$. Petal color and petal number were found to follow a pattern of correlated evolution ($p = 0.00$), suggesting concerted shifts in floral traits may play an important role in shaping diversification of Onagraceae. Further analyses should be performed to test whether shifts in floral traits are correlated with pollinator shifts, and to determine how these shifts affect diversification rates. Furthermore, this work should be extended to test historical biogeographical hypotheses that explain the intercontinental disjunct distribution of Onagraceae.

*Corresponding author

Email address: freyman@berkeley.edu (William A. Freyman)

1. Introduction

Biologists are increasingly using massive phylogenetic datasets to reconstruct ever-growing portions of the Tree of Life in pursuit of large-scale evolutionary questions. A common approach to building large phylogenetic trees has been the supermatrix approach, in which multiple datasets are combined and analyzed simultaneously (de Queiroz and Gatesy, 2007). However, assembling and utilizing supermatrices is not trivial due to difficulties such as determining homology of molecular sequences, defining operational taxonomic units, and managing the amount of missing data (Roure et al., 2007; Yan et al., 2005). `phlawd` and `phylota`... Supermatrix approaches have been successfully used in a number of large scale angiosperm evolutionary studies including, `bling`, `blang`, `blot`. In this study I use a supermatrix to build a phylogeny of the family Onagraceae, and implement a novel method for supermatrix construction in a Python package.

Onagraceae, the evening primrose family, is a well-studied family of flowering plants with 22 described genera and around 660 species (Wagner et al., 2007). The family is found on all continents except Antarctica, and pollen fossils date Onagraceae to the Upper Cretaceous (Grímsson et al., 2012). Onagraceae falls within the order Myrtales, and previous molecular phylogenetic studies find it most closely related to Lythraceae (Conti et al., 1997; Sytsma et al., 2004). Though there have been many genus-level molecular phylogenetic studies in Onagraceae (Berry et al., 2004; Evans et al., 2009; Hoggard and Kores, 2004; Xie et al., 2009; Baum et al., 1994; Wagner and Lammers, 2005), the most recent family-level analyses (Levin et al., 2003, 2004) used only one or two exemplars from some genera. Furthermore, the historical biogeography, divergence times of the major clades, and family-level patterns and processes of diversification remain poorly understood. With over 5000 DNA sequences available in GenBank from previous studies, Onagraceae is an ideal group to apply a data-mining supermatrix approach with the goal of constructing a densely sampled family-level phylogeny.

Here, I use a well-resolved supermatrix phylogeny of Onagraceae to test

whether the taxonomic groups described in Wagner et al. (2007) are monophyletic, and to answer the following questions: 1) When did the major clades diverge? 2) What is the pattern of chromosome number evolution? 3) What are the patterns of petal color and petal number evolution, and are their evolution correlated? Lastly, I discuss ways in which these analyses can be extended to further examine the evolution of Onagraceae, and describe some recently developed metrics that can be used to assess the informativeness of phylogenetic supermatrices.

2. Methods

Supermatrix assembly. Lythraceae was selected as an outgroup since previous molecular phylogenetic analyses place it sister to Onagraceae (Conti et al., 1997; Sytsma et al., 2004). I downloaded all DNA sequences from GenBank release 200 PLN division and performed an exhaustive all-by-all BLASTn (Camacho et al., 2009) comparison of sequences in Onagraceae and Lythraceae. Using a BLASTn e-value of 1.0×10^{-10} threshold and a sequence length percent similarity cutoff of 0.5, I constructed clusters of putative homologs using a single-linkage hierarchical clustering algorithm. Subspecies names were removed from all sequences, and all but one sequence of each species was pruned from each cluster. Clusters that were not phylogenetically informative (< 4 taxa) were discarded, and each cluster was aligned using MUSCLE (Edgar, 2004). The alignments were concatenated by species, and any species that was not present in at least two clusters was removed from the supermatrix. The program written to data-mine GenBank and assemble the supermatrix is available as the open source Python module SUMAC (Freyman, 2014). SUMAC will assemble supermatrices for any taxonomic group recognized in GenBank, and is optimized to run on multicore processors and clusters by utilizing multiple parallel processes.

Phylogenetic analyses. Maximum likelihood (ML) analyses were performed with RAxML-HPG (Stamatakis, 2014) on the CIPRES Scientific Gateway (Miller et al., 2010) using the rapid bootstrap heuristic and the GTRCAT nucleotide

Group	Age (Mya)	Prior Distribution	Mean	SD	Offset	Reference
<i>Circaea</i> (Onagraceae)	12	lognormal	0.0	2.0	12	(Grímsson et al., 2012)
<i>Epilobium</i> (Onagraceae)	12	lognormal	0.0	2.0	12	(Grímsson et al., 2012)
S. Pacific <i>Fuschia</i> (Onagraceae)	23	lognormal	0.0	1.0	23	(Lee et al., 2013)
<i>Ludwigia</i> (Onagraceae)	Paleocene	normal	60.0	3.0	-	(Zhi-Chen et al., 2004)
Lythraceae	82	lognormal	0.0	2.0	82	(Graham, 2013)

Table 1: Fossils used as priors in the Bayesian divergence time analysis.

substitution model. I used the ML tree to select 15 taxa phylogenetically widely distributed in Lythraceae to act as outgroup for the divergence time analysis; all other members of Lythraceae were subsequently removed from the supermatrix. Bayesian estimates of divergence times were inferred using BEAST v1.8 (Drummond and Rambaut, 2007; Suchard and Rambaut, 2009) on CIPRES and calibrated with five fossils identified with morphological synapomorphies (Table 1). The *Ludwigia* fossil pollen was dated broadly to the Paleocene (Grímsson et al., 2012), so I set the prior to a normal distribution with a wide standard deviation to cover the entire time period. For all other calibration points I used a lognormal prior distribution with the offset (the minimum age of the node) corresponding to the fossil age. The BEAST analysis utilized the GTR+Γ nucleotide substitution model with a relaxed molecular clock (uncorrelated lognormal model) and a Yule process tree prior. The Markov Chain Monte Carlo (MCMC) was run for 100 million generations, sampling every 10 thousand generations. Tracer v1.6 (Rambaut et al., 2013) was used to assess the MCMC output for parameter convergence and ensure that the effective sample size for all parameters was above 200. The first 1000 trees were discarded as burn-in, and the remaining 9000 trees were summarized as a maximum clade credibility (MCC) tree with mean divergence times.

Character state reconstruction. I scored six characters, including chromosome number, floral merosity, petal color, and self-compatibility/incompatibility. Character data was assembled from the comprehensive Wagner et al. (2007) Onagraceae monograph. Ancestral chromosome numbers were inferred using maxi-

maximum likelihood and Bayesian methods as implemented in ChromEvol 2.0 (Glick and Mayrose, 2014). Eight different models of chromosome evolution were fit to the Bayesian MCC phylogeny using ChromEvol, and the best fit model was selected using Akaike’s information criterion (Akaike, 1981). Ancestral character state reconstructions of petal number and petal color were performed using Mesquite v2.75 (Maddison and Maddison, 2011) over the Bayesian MCC tree. Characters were treated as unordered categorical data, and optimized using maximum likelihood with the Markov k-state 1 parameter (Mk1) model (Lewis, 2001). Additionally, Bayesian stochastic character mapping (Huelsenbeck et al., 2003) was used to test whether petal color and petal number covaried over the phylogeny. Models of petal color and petal number evolution were configured in the program SIMMAP v1.5 (Bollback, 2006) using unordered states, gamma rate priors, and equal state bias priors. The correlation analysis used SIMMAP’s default predictive sampling configuration and calculated the following correlation statistics: D the overall association between the two characters, and d_{ij} the association between the individual states of each character (Huelsenbeck et al., 2003).

3. Results

Supermatrix assembly. SUMAC evaluated 5571 Onagraceae and 2832 Lythraceae nucleotide sequences to construct the supermatrix. The completed supermatrix consisted of 11 clusters of homologous sequences (Table 2). As used in the maximum likelihood analyses (before pruning the number of outgroup taxa), the supermatrix contained 521 taxa, was 31862 nucleotides long, and contained 93.0% missing data.

Phylogeny and divergence time estimates. The topologies of the ML and Bayesian phylogenies were nearly identical for all major clades within Onagraceae, so only the Bayesian MCC tree (Figures 1 and 2) is shown here. The major exception is that in the ML tree the genus *Chamerion* was recovered as sister to *Epilobium*, whereas in the MCC tree XXXXXX..... In the MCC tree all Onagraceae

DNA Region	# of Taxa	Aligned Length	Missing data (%)	Taxon Coverage Density
ITS	453	1746	13.2	0.87
trnL	234	1429	55.2	0.45
rpl16	91	1414	82.6	0.17
rbcL	77	1474	85.2	0.15
rps16	74	1016	85.8	0.14
rbcL2	64	1310	87.7	0.12
PgiC2	47	4028	91.0	0.09
matK	37	921	92.9	0.07
ndhF	37	2063	92.9	0.07
pgiC	26	14709	95.0	0.05
R5	18	3129	96.6	0.03

Table 2: Clusters of homologous sequences used to assemble the supermatrix.

genera described in Wagner et al. (2007) were recovered as monophyletic clades with posterior probabilities > 0.95 except for sister genera *Neoholmgrenia* and *Camissoniopsis* (posterior = 0.31) (Figure 1). Onagraceae was found to diverge from Lythraceae at 109 Mya (Figure 2). Divergence time estimates of other major clades and 95% highest posterior density (HPD) intervals can be seen in Table 3.

Chromosome number evolution. Out of the eight models of chromosome evolution tested by ChromEvol, the four parameter *CONST_RATE_DEMI_EST* model was the best fit according to the AIC scores. This model has separate rate parameters for chromosome gains (λ), losses (δ), demi-polyploid events (μ), and polyploid events (ρ). Demi-polyploid events occur when reduced and unreduced gametes unite resulting in a $3x$ cytotype. The base chromosome number of Onagraceae was estimated to be $x = 5$, and the ancestral number of the large genus *Epilobium* was estimated to be $n = 9$, though the extant taxa are split in two large monophyletic clades of $n = 10$ and $n = 18$, respectively (Figure 3).

Floral evolution. The maximum likelihood estimate of petal color for the last common ancestor of Onagraceae was pink with a proportional likelihood (*PL*)

Clade	Mean Age (Mya)	95% HPD Min	95% HPD Max
Onagraceae / Lythraceae	109	88	131
<i>Ludwigia</i>	97	76	118
<i>Hauya</i>	49	35	64
<i>Circaea</i> / <i>Fuchsia</i>	37	28	47
<i>Lopezia</i>	71	55	68
<i>Gongylocarpus</i>	60	45	77
<i>Epilobium</i>	49	38	60
<i>Chamerion</i>	47	36	57
<i>Xylonagra</i>	43	33	52
<i>Clarkia</i>	40	32	48
<i>Terapteron</i>	19	10	29
<i>Camissoniopsis</i> / <i>Neoholmgrenia</i>	14	5	23
<i>Eremothera</i> / <i>Camissonia</i>	24	16	33
<i>Taraxia</i>	30	22	38
<i>Chylismiella</i> / <i>Gayophytum</i>	20	10	30
<i>Eulobus</i>	26	19	34
<i>Chylismia</i> / <i>Oenothera</i>	25	18	31

Table 3: Bayesian divergence time estimates of major clades.

	Number of Petals		
	2	4	5
Petal Color			
Pink	-0.005	.011	<i>ns</i>
Yellow	-0.008	.021	<i>ns</i>
White	0.008	0.013	-0.006
Green	<i>ns</i>	-0.011	<i>ns</i>
Red	<i>ns</i>	<i>ns</i>	<i>ns</i>

Table 4: Test statistics for the correlation between petal color evolution and petal number evolution. d_{ij} values are shown for the pairwise comparison of states with $p < 0.01$. *ns* indicates no significant association. Negative d_{ij} values express negative correlation. The overall D value was 0.263 ($p = 0.00$).

of 0.56 (Figure 4). White was the second most likely estimate ($PL = 0.19$). The maximum likelihood estimate of petal number for the last common ancestor of Onagraceae was 4 ($PL = 1.0$), with 8 independent transitions to 0, 2, or 5 petals (Figure 5). Petal color and petal number were found to significantly covary over the phylogeny ($D = 0.263, p = 0.00$). Multiple significant correlations (d_{ij}) were found between individual states of each character and are summarized in Table 4.

4. Discussion

discuss monophyly of tribe Epilobieae, which was supported in maximum likelihood analysis but not supported in Bayesian MCC, could be due to fossil constraints.....

References

Akaike, H., 1981. Likelihood of a model and information criteria. Journal of econometrics 16, 3–14.




Figure 2: Bayesian chronogram of 280 Onagraceae taxa and 15 Lythraceae taxa. Approximate positions of fossil calibration points are shown as black circles. All genera described in Wagner et al. (2007) are colored, and their divergence time estimates and %95 HPD intervals can be seen in Table 3.




Figure 3: Maximum likelihood estimates of ancestral chromosome numbers over the Bayesian MCC phylogeny using the 4 parameter (λ , δ , μ , ρ) model of chromosome evolution selected by Akaike's information criterion. The base number of Onagraceae is inferred to be $x = 5$.




Figure 4: Maximum likelihood ancestral state reconstruction of petal color. Black represents absence of petals. Vertical time scale is in millions of years before present.




Figure 5: Maximum likelihood ancestral state reconstructions of petal number. White = 0, Blue = 2, Green = 4, Orange = 5. Vertical time scale is in millions of years before present.

- Baum, D., Sytsma, K., Hoch, P., 1994. A phylogenetic analysis of *Epilobium* (Onagraceae) based on nuclear ribosomal DNA sequences. *Systematic Botany* 19, 363–388.
- Berry, P.E., Hahn, W.J., Sytsma, K.J., Hall, J.C., Mast, A., 2004. Phylogenetic relationships and biogeography of *Fuchsia* (Onagraceae) based on noncoding nuclear and chloroplast DNA data. *American Journal of Botany* 91, 601–14.
- Bollback, J.P., 2006. Simmap: stochastic character mapping of discrete traits on phylogenies. *BMC Bioinformatics* 7, 88.
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10, 421.
- Conti, E., Litt, A., Wilson, P.G., Graham, S.A., Briggs, B.G., Johnson, L., Sytsma, K.J., 1997. Interfamilial relationships in Myrtales: molecular phylogeny and patterns of morphological evolution. *Systematic Botany* 22, 629–647.
- Drummond, A.J., Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7, 214.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32, 1792–1797.
- Evans, M.E.K., Smith, S.a., Flynn, R.S., Donoghue, M.J., 2009. Climate, niche evolution, and diversification of the "bird-cage" evening primroses (*Oenothera*, sections *Anogra* and *Kleinia*). *The American Naturalist* 173, 225–40.
- Freyman, W.A., 2014. SUMAC: Supermatrix Constructor. <https://github.com/wf8/sumac>.
- Glick, L., Mayrose, I., 2014. Chromevol: Assessing the pattern of chromosome number evolution and the inference of polyploidy along a phylogeny. *Molecular biology and evolution* , msu122.

- Graham, S.A., 2013. Fossil records in the Lythraceae. *The Botanical Review* 79, 48–145.
- Grímsson, F., Zetter, R., Leng, Q., 2012. Diverse fossil Onagraceae pollen from a Miocene palynoflora of north-east China: early steps in resolving the phytogeographic history of the family. *Plant Systematics and Evolution* 298, 671–687.
- Hoggard, G., Kores, P., 2004. The phylogeny of *Gaura* (Onagraceae) based on ITS, ETS, and trnL-F sequence data. *American Journal of Botany* 91, 139–148.
- Huelsenbeck, J.P., Nielsen, R., Bollback, J.P., 2003. Stochastic Mapping of Morphological Characters. *Systematic Biology* 52, 131–158.
- Lee, D.E., Conran, J.G., Bannister, J.M., Kaulfuss, U., Mildenhall, D.C., 2013. A fossil *Fuchsia* (Onagraceae) flower and an anther mass with in situ pollen from the early Miocene of New Zealand. *American Journal of Botany* 100, 2052–2065.
- Levin, R., Wagner, W., Hoch, P., 2004. Paraphyly in tribe Onagreae: insights into phylogenetic relationships of Onagraceae based on nuclear and chloroplast sequence data. *Systematic Botany* 29, 147–164.
- Levin, R.a., Wagner, W.L., Hoch, P.C., Nepokroeff, M., Pires, J.C., Zimmer, E.a., Sytsma, K.J., 2003. Family-level relationships of Onagraceae based on chloroplast rbcL and ndhF data. *American Journal of Botany* 90, 107–15.
- Lewis, P.O., 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. *Systematic Biology* 50, 913–925.
- Maddison, W.P., Maddison, D., 2011. Mesquite: a modular system for evolutionary analysis. Version 2.75. <http://mesquiteproject.org>.
- Miller, M.A., Pfeiffer, W., Schwartz, T., 2010. Creating the CIPRES science gateway for inference of large phylogenetic trees, in: Gateway Computing Environments Workshop (GCE), 2010, IEEE. pp. 1–8.

- de Queiroz, A., Gatesy, J., 2007. The supermatrix approach to systematics. *Trends in Ecology & Evolution* 22, 34–41.
- Rambaut, A., Suchard, M., Drummond, A.J., 2013. Tracer v1.6. <http://tree.bio.ed.ac.uk/software/tracer/>.
- Roure, B., Rodriguez-Ezpeleta, N., Philippe, H., 2007. SCaFoS: a tool for selection, concatenation and fusion of sequences for phylogenomics. *BMC Evolutionary Biology* 7 Suppl 1, S2.
- Stamatakis, A., 2014. RAxML Version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* .
- Suchard, M.A., Rambaut, A., 2009. Many-core algorithms for statistical phylogenetics. *Bioinformatics* 25, 1370–1376.
- Sytsma, K., Litt, A., Zjhra, M., 2004. Clades, clocks, and continents: Historical and biogeographical analysis of Myrtaceae, Vochysiaceae, and relatives in the southern Hemisphere source. *International Journal of Plant Sciences* 165, S85–S105.
- Wagner, W., Lammers, T., 2005. Systematics of *Oenothera* Sections Contortae, Eremia, and Ravenia (Onagraceae). *Systematic Botany* 30, 332–355.
- Wagner, W.L., Hoch, P.C., Raven, P.H., 2007. Revised classification of the Onagraceae. *Systematic Botany Monographs* 83.
- Xie, L., Wagner, W.L., Ree, R.H., Berry, P.E., Wen, J., 2009. Molecular phylogeny, divergence time estimates, and historical biogeography of *Circaea* (Onagraceae) in the Northern Hemisphere. *Molecular Phylogenetics and Evolution* 53, 995–1009.
- Yan, C., Burleigh, J.G., Eulenstein, O., 2005. Identifying optimal incomplete phylogenetic data sets from sequence databases. *Molecular phylogenetics and evolution* 35, 528–35.

Zhi-Chen, S., Wei-Ming, W., Fei, H., 2004. Fossil pollen records of extant angiosperms in China. *The Botanical Review* 70, 425–458.