

Warped Memory Geometry: Feedback Between Coherence Fields and Compactified Curvature

Wayne Fortes

June 2025

Abstract

We extend our six-dimensional emergent spacetime model by introducing a dynamic warp field $\sigma(x)$ that governs the geometry of a compactified 2-sphere. The coherence field $\tau_2(\theta, \phi, t)$, defined on this internal space, is shown to both influence and respond to its curvature. This bidirectional feedback creates a coherent memory mechanism wherein localized emergent phenomena (τ_3) result from resonance between field density and compactification geometry. The resulting model enriches the coupling between hidden coherence structures and observable 4D spacetime, offering testable predictions involving localized curvature shifts and oscillatory gravitational effects.

1. Introduction

The previous formulation of a six-dimensional emergent curvature model demonstrated how a scalar field τ_2 on a compact 2-sphere could influence the effective 4D Einstein equations. However, the geometry of the internal space was held static. In this paper, we upgrade the model by making the internal curvature dynamic. This allows τ_2 to act as both a field and a geometric agent, and unlocks feedback behavior critical to understanding how coherence and emergence propagate in a curved manifold.

2. Geometric Framework

We define the 6D spacetime manifold $\mathcal{M}^6 = M^4 \times S^2$, with coordinates:

- $x^\mu = (t, x, y, z)$ on the observable 4D spacetime
- (θ, ϕ) on the compactified 2-sphere

The metric ansatz is:

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu + e^{2\sigma(x)} (d\theta^2 + \sin^2 \theta d\phi^2)$$

Here $\sigma(x)$ is a scalar warp field on 4D spacetime that determines the effective radius and curvature of the internal sphere.

3. Coherence Field Dynamics

We define a real scalar field $\tau_2(\theta, \phi, t)$ with Lagrangian density:

$$\mathcal{L}_{\tau_2} = -\frac{1}{2} \left[(\partial_t \tau_2)^2 + \frac{1}{e^{2\sigma}} |\nabla_{S^2} \tau_2|^2 + m^2 \tau_2^2 \right]$$

The corresponding field equation is:

$$-\partial_t^2 \tau_2 + \frac{1}{e^{2\sigma}} \Delta_{S^2} \tau_2 = m^2 \tau_2$$

Solutions decompose via spherical harmonics:

$$\tau_2(\theta, \phi, t) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m}(t) Y_{\ell m}(\theta, \phi)$$

4. Curvature Feedback Equation

We define the energy density of τ_2 as:

$$T_{00}(\tau_2) = \frac{1}{2} \left[(\partial_t \tau_2)^2 + \frac{1}{e^{2\sigma}} |\nabla_{S^2} \tau_2|^2 + m^2 \tau_2^2 \right]$$

Integrating over S^2 :

$$E_{\tau_2}(x) = \int_{S^2} d\Omega T_{00}(\tau_2)$$

The warp field $\sigma(x)$ then satisfies:

$$\square_4 \sigma = \alpha E_{\tau_2}(x)$$

where \square_4 is the d'Alembertian in 4D and α is a coupling constant.

5. Modified Einstein Equations

The effective 4D Einstein tensor incorporates contributions from both E_{τ_2} and the kinetic term of σ :

$$G_{\mu\nu} = \kappa \left(T_{\mu\nu}^{(\tau_2)} + \partial_\mu \sigma \partial_\nu \sigma - \frac{1}{2} \eta_{\mu\nu} (\partial\sigma)^2 \right)$$

This establishes σ as both a warp field and a mediator of coherence feedback.

6. Emergence Function τ_3

We redefine $\tau_3(t)$ not as binary, but as a smooth activation:

$$\tau_3(t) = \frac{1}{1 + \exp(-\beta(E_{\tau_2} - E_{\text{crit}}))}$$

This sigmoid transition captures partial coherence, allowing graded emergence and stable attractor behavior.

7. Conclusion and Future Work

We have proposed a bidirectional geometric model where curvature and coherence co-evolve within a six-dimensional framework. The introduction of a warp field σ allows curvature to respond to τ_2 dynamics and close the loop on emergent behavior. Future work includes coupling matter fields to τ_2 , simulating τ_3 cascades, and identifying measurable signatures in gravitational waveforms or localized curvature fluctuations.

References

- [1] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, W. H. Freeman, 1973.
- [2] T. Appelquist, A. Chodos, and P. G. O. Freund, *Modern Kaluza-Klein Theories*, Addison-Wesley, 1987.
- [3] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics, 1982.
- [4] L. Randall and R. Sundrum, “An Alternative to Compactification,” *Physical Review Letters*, vol. 83, no. 23, pp. 4690–4693, 1999.
- [5] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, “The Hierarchy Problem and New Dimensions at a Millimeter,” *Physics Letters B*, vol. 429, pp. 263–272, 1998.