


PHYS1408-2 Exam 2 October 23, 2015

1. A skateboarder rides his skateboard down the wall of a halfpipe from an initial height of 7.0 m. On the bottom of the otherwise frictionless halfpipe he encounters a region 5.0 m long that has a coefficient of friction $\mu=0.1$. How high up the far side of the halfpipe does he rise before he comes to rest?




a) 6.9m b) 6.8m c) 6.7m d) 6.6m e) 6.5m

2. A mass is fired from a height of 3 m above the flat ground below (as shown) at 20 m/s at an angle of 45° above the horizontal. How high above the ground below does the mass fly at the highest point in its trajectory?




a) 3.0m b) 10.2m c) 13.2m d) 20.4m e) 23.4m

3. In order to go all the way around a circular loop of radius $R=10\text{m}$, how fast must a mass be moving as it enters the bottom of the loop?



a) 17 m/s b) 19 m/s c) 21 m/s d) 23 m/s e) 25 m/s


4. A mass of 5 kg is sliding up a ramp ($\mu=0.6$ and $\theta=5^\circ$), initially sliding at 7 m/s. How long does it continue to slide along the ramp before it comes to rest?



a) 0.6s b) 1.0s c) 2.1s d) 3.6s e) 6.7s

5. A 0.05 kg mass is placed on a vertical spring of spring constant $k=2000$ N/m. The spring is compressed 0.02 m from its equilibrium length. When the mass on the compressed spring is released it shoots the mass straight up. How far does the mass fly above the release point?


- a) 0.06 m b) 0.8 m c) 2 m d) 3.5 m e) 14 m



6. A 1 kg mass moving to the right at 5 m/s collides **elastically** in a head on collision with a 2 kg mass moving at 2 m/s to the left. How fast and in what direction is the 2 kg mass moving after the collision?

Positive to the right/ Negative to the left.


- a) +4.33 m/s b) +2.67 m/s c) +1.67 m/s d) -2.67 m/s e) -4.33 m/s



7. A railroad car of mass $2m$ and velocity $2v$ collides completely **inelastically** with another railroad car of mass m and velocity v . The two cars collide and couple together. What fraction of the initial kinetic energy is still retained by the cars after the collision?


8. A bullet (mass 0.005 kg) is fired into a suspended block of wood (10 kg). Initially the wood is not moving. The bullet lodges in the wood. Recoil of the wood causes it to swing (rising 0.01 m). What fraction of the initial kinetic energy remained in the motion of the block of wood? FYI- this is very close to a real situation (i.e., realistic numbers!)

- a) <0.1% b) 0.1-0.5% c) 0.5-1% d) 1-10% e) >10%




9. A mass of 0.5 kg hangs from a pulley ($I = \frac{3}{5}MR^2$, $M=10$ kg, $R=0.3$ m) as shown. What is the downward linear acceleration of the hanging mass?

- a) 0.25 m/s^2 b) 0.50 m/s^2 c) 0.75 m/s^2 d) 1.00 m/s^2 e) 1.25 m/s^2




10. A sphere ($M=2$ kg, $R=0.01$ m, $I = \frac{2}{5}MR^2$) rolls down an incline from a height of 1 m. At the bottom of the incline, what is the sphere's rotational velocity, ω ?



- a) 3.7 rad/s b) 14 rad/s c) 91 rad/s d) 370 rad/s e) 140,000 rad/s

11. A disk $I = \frac{1}{2}MR^2$, mass 200 kg, radius 3 m rotates at 4 rad/s. A wheel ($I = 0.8MR^2$ mass 100 kg, radius 2.5 m initially rotating in the opposite direction at -2 rad/s) is dropped onto the disk. Their collision is completely inelastic. After they stop bouncing what is the angular velocity (rad/s) of the two-object stack?




- a) -0.5 rad/s b) 0 rad/s c) +1 rad/s d) +1.5 rad/s e) +2 rad/s

12. A merry go round (a disk $I = \frac{1}{2}MR^2$, $M=500$ kg, $R=2$ m) rotates at +0.7 rad/s. A child with a mass of 50 kg jumps onto the merry go round (going same direction) at the rim. The child was running at 10 m/s as she jumps on. What is the new rotation rate for the merry go round WITH the child?

- a) 0.8 rad/s b) 1.1 rad/s c) 1.4 rad/s d) 1.7 rad/s e) 2.0 rad/s

13. A mass of 5 kg slides down a 30° inclined surface (with $\mu=0.1$) at 0.80 m/s. It is connected to a pulley (as shown). The pulley is a disk has mass 2 kg, radius 0.05 m, and moment of inertia $0.5 MR^2$. How fast is the mass sliding when it has slid 0.6 m down along the incline?



- a) 2.17 m/s b) 2.04 m/s c) 1.82 m/s d) 0.57 m/s
e) it stops before it slides the whole 0.6 m