

A complete classification of metrizable theta graphs

Guangfu Wang^{a,*}

^a*School of Mathematics and Information Sciences, Yantai University, Yantai, 264005, Shandong Province, China*

Abstract

Cizma and Linial introduced graph metrizability as the problem of deciding whether every consistent system of prescribed paths in a graph can be realized by shortest paths for some positive edge lengths. They asked for a classification of the metrizable theta graphs. We give the complete classification. If $a \leq b \leq c$, then the theta graph $\Theta_{a,b,c}$ is metrizable if and only if $a \leq 2$ or $(a, b, c) = (3, 3, 3)$. The non-metrizable direction follows from the known obstruction $\Theta_{3,3,4}$ and topological-minor closure. The positive direction is constructive. For the family $\Theta_{2,b,c}$, consistency forces certain same-arm and cross-arm choices to be Ferrers relations, and these relations are realized by one-dimensional potentials. The exceptional graph $\Theta_{3,3,3}$ is handled by a two-threshold version of the same construction. The proof is structural and does not rely on enumeration of path systems.

Keywords: Metrizable graph, consistent path system, shortest path, theta graph, Ferrers relation, potential construction

2020 MSC: 05C12, 05C22, 05C75, 05C85

1. Introduction

Shortest paths in weighted graphs are usually studied from a metric viewpoint: one assigns positive lengths to the edges and then analyzes the resulting geodesics. Cizma and Linial proposed the reverse problem [4]. Given a graph G , choose one simple path $P_{u,v}$ between every pair of vertices u, v . The system is *consistent* if every subpath of a chosen path is itself the chosen path between its endpoints. It is *metrizable* if there are positive edge

*Corresponding author.

Email address: gfwang@ytu.edu.cn (Guangfu Wang)

lengths for which every prescribed path is a shortest path. A graph is called metrizable if every consistent path system in it is metrizable.

This definition isolates a graph-theoretic analogue of geodesic geometry. It is close to, but distinct from, Bodwin's topological characterization of partial systems of unique shortest paths [1]: graph metrizability concerns full path systems in a fixed graph and allows ties among shortest paths. Cizma and Linial proved the structural results that make the notion tractable: cycles and outerplanar graphs are strictly metrizable, metrizability is closed downward under topological minors, the class has a finite forbidden-topological-minor characterization, and metrizability is decidable in polynomial time in principle [4]. Subsequent work constructed irreducible non-metrizable systems [5], developed a structure theorem for metrizable graphs [3], showed that the strict variant is minor-closed [2], and studied both the enumerative gap between consistent and metric path systems [7] and quantitative approximation of non-metric systems [6].

One concrete problem left open in [4] is the classification of metrizable theta graphs. The theta graph $\Theta_{a,b,c}$ consists of two branch vertices s, t joined by three internally disjoint s - t paths, called arms, of lengths a, b, c . Since the parameters may be reordered, we assume $a \leq b \leq c$. Cizma and Linial observed that $\Theta_{a,b,c}$ is metrizable when $a = 1$, because such a graph is outerplanar, and they exhibited $\Theta_{3,3,4}$ as a non-metrizable obstruction. Topological-minor closure then makes every $\Theta_{a,b,c}$ with $a \geq 3$, $b \geq 3$, and $c \geq 4$ non-metrizable. Thus the unresolved cases are exactly the infinite family with $a = 2$ and the single graph $\Theta_{3,3,3}$.

Our main theorem resolves these cases.

Theorem 1.1. *Let $a \leq b \leq c$. Then*

$$\Theta_{a,b,c} \text{ is metrizable} \iff a \leq 2 \text{ or } (a, b, c) = (3, 3, 3).$$

Equivalently, $\Theta_{a,b,c}$ is non-metrizable if and only if

$$a \geq 3, \quad b \geq 3, \quad c \geq 4.$$

The theorem gives the promised complete answer for theta graphs. Its significance is twofold. First, it removes the last undetermined theta cases left after the obstruction $\Theta_{3,3,4}$ and topological-minor closure: the infinite family with one arm of length 2 and the exceptional graph $\Theta_{3,3,3}$. Second, the proof gives a structural realization mechanism rather than a computer-aided enumeration of consistent systems. The mechanism is a Ferrers-potential construction: consistency turns the possible detours through the short arm

into nested Ferrers relations, and these relations are then encoded by one-dimensional potentials whose successive differences become the desired edge lengths.

First we reduce to neighborly path systems, since a non-neighborly system on a theta graph avoids some edge and is therefore supported on an outerplanar graph. If the length-two arm of $\Theta_{2,b,c}$ is not the prescribed s – t path, a cycle-cut realization lemma reduces the problem to the metrizability of cycles. The main case is therefore $P_{s,t} = s - a - t$. There, consistency partitions each long arm into a prefix, a middle interval, and a suffix. Same-arm prescribed paths through the short arm form difference Ferrers relations, while cross-arm choices in the middle rectangle form an additive Ferrers relation. We realize these relations by potentials along the two long arms and then define the edge lengths from successive potential differences. The graph $\Theta_{3,3,3}$ is exceptional but positive: the selected length-three arm supplies two nested cuts, and because each remaining middle interval has size at most two, a two-threshold Ferrers representation suffices.

The paper is organized as follows. Section 2 contains definitions and reductions, including the non-metrizable side of the theorem. Section 3 proves the cycle-cut and Ferrers representation lemmas. Section 4 proves metrizability of $\Theta_{2,b,c}$ and $\Theta_{3,3,3}$ and completes the classification.

2. Preliminaries and reductions for theta graphs

All graphs in this paper are finite and connected, and all paths are simple unless explicitly stated otherwise. The length of a path is its number of edges. If Q is a path and x, y are vertices of Q , then $Q[x, y]$ denotes the subpath of Q with endpoints x and y , read in either direction as needed. Since our path systems are indexed by unordered vertex pairs, two opposite orientations of the same path are identified.

If $w : E(G) \rightarrow (0, \infty)$ is an edge weighting and Q is a path, write

$$w(Q) = \sum_{e \in E(Q)} w(e).$$

The corresponding distance is

$$d_w(x, y) = \min\{w(Q) : Q \text{ is an } x\text{--}y \text{ path in } G\}.$$

Thus a path is w -shortest, or a geodesic, if its w -length equals this minimum. A weighting *strictly* induces a path system if each prescribed path is strictly shorter than every other simple path with the same endpoints; ordinary metrizability in this paper allows ties.

We use the following notation for paths on theta graphs. If

$$D : s = d_0, d_1, \dots, d_m = t$$

is an oriented s – t arm and $x, y \in V(D)$, then $x Dy$ denotes the subpath of D from x to y . Concatenations such as $d_i D s a t$ mean the path obtained by first following the subpath $d_i D s$ and then the edges sa and at ; repeated vertices at concatenation points are suppressed. A *prefix* of D means a set of the form $\{d_0, d_1, \dots, d_p\}$, and a *suffix* means a set of the form $\{d_q, d_{q+1}, \dots, d_m\}$.

Definition 2.1. Let $G = (V, E)$ be a graph. A *path system* $\mathcal{P}$ in G is a collection

$$\mathcal{P} = \{P_{u,v} : \{u, v\} \in \binom{V}{2}\},$$

where $P_{u,v}$ is a simple u – v path. The system is *consistent* if, whenever x and y lie on $P_{u,v}$, the subpath $P_{u,v}[x, y]$ is the prescribed path $P_{x,y}$. It is *neighborly* if $P_{u,v} = uv$ for every edge $uv \in E$.

Definition 2.2. A positive edge weighting $w : E \rightarrow (0, \infty)$ *induces* a path system $\mathcal{P}$ if every $P_{u,v}$ is a w -shortest u – v path. A path system is *metrizable* if it is induced by some positive edge weighting. A graph is *metrizable* if every consistent path system in it is metrizable.

For positive integers a, b, c , the theta graph $\Theta_{a,b,c}$ consists of two branch vertices s, t and three internally disjoint s – t paths, called arms, of lengths a, b, c . We always reorder the parameters as $a \leq b \leq c$. In the case $\Theta_{2,b,c}$ the length-two arm is written

$$A = s - a - t,$$

and the two remaining arms are usually written B and C .

We use the following facts from Cizma and Linial [4].

Proposition 2.3. *The following hold.*

- (i) *Cycles are strictly metrizable.*
- (ii) *Outerplanar graphs are strictly metrizable.*
- (iii) *Metrizability is closed downward under topological minors. Equivalently for our purposes, if a graph contains a subdivision of a non-metrizable graph, then it is non-metrizable.*

We also use the cycle structure theorem of Cizma and Linial: after all persistent edges are contracted, every non-trivial consistent path system on a cycle becomes the standard geodesic system on an odd cycle [4, Proposition 4.8]. Here the *standard geodesic system* on an odd cycle means the system that chooses, for every pair of vertices, the shorter of the two cyclic arcs; on an odd cycle this shorter arc is unique. The trivial cycle systems are those supported on a single spanning path of the cycle; they will not arise in the application of Lemma 3.2, because there the restricted cycle system is neighborly.

The next reductions isolate the only theta path systems that require new work.

Proposition 2.4. *Let $G = \Theta_{a,b,c}$. Every non-neighborly consistent path system on G is metrizable.*

Proof. Let $\mathcal{P}$ be non-neighborly. Choose an edge $e = xy$ for which $P_{x,y} \neq xy$. No path of $\mathcal{P}$ can use e : if e appeared in some $P_{u,v}$, then the x - y subpath of $P_{u,v}$ would be the edge xy , and consistency would force $P_{x,y} = xy$.

Thus all paths of $\mathcal{P}$ lie in $G - e$. The graph $G - e$ is outerplanar, so by Proposition 2.3 it has a positive edge weighting w_0 inducing $\mathcal{P}$ as a path system on $G - e$. Extend w_0 to G by assigning e a weight larger than the total w_0 -weight of all edges in $G - e$. Then no path using e can be shorter than a prescribed path, since every prescribed path avoids e . Hence the extended weighting induces $\mathcal{P}$ on G . $\square$

Corollary 2.5. *Any non-metrizable path system on a theta graph is neighborly.*

Proposition 2.6. *Let $a \leq b \leq c$. If $a \geq 3$, $b \geq 3$, and $c \geq 4$, then $\Theta_{a,b,c}$ is non-metrizable.*

Proof. The graph $\Theta_{a,b,c}$ contains a subdivision of $\Theta_{3,3,4}$. The graph $\Theta_{3,3,4}$ is non-metrizable by the certificate displayed as Fig. 22k in [4]. The conclusion follows from topological-minor closure, Proposition 2.3(iii). $\square$

For reference, we record the local algebraic form of the obstruction.

Proposition 2.7. *Suppose a graph contains internally disjoint subpaths*

$$A : s - a_1 - a_2 - t, \quad B : s - b_1 - b_2 - t, \quad C : s - c_1 - c_2 - c_3 - t.$$

If a path system contains the six prescribed paths

$$P_{a_1,b_2} = a_1 s b_1 b_2, \quad P_{a_2,b_1} = a_2 t b_2 b_1,$$

$$\begin{aligned}
P_{a_1, c_3} &= a_1 a_2 t c_3, & P_{a_2, c_2} &= a_2 a_1 s c_1 c_2, \\
P_{b_1, c_3} &= b_1 s c_1 c_2 c_3, & P_{b_2, c_1} &= b_2 t c_3 c_2 c_1,
\end{aligned}$$

then no positive edge weighting can induce all six paths.

Proof. Write

$$\begin{aligned}
A_1 &= w(sa_1), & A_2 &= w(a_1 a_2), & A_3 &= w(a_2 t), \\
B_1 &= w(sb_1), & B_2 &= w(b_1 b_2), & B_3 &= w(b_2 t), \\
C_1 &= w(sc_1), & C_2 &= w(c_1 c_2), & C_3 &= w(c_2 c_3), & C_4 &= w(c_3 t).
\end{aligned}$$

If the six displayed paths are geodesics, then comparison with the natural alternative on the same endpoints gives

$$\begin{aligned}
A_1 + B_1 + B_2 &\leq A_2 + A_3 + B_3, \\
A_3 + B_3 + B_2 &\leq A_2 + A_1 + B_1, \\
A_2 + A_3 + C_4 &\leq A_1 + C_1 + C_2 + C_3, \\
A_2 + A_1 + C_1 + C_2 &\leq A_3 + C_4 + C_3, \\
B_1 + C_1 + C_2 + C_3 &\leq B_2 + B_3 + C_4, \\
B_3 + C_4 + C_3 + C_2 &\leq B_2 + B_1 + C_1.
\end{aligned}$$

Adding these inequalities and cancelling common terms yields $2C_2 \leq 0$, impossible for positive edge weights. $\square$

3. Cycle cuts and Ferrers representations

This section contains the two auxiliary realization tools used in the positive part of the classification. The cycle-cut lemma handles the case in which the length-two arm is not the prescribed s - t path. The Ferrers lemmas handle the remaining case $P_{s,t} = s - a - t$ by converting monotone choices of paths into threshold inequalities.

Lemma 3.1. *Let H be a cycle with distinct vertices s, t , let $\mathcal{Q}$ be a non-trivial consistent path system on H , and let $S \subseteq V(H)$ satisfy the following compatibility conditions:*

- (a) $s \in S$ and $t \notin S$;
- (b) on each of the two s - t arcs of H , the set S is a prefix starting at s ;
- (c) if $v \notin S$, then the prescribed t - v path in $\mathcal{Q}$ contains no vertex of S .

Then, for the purpose of constructing a weighting that both induces $\mathcal{Q}$ and strictly separates S from its complement by a threshold in the values

$$d(s, v) - d(t, v),$$

the Cizma–Linial contraction of persistent components may be performed in a cut-respecting way. More precisely, because S is a prefix on each of the two s – t arcs, there are at most two cycle edges with one endpoint in S and the other in $V(H) \setminus S$. Contract persistent components as usual except at these boundary edges: whenever a persistent component contains such a boundary edge uv , contract the maximal persistent subpath on the S side of uv and the maximal persistent subpath on the complementary side separately, and retain the edge uv between the two contracted vertices. Components not meeting a boundary edge are contracted in the ordinary way. The resulting quotient is a cycle which, after suppressing at most two retained boundary subdivision edges, is the standard odd-cycle quotient of Cizma–Linial. Its path system is the corresponding subdivision of the standard odd-cycle system.

Suppose this cut-respecting quotient has a positive edge weighting $w^\#$ which strictly induces its quotient path system and, for some real number h , satisfies

$$\begin{aligned} d_{w^\#}(s, v) - d_{w^\#}(t, v) &< h & (v \in S^\#), \\ d_{w^\#}(s, v) - d_{w^\#}(t, v) &> h & (v \notin S^\#), \end{aligned}$$

where $S^\#$ is the image of S in the cut-respecting quotient. Then $w^\#$ expands to a positive edge weighting w of the original cycle H which induces $\mathcal{Q}$ and satisfies the same two strict threshold inequalities for all original vertices.

Proof. The first assertions are consequences of the prefix assumptions. The set S has at most one boundary edge on each of the two s – t arcs, hence at most two boundary edges in the whole cycle. Splitting a persistent component only at these boundary edges leaves every contracted piece on a single side of the cut and keeps each boundary edge as an explicit edge of the quotient. Suppressing the retained boundary edges recovers exactly the usual persistent-edge quotient of the cycle system; before suppression, the quotient is just a subdivision of that odd-cycle quotient at at most two places.

It remains to justify the expansion step. The ordinary persistent-edge expansion lemma of Cizma and Linial [4, Lemma 4.6] says that contracted persistent components may be replaced by sufficiently small positive edge weights without changing the induced cycle system. We need only record that the same small expansion can be chosen to preserve the cut inequalities.

Indeed, on the cut-respecting quotient all relevant inequalities are strict and finite in number. They are of two types:

- (i) for every pair of quotient vertices x, y and every non-prescribed simple x - y path R ,

$$w^\sharp(Q_{x,y}^\sharp) < w^\sharp(R);$$

- (ii) for every quotient vertex v ,

$$d_{w^\sharp}(s, v) - d_{w^\sharp}(t, v) < h \quad \text{if } v \in S^\sharp,$$

$$d_{w^\sharp}(s, v) - d_{w^\sharp}(t, v) > h \quad \text{if } v \notin S^\sharp.$$

Let $\gamma > 0$ be smaller than the minimum slack in these inequalities. Expand every contracted component by assigning positive edge weights whose total weight is less than $\gamma/(10|V(H)|)$. All path lengths and all quantities $d(s, v) - d(t, v)$ vary continuously with these inserted weights; hence the strict inequalities involving quotient vertices persist.

For vertices internal to an expanded component contained wholly in one side of the cut, the value of $d(s, v) - d(t, v)$ differs from the value at the contracted quotient vertex by less than $\gamma/5$, so the vertex remains on the same side of the threshold. For a crossing persistent component, the boundary edge was retained in the quotient and already separates an S -side endpoint from a complementary endpoint. Only the two one-sided pieces adjacent to that boundary edge are expanded, and the same smallness estimate keeps all internal vertices of those pieces on their prescribed sides. Thus both the geodesic inequalities and the cut inequalities survive the expansion. The resulting positive weighting of H induces $\mathcal{Q}$ and has the required strict threshold separation. $\square$

Lemma 3.2. *Let H be a cycle with two distinct vertices s, t , and let $\mathcal{Q}$ be a non-trivial consistent path system on H . Suppose that a new vertex a is joined to s and t , and that a consistent extension is neighborly on the new edges, so $P_{a,s} = as$ and $P_{a,t} = at$. Assume further that each a - v path, $v \in V(H)$, begins either with as or with at . Then there is a positive edge weighting w_H inducing $\mathcal{Q}$ and a real number h such that*

$$P_{a,v} \text{ begins with } as \implies d_{w_H}(s, v) - d_{w_H}(t, v) < h,$$

$$P_{a,v} \text{ begins with } at \implies d_{w_H}(s, v) - d_{w_H}(t, v) > h.$$

Proof. Let

$$S = \{v \in V(H) : P_{a,v} \text{ begins with } as\}.$$

Consistency of the extension gives three elementary consequences. First, $s \in S$ and $t \notin S$. Second, on each of the two s – t arcs of H , the set S is a prefix starting at s : if $v \in S$ and u lies on the prescribed s – v subpath of $P_{a,v}$, then the a – u subpath of $P_{a,v}$ begins with as . Third, if $v \notin S$, then the prescribed t – v subpath of $P_{a,v}$ contains no vertex of S , for otherwise the a – u subpath at the first such vertex $u \in S$ would begin in two different ways.

By Lemma 3.1, it is enough to construct the desired strict threshold separation after the cut-respecting cycle reduction. After suppressing at most two retained boundary subdivisions, the quotient is the standard geodesic system on an odd cycle. Such retained subdivision edges can be assigned arbitrarily small positive weights after the strict odd-cycle inequalities below have been obtained, so it suffices to treat the reduced odd cycle itself.

Label the reduced odd cycle by $0, 1, \dots, 2n$ in cyclic order, put $s = 0$, and write $t = d$ with $1 \leq d \leq n$. By the prefix property just proved, there are integers q and r with

$$0 \leq q < d, \quad n - d \leq r \leq n,$$

such that, writing $N = 2n + 1$,

$$S = \{N - r, \dots, N - 1, 0, 1, \dots, q\}.$$

The lower bound on r is exactly the assertion that the standard shortest path from t to every vertex outside S avoids S . Put

$$p = N - r - 1,$$

with indices read modulo N when necessary. Thus the two boundary pairs of the cut are $(q, q + 1)$ and $(p, p + 1)$.

Let e_i be the edge joining i to $i + 1$ modulo N . For $k \in \mathbb{Z}/N\mathbb{Z}$, let R_k be the non-negative edge weighting that assigns weight 1 to e_k and e_{k+n} and weight 0 to every other edge. Each R_k induces the standard odd-cycle system non-strictly: every standard arc contains at most one of the two weighted edges, whereas the complementary arc contains at least one. Set

$$w_0 = R_q + R_p + \frac{1}{2}R_0$$

and write

$$\phi(v) = d_{w_0}(0, v) - d_{w_0}(d, v).$$

For the two boundary gaps define

$$F_1 = \phi(p) - \phi(q), \quad F_2 = \phi(q+1) - \phi(p+1).$$

A direct check on the odd cycle gives

$$F_1(R_q + R_p) = 2, \quad F_2(R_q + R_p) \in \{0, 2, 4\},$$

and

$$F_1(R_0) \in \{-2, 0, 2\}, \quad F_2(R_0) = 2.$$

Consequently $F_1(w_0) > 0$ and $F_2(w_0) > 0$, that is,

$$\phi(q) < \phi(p), \quad \phi(p+1) < \phi(q+1).$$

For every non-negative weighting that induces the standard odd-cycle paths, the function ϕ is monotone on each of the two s - t arcs: as one moves away from s along an arc, the distance from s weakly increases and the distance from t weakly decreases. The two boundary inequalities therefore imply

$$\max_{v \in S} \phi(v) < \min_{v \notin S} \phi(v).$$

Choose h strictly between these two values.

Finally add a sufficiently small positive multiple of the uniform edge weighting. Since in an odd cycle the standard arc between two vertices has fewer edges than the complementary arc, the standard path system becomes strictly induced; the strict cut separation remains true. If the cut-respecting quotient contains retained boundary subdivisions, give those subdivision edges sufficiently small positive weights. Lemma 3.1 then expands the remaining persistent components and gives a positive weighting of the original cycle that induces $\mathcal{Q}$ and satisfies the two desired strict threshold inequalities. $\square$

Proposition 3.3. *Let $G = \Theta_{2,b,c}$, and let its length-two arm be $A = s - a - t$. If $\mathcal{P}$ is neighborly and $P_{s,t} \neq A$, then $\mathcal{P}$ is metrizable.*

Proof. Let $H = G - a$. Then H is the cycle formed by the two long arms. No prescribed path whose endpoints lie in H can pass through a , because such a path would contain A as its s - t subpath, forcing $P_{s,t} = A$ by consistency. Therefore $\mathcal{P}$ restricts to a cycle path system $\mathcal{Q}$ on H .

Because $\mathcal{P}$ is neighborly, the restricted cycle system $\mathcal{Q}$ is neighborly and hence non-trivial. Apply Lemma 3.2 to obtain w_H and h . Choose $M > |h|$

so large that no shortest path between two vertices of H can improve by going through a , and set

$$w(as) = M, \quad w(at) = M + h.$$

Both weights are positive. For $v \in V(H)$, the route from a through s is no longer than the route through t exactly when

$$d_{w_H}(s, v) - d_{w_H}(t, v) \leq h.$$

Thus every prescribed a - v path is a geodesic, and the choice of M preserves all prescribed paths inside H . $\square$

We now spell out the Ferrers terminology used in the proof of $\Theta_{2,b,c}$. This is the threshold viewpoint on Ferrers or chain graphs, also known in an equivalent weighted form as difference graphs; see, for example, [8, 9].

Definition 3.4. A *chain* is a finite totally ordered set. For the standard chain $I = \{1, \dots, m\}$, a prefix is an initial interval $\{1, \dots, p\}$, possibly empty or all of I ; a suffix is a final interval $\{q, \dots, m\}$, possibly empty or all of I .

Let I and J be chains. A relation $F \subseteq I \times J$ is an *additive Ferrers relation* if it is downward closed in both coordinates: whenever $(i, j) \in F$, $i' \leq i$, and $j' \leq j$, then $(i', j') \in F$. Equivalently, the 0–1 matrix of F has left-justified rows and the row lengths are weakly decreasing as i increases. An *additive threshold representation* of F is a choice of real labels x_i for $i \in I$ and y_j for $j \in J$ such that membership in F is determined by the sign of $x_i + y_j$.

For a single ordered arm $D : s = d_0, d_1, \dots, d_m = t$, a *potential* is simply a real label X_i assigned to d_i . In the construction below the potentials are strictly increasing along D , and the edge length of $d_{i-1}d_i$ is set to $(X_i - X_{i-1})/2$. This choice is useful because the total length of D is then $(X_m - X_0)/2$, and, writing d_D for the path-length distance restricted to the arm D ,

$$d_D(s, d_i) - d_D(t, d_i) = X_i$$

when $X_0 = -X_m$. A relation $F \subseteq D^- \times D^+$, where D^- is a prefix and D^+ is a suffix of D , is a *difference Ferrers relation* if it is downward closed in the prefix coordinate and upward closed in the suffix coordinate:

$$(d_i, d_j) \in F, \quad i' \leq i, \quad j' \geq j \quad \implies \quad (d_{i'}, d_{j'}) \in F.$$

After reversing the order on D^+ , this is exactly an additive Ferrers relation. A *difference threshold representation* realizes membership in F by an inequality

of the form $X_j - X_i > \text{constant}$. A *prefix cut represented by a threshold* means that a prescribed prefix is exactly the set of chain elements whose potential is below that threshold.

Lemma 3.5. *Let $I = \{1, \dots, m\}$ and $J = \{1, \dots, n\}$ be chains. Let $F \subseteq I \times J$ satisfy*

$$(i, j) \in F, \quad i' \leq i, \quad j' \leq j \quad \implies \quad (i', j') \in F.$$

Then there are strictly increasing real numbers x_i and y_j such that

$$(i, j) \in F \quad \iff \quad x_i + y_j < 0.$$

Moreover, if a prefix cut is prescribed in I and a prefix cut is prescribed in J , the representation may be chosen so that both cuts are represented by a common threshold h :

$$i \text{ lies in the prescribed prefix of } I \iff x_i < h,$$

$$j \text{ lies in the prescribed prefix of } J \iff y_j < h.$$

Proof. For each row i , let r_i be the largest column index j such that $(i, j) \in F$, and set $r_i = 0$ if the i th row is empty. The Ferrers condition says exactly that the row lengths are nested:

$$r_1 \geq r_2 \geq \dots \geq r_m.$$

Put $x_i = i$. For $j \in J$, let

$$k_j = \max\{i : r_i \geq j\},$$

with $k_j = 0$ if the set is empty. Since the row lengths are non-increasing, the sequence k_j is non-increasing in j . Choose $0 < \delta < 1/(2n)$ and set

$$y_j = -k_j - \frac{1}{2} + \delta j.$$

Then $y_1 < y_2 < \dots < y_n$. Moreover,

$$x_i + y_j = i - k_j - \frac{1}{2} + \delta j.$$

If $i \leq k_j$, then $x_i + y_j < 0$; if $i > k_j$, then $x_i + y_j > 0$. Hence $x_i + y_j < 0$ holds exactly when $i \leq k_j$, equivalently when $j \leq r_i$, which is precisely $(i, j) \in F$.

Now suppose the prescribed prefixes are $I_0 = \{1, \dots, p\}$ and $J_0 = \{1, \dots, q\}$, allowing $p = 0, m$ and $q = 0, n$. Starting with the representation just constructed, replace

$$x_i \mapsto \lambda x_i + \alpha, \quad y_j \mapsto \lambda y_j - \alpha,$$

where $\lambda > 0$. This multiplies every sum $x_i + y_j$ by λ , so it preserves all signs and therefore preserves F . A common threshold h must lie in both intervals

$$(\lambda x_p + \alpha, \lambda x_{p+1} + \alpha), \quad (\lambda y_q - \alpha, \lambda y_{q+1} - \alpha),$$

where $x_0 = y_0 = -\infty$ and $x_{m+1} = y_{n+1} = +\infty$. In the non-extreme case, these intervals overlap whenever

$$\frac{\lambda(y_q - x_{p+1})}{2} < \alpha < \frac{\lambda(y_{q+1} - x_p)}{2},$$

and the interval for α is non-empty because $x_p < x_{p+1}$ and $y_q < y_{q+1}$. Empty and full prefixes are handled by taking h outside the corresponding finite range of labels. Choosing h in the overlap represents both prescribed prefix cuts. $\square$

Lemma 3.6. *Let $D : s = d_0, d_1, \dots, d_m = t$ be a chain. Let D^- be a prefix containing s , let D^+ be a suffix containing t , assume $D^- \cap D^+ = \emptyset$, and let $D^0 = D \setminus (D^- \cup D^+)$. Suppose $F \subseteq D^- \times D^+$ satisfies*

$$(d_i, d_j) \in F, \quad i' \leq i, \quad j' \geq j \implies (d_{i'}, d_{j'}) \in F,$$

and assume that the row s and the column t belong entirely to F . Fix $L > 1$. Given any strictly increasing prescribed potentials in $(-1, 1)$ on D^0 , there are $M > L$ and a strictly increasing sequence

$$-M = X_0 < X_1 < \dots < X_m = M$$

which extends those middle potentials, has $X_i < -L$ on $D^- \setminus \{s\}$ and $X_i > L$ on $D^+ \setminus \{t\}$, and satisfies

$$(d_i, d_j) \in F \iff X_j - X_i > M + L$$

for all $d_i \in D^-$ and $d_j \in D^+$ with $i < j$.

Proof. Reverse the order on D^+ : in the reversed order, d_j comes before $d_{j'}$ exactly when $j \geq j'$. With this reversed order on the second coordinate, the displayed hypothesis is the additive Ferrers condition. Applying Lemma 3.5,

and then multiplying all potentials and the threshold by a positive constant if necessary, gives real numbers U_i on D^- , real numbers V_j on D^+ , and a threshold $R > 0$ such that

$$(d_i, d_j) \in F \iff U_i + V_j < R.$$

The U_i are strictly increasing in the natural order on D^- , while the V_j are strictly increasing in the reversed order on D^+ . Translate both families and the threshold by replacing U_i with $U_i - U_s$, V_j with $V_j - V_t$, and R with $R - U_s - V_t$. This preserves the displayed equivalence and gives $U_s = 0$ and $V_t = 0$. Since the row s and the column t belong entirely to F , we have $V_j < R$ for every $d_j \in D^+$ and $U_i < R$ for every $d_i \in D^-$.

Set $M = R + L$. For the prefix and suffix vertices define

$$X_i = -M + U_i \quad (d_i \in D^-), \quad X_j = M - V_j \quad (d_j \in D^+).$$

Then $X_s = -M$ and $X_t = M$. The values on D^- are strictly increasing because the U_i are strictly increasing. The values on D^+ are also strictly increasing in the natural order along D : as j increases, the reversed-order potential V_j strictly decreases, and hence $M - V_j$ strictly increases. Because $U_i < R$ and $V_j < R$, all non-end prefix values are below $-L$ and all non-end suffix values are above L .

Insert the prescribed middle potentials on D^0 . These lie in $(-1, 1)$, while the prefix values lie below $-L$ and the suffix values lie above L with $L > 1$; therefore the full sequence $X_0 < X_1 < \dots < X_m$ is strictly increasing. Finally,

$$X_j - X_i = 2M - (U_i + V_j),$$

and, since $M = R + L$,

$$X_j - X_i > M + L \iff U_i + V_j < R.$$

Thus the sequence has all required properties. $\square$

Remark 3.7. The disjointness hypothesis in Lemma 3.6 is essential for the stated range conditions. It is automatically satisfied in the two applications below: there, an internal vertex belonging to both D^- and D^+ would force two different prescribed s - d_i subpaths, contradicting consistency.

Lemma 3.8. *Let I and J be possibly empty finite chains, each of size at most 2, and let $F \subseteq I \times J$ be an additive Ferrers relation. Let*

$$I_b \subseteq I_a \subseteq I, \quad J_b \subseteq J_a \subseteq J$$

be prescribed prefixes. Then there are strictly increasing real labels x_i for $i \in I$, strictly increasing real labels y_j for $j \in J$, and two real thresholds $h_b < h_a$ such that

$$(i, j) \in F \implies x_i + y_j \leq 0, \quad (i, j) \notin F \implies x_i + y_j \geq 0,$$

$$i \in I_b \implies x_i \leq h_b, \quad i \notin I_b \implies x_i \geq h_b,$$

$$i \in I_a \implies x_i \leq h_a, \quad i \notin I_a \implies x_i \geq h_a,$$

and the same two threshold conditions hold for $J_b \subseteq J_a$ with the labels y_j . The labels and thresholds may also be made arbitrarily small by a common positive rescaling.

Proof. Assign to each element of I a type 0, 1, or 2: type 0 means membership in I_b , type 1 means membership in $I_a \setminus I_b$, and type 2 means membership in $I \setminus I_a$. Define the types in J similarly. Since all four sets are prefixes, the type sequence along each chain is non-decreasing.

We first choose auxiliary labels ξ_i, η_j and a real number T with the following properties:

$$(i, j) \in F \implies \xi_i + \eta_j \leq T, \quad (i, j) \notin F \implies \xi_i + \eta_j \geq T,$$

and each auxiliary label lies in the interval prescribed by its type,

$$\text{type } 0 : (-\infty, -1], \quad \text{type } 1 : [-1, 1], \quad \text{type } 2 : [1, \infty).$$

The auxiliary labels are required to be strictly increasing along their chains.

If one of the chains is empty, there is nothing to separate. If one chain has a single element, say $I = \{1\}$, then a Ferrers relation in $I \times J$ is simply an initial segment of the chain J . Choose ξ_1 in its prescribed type interval and choose the labels η_j strictly increasing inside their prescribed type intervals; if necessary, move the labels in the unbounded type-0 or type-2 intervals farther apart. The sums $\xi_1 + \eta_j$ are then strictly increasing in j , so a threshold T separates the initial segment. The case $|J| = 1$ is symmetric. Suppose therefore that $I = \{1, 2\}$ and $J = \{1, 2\}$. Put

$$s_{ij} = \xi_i + \eta_j.$$

Since $\xi_1 < \xi_2$ and $\eta_1 < \eta_2$, we always have

$$s_{11} < s_{12} < s_{22}, \quad s_{11} < s_{21} < s_{22}.$$

Thus every Ferrers lower ideal in the 2×2 grid is automatically separable by a threshold except possibly the two ideals of size two. If

$$F = \{(1, 1), (1, 2)\},$$

then separation is equivalent to $s_{12} \leq s_{21}$, that is,

$$\eta_2 - \eta_1 \leq \xi_2 - \xi_1.$$

If

$$F = \{(1, 1), (2, 1)\},$$

then separation is equivalent to $s_{21} \leq s_{12}$, that is,

$$\xi_2 - \xi_1 \leq \eta_2 - \eta_1.$$

It remains only to know that the two gaps can be chosen with either prescribed weak order while respecting the type intervals. For a non-decreasing two-term type sequence, the possible positive gaps are as follows: the type sequence $(1, 1)$ allows exactly the gaps $0 < g \leq 2$; the type sequence $(0, 2)$ allows exactly the gaps $g \geq 2$; and the four remaining type sequences allow every positive gap. Hence, to impose $\eta_2 - \eta_1 \leq \xi_2 - \xi_1$, choose the ξ -gap at least as large as the η -gap; in the only extremal case $(\xi\text{-types}) = (1, 1)$ and $(\eta\text{-types}) = (0, 2)$, take both gaps equal to 2. The opposite inequality is handled symmetrically. This proves the existence of the auxiliary labels and the separating threshold T .

Now set

$$H = -\frac{T}{2}, \quad x_i = H + \xi_i, \quad y_j = H + \eta_j,$$

and choose

$$h_b = H - 1, \quad h_a = H + 1.$$

Then $h_b < h_a$, and the type intervals above translate exactly into the two required prefix-threshold conditions. Moreover

$$x_i + y_j \leq 0 \iff \xi_i + \eta_j \leq T,$$

and similarly for the reversed weak inequality. A common positive rescaling preserves all weak inequalities, all represented cuts, and the strict inequality $h_b < h_a$. The lemma follows. $\square$

4. Metrizable cases and proof of the classification

We now prove the two positive cases not already covered by outerplanarity, and then complete the proof of Theorem 1.1. The first case is the infinite family $\Theta_{2,b,c}$; the second is the exceptional graph $\Theta_{3,3,3}$.

Theorem 4.1. *For every $2 \leq b \leq c$, the theta graph $\Theta_{2,b,c}$ is metrizable.*

Proof. Let the length-two arm be

$$A = s - a - t,$$

and write the other arms as

$$B : s = b_0, b_1, \dots, b_b = t, \quad C : s = c_0, c_1, \dots, c_c = t.$$

Let $\mathcal{P}$ be a consistent path system. By Proposition 2.4, we may assume $\mathcal{P}$ is neighborly. By Proposition 3.3, we may assume

$$P_{s,t} = A = s - a - t.$$

For a long arm $D : s = d_0, d_1, \dots, d_m = t$, define three vertex sets

$$D^- = \{s\} \cup \{d_i : 1 \leq i \leq m-1, P_{d_i,t} = d_i D s a t\},$$

$$D^+ = \{t\} \cup \{d_i : 1 \leq i \leq m-1, P_{s,d_i} = s a t D d_i\},$$

and

$$D^0 = D \setminus (D^- \cup D^+).$$

Thus D^- consists of vertices whose prescribed path to t first goes backward along D to s and then crosses the length-two arm A ; D^+ consists of vertices whose prescribed path from s first crosses A and then follows D backward from t ; and D^0 is the middle part where neither of these two forced behaviours occurs.

Consistency implies that D^- is a prefix, D^+ is a suffix, and $D^- \cap D^+ = \emptyset$. For example, if $d_i \in D^-$ and $i' \leq i$, then the $d_{i'}-t$ subpath of $P_{d_i,t}$ is $d_{i'} D s a t$, so $d_{i'} \in D^-$. The assertion for D^+ is symmetric. If an internal vertex d_i belonged to both D^- and D^+ , then the $s-d_i$ subpath of $P_{d_i,t}$ would be the direct subpath $s D d_i$, whereas P_{s,d_i} would be $s a t D d_i$, contradicting consistency.

For $i < j$ on the same long arm D , if P_{d_i,d_j} uses A , then $d_i \in D^-$ and $d_j \in D^+$. The set of such same-arm pairs is a difference Ferrers relation

on $D^- \times D^+$: if (d_i, d_j) uses A , $i' \leq i$, and $j' \geq j$, then P_{d_i, d_j} contains $d_{i'} D s a t D d_{j'}$ as a subpath, so $P_{d_{i'}, d_{j'}}$ also uses A .

Next consider a cross-arm pair $b_i \in V(B) \setminus \{s, t\}$ and $c_j \in V(C) \setminus \{s, t\}$. There are four relevant simple routes between them:

$$S : b_i B s C c_j, \quad T : b_i B t C c_j, \quad U : b_i B s a t C c_j, \quad V : b_i B t a s C c_j.$$

The letters record which junctions are used: S goes through the branch vertex s , T goes through t , U crosses A from s to t , and V crosses A from t to s . Consistency gives

$$P_{b_i, c_j} = U \iff b_i \in B^- \text{ and } c_j \in C^+,$$

$$P_{b_i, c_j} = V \iff b_i \in B^+ \text{ and } c_j \in C^-.$$

Indeed, the forward implication follows by taking the b_i - t and s - c_j subpaths. Conversely, if $b_i \in B^-$ and $c_j \in C^+$, then the routes S, T, V would give a wrong b_i - t or s - c_j subpath, so only U is possible. The proof for V is symmetric.

Outside the two exceptional rectangles $B^- \times C^+$ and $B^+ \times C^-$, the route is either S or T . More precisely, away from these rectangles, membership in B^- or C^- forces the route through s , while membership in B^+ or C^+ forces the route through t ; otherwise one obtains a wrong b_i - t , s - c_j , s - b_i , or c_j - t subpath. On the central rectangle $B^0 \times C^0$, the S -region is an additive Ferrers relation: if $P_{b_i, c_j} = S$, $i' \leq i$, and $j' \leq j$, then $P_{b_{i'}, c_{j'}}$ contains $b_{i'} B s C c_{j'}$ as a subpath, hence $P_{b_{i'}, c_{j'}} = S$.

Consistency also controls the paths from a to the long arms. Vertices in D^- have their a - d_i path beginning with as , and vertices in D^+ have their a - d_i path beginning with at . The opposite choice would either give the direct d_i - t or s - d_i subpath where the definition of D^- or D^+ prescribes a path through A , or would contain an s - t subpath different from A .

Fix $L = 2$. The vertices d_i of any long arm whose a - d_i path begins with as form a prefix: if $i' \leq i$ and P_{a, d_i} begins with as , then the a - $d_{i'}$ subpath of P_{a, d_i} also begins with as . If $B^0 \times C^0$ is empty, choose arbitrary strictly increasing middle potentials in $(-1, 1)$ on the non-empty middle chain or chains, and choose a real threshold h representing the relevant prefix cuts there; if both middle chains are empty, take $h = 0$. Since a finite ordered set with a prescribed prefix is separated by an open interval of possible thresholds, after a common small rescaling we may also require $|h| < 1$. If both B^0 and C^0 are non-empty, apply Lemma 3.5 to the Ferrers S -region on $B^0 \times C^0$, together with the prefix cuts on B^0 and C^0 determined by whether

P_{a,b_i} and P_{a,c_j} begin with as . This gives strictly increasing middle potentials X_i for $b_i \in B^0$, Y_j for $c_j \in C^0$, and a threshold h , such that

$$P_{a,b_i} \text{ begins with } as \iff X_i < h, \quad P_{a,c_j} \text{ begins with } as \iff Y_j < h,$$

and

$$P_{b_i,c_j} = S \iff X_i + Y_j < 0 \quad (b_i \in B^0, c_j \in C^0).$$

Scaling all middle potentials and, when necessary, the threshold by a common small positive factor, assume $|X_i|, |Y_j|, |h| < 1$ whenever the corresponding quantities are present.

Now apply Lemma 3.6 separately to the same-arm Ferrers relations on $B^- \times B^+$ and $C^- \times C^+$, preserving the already chosen middle potentials. We obtain strictly increasing potential sequences

$$-M_B = X_0 < X_1 < \dots < X_b = M_B,$$

$$-M_C = Y_0 < Y_1 < \dots < Y_c = M_C,$$

with prefix values below $-L$, middle values in $(-1, 1)$, suffix values above L , and with same-arm paths through A represented exactly by

$$X_j - X_i > M_B + L \quad \text{on } B^- \times B^+,$$

$$Y_j - Y_i > M_C + L \quad \text{on } C^- \times C^+.$$

Define edge weights by

$$w(sa) = \frac{L-h}{2}, \quad w(at) = \frac{L+h}{2},$$

$$w(b_{i-1}b_i) = \frac{X_i - X_{i-1}}{2}, \quad w(c_{j-1}c_j) = \frac{Y_j - Y_{j-1}}{2}.$$

All weights are positive because $|h| < L$ and the potential sequences are strictly increasing. The total lengths of B and C are M_B and M_C , respectively, and the potential normalization gives

$$d_B(s, b_i) - d_B(t, b_i) = X_i, \quad d_C(s, c_j) - d_C(t, c_j) = Y_j.$$

We verify that every prescribed path is a geodesic. First consider a and a vertex d_i on a long arm D , and let E be the other long arm. Write Z_i for the corresponding potential and M_D, M_E for the total lengths of D, E . The two direct routes through the endpoints of A have lengths

$$R_s = \frac{L-h}{2} + \frac{M_D + Z_i}{2}, \quad R_t = \frac{L+h}{2} + \frac{M_D - Z_i}{2}.$$

Thus $R_s \leq R_t$ exactly when $Z_i \leq h$. There are also two routes that use the other long arm E : their lengths are

$$R_s + M_E - Z_i, \quad R_t + M_E + Z_i.$$

If the prescribed path begins with as , then either $d_i \in D^-$, so $Z_i < -L$, or $d_i \in D^0$ and the threshold construction gives $Z_i < h < 1$. Hence $R_s \leq R_t$ and

$$\begin{aligned} (R_s + M_E - Z_i) - R_s &= M_E - Z_i > 0, \\ (R_t + M_E + Z_i) - R_s &= M_E + h > 0, \end{aligned}$$

because $M_E > L = 2$ and $|h| < 1$. If the prescribed path begins with at , then either $d_i \in D^+$, so $Z_i > L$, or $d_i \in D^0$ and $Z_i \geq h > -1$. Hence $R_t \leq R_s$ and

$$\begin{aligned} (R_t + M_E + Z_i) - R_t &= M_E + Z_i > 0, \\ (R_s + M_E - Z_i) - R_t &= M_E - h > 0. \end{aligned}$$

Therefore every prescribed $a-d_i$ path is a geodesic.

For two vertices d_i, d_j on the same long arm D , $i < j$, let Z denote the corresponding potential sequence and let M_D be the total length of the arm. The direct route along D has length $(Z_j - Z_i)/2$, while the route through A has length

$$M_D + L - \frac{Z_j - Z_i}{2}.$$

Thus the route through A is no longer than the direct route exactly when $Z_j - Z_i \geq M_D + L$. If the prescribed path uses A , then $(d_i, d_j) \in D^- \times D^+$ and Lemma 3.6 gives $Z_j - Z_i > M_D + L$. If the prescribed path is direct and $(d_i, d_j) \in D^- \times D^+$, the same lemma gives $Z_j - Z_i \leq M_D + L$. In all remaining direct cases this inequality is automatic from the potential ranges: two prefix vertices, two suffix vertices, or two middle vertices differ by less than $M_D + L$, and a prefix-to-middle or middle-to-suffix pair differs by less than $M_D + 1 < M_D + L$. The route through the other long arm is at least as long as the route through A , because the other long arm has total length larger than L .

Finally take $b_i \in V(B) \setminus \{s, t\}$ and $c_j \in V(C) \setminus \{s, t\}$. Direct comparison of the four possible routes gives the following criteria for the named route to be geodesic:

$$\begin{aligned} S: & \quad X_i \leq L, \quad Y_j \leq L, \quad X_i + Y_j \leq 0, \\ T: & \quad X_i \geq -L, \quad Y_j \geq -L, \quad X_i + Y_j \geq 0, \\ U: & \quad X_i \leq -L, \quad Y_j \geq L, \\ V: & \quad X_i \geq L, \quad Y_j \leq -L. \end{aligned}$$

The rectangles $B^- \times C^+$ and $B^+ \times C^-$ are exactly the U and V regions, and the potential ranges give the corresponding inequalities. Suppose next that $b_i \in B^-$ and $c_j \notin C^+$. Then $X_i \leq -L$, while $Y_j < L$ because c_j is either in C^- or in C^0 ; hence the three inequalities for S hold. The case $c_j \in C^-$ and $b_i \notin B^+$ is symmetric. Similarly, if $b_i \in B^+$ and $c_j \notin C^-$, then $X_i \geq L$ and $Y_j > -L$, so the inequalities for T hold; the case $c_j \in C^+$ and $b_i \notin B^-$ is symmetric. The only remaining pairs lie in $B^0 \times C^0$, where the sign of $X_i + Y_j$ gives exactly the prescribed Ferrers S/T matrix. Hence every cross-arm prescribed path is a geodesic. This completes the proof. $\square$

The remaining positive case is the exceptional graph with three arms of length three. Its proof uses the same potentials as Theorem 4.1, but the chosen s - t arm now has two internal vertices and therefore produces two thresholds instead of one.

Proposition 4.2. *The graph $\Theta_{3,3,3}$ is metrizable.*

Proof. Let $\mathcal{P}$ be a consistent path system on $\Theta_{3,3,3}$. By Proposition 2.4, we may assume that $\mathcal{P}$ is neighborly. The prescribed s - t path is one of the three length-three arms. By symmetry, write it as

$$A : s - a - b - t.$$

Write the other two arms as

$$B : s = b_0, b_1, b_2, b_3 = t, \quad C : s = c_0, c_1, c_2, c_3 = t.$$

For $D \in \{B, C\}$, written as $D : s = d_0, d_1, d_2, d_3 = t$, define

$$D^- = \{s\} \cup \{d_i : 1 \leq i \leq 2, P_{d_i, t} = d_i D s a b t\},$$

$$D^+ = \{t\} \cup \{d_i : 1 \leq i \leq 2, P_{s, d_i} = s a b t D d_i\},$$

and set $D^0 = D \setminus (D^- \cup D^+)$. As in the proof of Theorem 4.1, consistency implies that D^- is a prefix, D^+ is a suffix, and $D^- \cap D^+ = \emptyset$. Since $\mathcal{P}$ is neighborly, $d_2 \notin D^-$ and $d_1 \notin D^+$; in particular D^0 has at most two vertices.

For comparable pairs $d_i \in D^-$ and $d_j \in D^+$ with $i < j$, let F_D be the set of pairs for which P_{d_i, d_j} uses the arm A , i.e.

$$P_{d_i, d_j} = d_i D s a b t D d_j.$$

Then F_D is a difference Ferrers relation. Indeed, if $(d_i, d_j) \in F_D$, $i' \leq i$, and $j' \geq j$, then the $d_{i'}-d_{j'}$ subpath of P_{d_i, d_j} is $d_{i'} D s a b t D d_{j'}$, so consistency

puts $(d_{i'}, d_{j'})$ in F_D . The row s and the column t are contained in F_D by the definitions of D^+ and D^- and by $P_{s,t} = A$.

Next consider cross-arm pairs $b_i \in V(B) \setminus \{s, t\}$ and $c_j \in V(C) \setminus \{s, t\}$. There are four relevant routes:

$$\begin{aligned} S &: b_i B s C c_j, & T &: b_i B t C c_j, \\ U &: b_i B s a b t C c_j, & V &: b_i B t b a s C c_j. \end{aligned}$$

The same consistency argument as in Theorem 4.1 gives

$$\begin{aligned} P_{b_i, c_j} = U &\iff b_i \in B^- \text{ and } c_j \in C^+, \\ P_{b_i, c_j} = V &\iff b_i \in B^+ \text{ and } c_j \in C^-. \end{aligned}$$

Outside these two rectangles the prescribed route is S or T , and on the central rectangle $B^0 \times C^0$ the S -region is an additive Ferrers relation: if $P_{b_i, c_j} = S$, $i' \leq i$, and $j' \leq j$, then $P_{b_{i'}, c_{j'}}$ contains $b_{i'} B s C c_{j'}$ as a subpath, so $P_{b_{i'}, c_{j'}} = S$.

We now record the two cuts produced by the internal vertices a and b of the arm A . For a vertex $v \in B \cup C$, let

$$v \in S_a \iff P_{a,v} \text{ begins with } as,$$

and

$$v \in S_b \iff P_{b,v} \text{ begins with } ba.$$

Consistency gives

$$D^- \subseteq S_b \cap D \subseteq S_a \cap D \subseteq D \setminus D^+ \quad (D = B, C).$$

For instance, if $v \in D^-$, then $P_{v,t} = v D s a b t$ contains the subpaths $v D s a$ and $v D s a b$, forcing both $P_{a,v}$ and $P_{b,v}$ to go through s . If $v \in D^+$, the path $P_{s,v} = s a b t D v$ forces both $P_{a,v}$ and $P_{b,v}$ to go through t . Finally, $S_b \subseteq S_a$ because a path $P_{b,v}$ beginning with ba contains the $a-v$ subpath beginning with as . On each D^0 , both $S_a \cap D^0$ and $S_b \cap D^0$ are prefixes, again by taking subpaths.

Apply Lemma 3.8 to the additive Ferrers S -region on $B^0 \times C^0$, with the nested prefixes

$$S_b \cap B^0 \subseteq S_a \cap B^0, \quad S_b \cap C^0 \subseteq S_a \cap C^0.$$

We obtain strictly increasing middle potentials X_i on B^0 , strictly increasing middle potentials Y_j on C^0 , and thresholds $h_b < h_a$ such that

$$P_{b_i, c_j} = S \implies X_i + Y_j \leq 0, \quad P_{b_i, c_j} = T \implies X_i + Y_j \geq 0$$

for $b_i \in B^0$, $c_j \in C^0$, and such that the two prefix cuts are represented by the same thresholds:

$$\begin{aligned} v \in S_b &\implies Z_v \leq h_b, & v \notin S_b &\implies Z_v \geq h_b, \\ v \in S_a &\implies Z_v \leq h_a, & v \notin S_a &\implies Z_v \geq h_a, \end{aligned}$$

where $Z_{b_i} = X_i$ and $Z_{c_j} = Y_j$ on the central vertices. Multiplying all middle potentials and the two thresholds by a sufficiently small positive constant, assume

$$|X_i| < 1, \quad |Y_j| < 1, \quad |h_b| < 1, \quad |h_a| < 1.$$

Set $L = 2$. Apply Lemma 3.6 separately to the difference Ferrers relations F_B and F_C , preserving the middle potentials just chosen. We obtain strictly increasing sequences

$$\begin{aligned} -M_B &= X_0 < X_1 < X_2 < X_3 = M_B, \\ -M_C &= Y_0 < Y_1 < Y_2 < Y_3 = M_C, \end{aligned}$$

where $M_B, M_C > L$, all non-end prefix values are below $-L$, all non-end suffix values are above L , and the same-arm paths through A are represented by

$$\begin{aligned} (b_i, b_j) \in F_B &\iff X_j - X_i > M_B + L, \\ (c_i, c_j) \in F_C &\iff Y_j - Y_i > M_C + L. \end{aligned}$$

Define the weights on the chosen arm A by

$$w(sa) = \frac{L - h_a}{2}, \quad w(ab) = \frac{h_a - h_b}{2}, \quad w(bt) = \frac{L + h_b}{2}.$$

These three numbers are positive because $|h_a|, |h_b| < L$ and $h_b < h_a$, and their sum is L . On the two remaining arms define

$$w(b_{i-1}b_i) = \frac{X_i - X_{i-1}}{2}, \quad w(c_{j-1}c_j) = \frac{Y_j - Y_{j-1}}{2}.$$

All these weights are positive. The total lengths of B and C are M_B and M_C , and

$$d_B(s, b_i) - d_B(t, b_i) = X_i, \quad d_C(s, c_j) - d_C(t, c_j) = Y_j.$$

We check that every prescribed path is a geodesic. First, the subpaths of A are geodesics: any alternative to a proper subpath of A must use one of the

other arms, whose total length is larger than L , while every proper subpath of A has length smaller than L ; and $P_{s,t} = A$ has length $L < M_B, M_C$.

Consider a and a vertex $v = d_i$ on $D \in \{B, C\}$. Let E be the other long arm, and let Z_i , M_D , and M_E denote the corresponding potential and total arm lengths. The two direct routes through s and through t have lengths

$$R_s^a = \frac{L - h_a}{2} + \frac{M_D + Z_i}{2},$$

$$R_t^a = \frac{h_a - h_b}{2} + \frac{L + h_b}{2} + \frac{M_D - Z_i}{2}.$$

Thus $R_s^a \leq R_t^a$ exactly when $Z_i \leq h_a$. If $P_{a,v}$ begins with as , the threshold construction, together with the inclusion $D^- \subseteq S_a$, gives $Z_i \leq h_a$; if $P_{a,v}$ begins with ab , it gives $Z_i \geq h_a$.

It remains only to compare with the two routes that use the other long arm. The route $a - s - E - t - D - v$ has length $R_s^a + M_E - Z_i$, and the route $a - b - t - E - s - D - v$ has length $R_t^a + M_E + Z_i$. If $Z_i \leq h_a$, then

$$M_E - Z_i > 0, \quad (R_t^a + M_E + Z_i) - R_s^a = M_E + h_a > 0.$$

If $Z_i \geq h_a$, then

$$M_E + Z_i > 0, \quad (R_s^a + M_E - Z_i) - R_t^a = M_E - h_a > 0.$$

Here we use $M_E > L > 1$ and $|h_a| < 1$, together with the facts that prefix potentials are $< -L$, suffix potentials are $> L$, and middle potentials lie in $(-1, 1)$. Hence every prescribed $a-v$ path is a geodesic.

The argument for b is identical, using the lower threshold. The route from b to $v = d_i$ through s has length

$$R_s^b = \frac{h_a - h_b}{2} + \frac{L - h_a}{2} + \frac{M_D + Z_i}{2},$$

whereas the route through t has length

$$R_t^b = \frac{L + h_b}{2} + \frac{M_D - Z_i}{2}.$$

Thus $R_s^b \leq R_t^b$ exactly when $Z_i \leq h_b$. The two routes through the other long arm are longer by the same calculation with h_b in place of h_a . Hence every prescribed $b-v$ path is a geodesic.

For two vertices d_i, d_j on the same arm $D \in \{B, C\}$, with $i < j$, the direct subpath of D has length $(Z_j - Z_i)/2$, while the route using A has length

$$M_D + L - \frac{Z_j - Z_i}{2}.$$

The route using the other long arm is still longer, since that arm has total length larger than L . Therefore the route through A is geodesic exactly when $Z_j - Z_i \geq M_D + L$, and the direct route is geodesic exactly when $Z_j - Z_i \leq M_D + L$. If the prescribed path uses A , then $(d_i, d_j) \in F_D$ and Lemma 3.6 gives $Z_j - Z_i > M_D + L$. If the prescribed path is direct and $(d_i, d_j) \in D^- \times D^+$, the same lemma gives $Z_j - Z_i \leq M_D + L$. In all other direct cases the inequality $Z_j - Z_i \leq M_D + L$ follows from the potential ranges: two prefix vertices, two suffix vertices, or two middle vertices differ by less than $M_D + L$, and a prefix-to-middle or middle-to-suffix pair differs by less than $M_D + 1 < M_D + L$.

Finally take $b_i \in V(B) \setminus \{s, t\}$ and $c_j \in V(C) \setminus \{s, t\}$. Direct comparison of the four routes S, T, U, V gives the following sufficient and necessary criteria for the named route to be geodesic:

$$\begin{aligned} S: & \quad X_i \leq L, \quad Y_j \leq L, \quad X_i + Y_j \leq 0, \\ T: & \quad X_i \geq -L, \quad Y_j \geq -L, \quad X_i + Y_j \geq 0, \\ U: & \quad X_i \leq -L, \quad Y_j \geq L, \\ V: & \quad X_i \geq L, \quad Y_j \leq -L. \end{aligned}$$

The rectangles $B^- \times C^+$ and $B^+ \times C^-$ are exactly the U and V regions, and the potential ranges give the corresponding inequalities. If $b_i \in B^-$ and $c_j \notin C^+$, then $X_i \leq -L$ and $Y_j < L$, so S is geodesic; the case $c_j \in C^-$ and $b_i \notin B^+$ is symmetric. If $b_i \in B^+$ and $c_j \notin C^-$, then $X_i \geq L$ and $Y_j > -L$, so T is geodesic; the case $c_j \in C^+$ and $b_i \notin B^-$ is symmetric. The only remaining pairs lie in $B^0 \times C^0$, where the two-threshold Ferrers representation gives the required weak sign of $X_i + Y_j$. These weak inequalities are sufficient because metrizable allows ties among shortest paths. Thus every prescribed cross-arm path is a geodesic.

All unordered pairs of vertices have now been checked, so the weighting w induces $\mathcal{P}$. Hence every neighborly consistent path system on $\Theta_{3,3,3}$ is metrizable, and the non-neighborly systems are metrizable by Proposition 2.4. $\square$

Proof of Theorem 1.1. If $a = 1$, then $\Theta_{a,b,c}$ is outerplanar, hence metrizable by Proposition 2.3. If $a = 2$, metrizability follows from Theorem 4.1. The remaining positive case $(a, b, c) = (3, 3, 3)$ is Proposition 4.2.

Conversely, if $a \geq 3$, $b \geq 3$, and $c \geq 4$, then non-metrizability follows from Proposition 2.6. These alternatives are exhaustive when $a \leq b \leq c$. $\square$

5. Concluding remarks

The classification shows that, among theta graphs, the first obstruction is exactly the six-inequality block realized by $\Theta_{3,3,4}$. All longer theta graphs with three arms of lengths at least 3, 3, 4 inherit this obstruction as a topological minor. On the positive side, a length-two arm suppresses the possible obstruction: the proof for $\Theta_{2,b,c}$ turns consistency into Ferrers relations and realizes them by one-dimensional potentials.

The exceptional case $\Theta_{3,3,3}$ shows that the potential method is not limited to a single short internal vertex. The two internal vertices of the chosen length-three arm give two nested prefix cuts, and the fact that the remaining central chains have size at most two is exactly what permits the two-threshold Ferrers representation.

Funding

This work was supported by the Natural Science Foundation of Shandong Province (Grant No. ZR2024MA073).

Data availability

No new research data were generated or analyzed in this theoretical study. All arguments are contained in the manuscript.

Declaration of competing interest

The author declares no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this submission, ChatGPT was used to assist with journal-format conversion and language editing. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the submitted version.

CRediT author statement

Guangfu Wang: Conceptualization, Methodology, Formal analysis, Investigation, Writing - original draft, Writing - review and editing, Funding acquisition.

References

- [1] G. Bodwin, On the structure of unique shortest paths in graphs, in *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, SIAM, 2019, pp. 2071–2089. doi:10.1137/1.9781611975482.125.
- [2] M. Chudnovsky, D. Cizma, and N. Linial, Strictly metrizable graphs are minor-closed, arXiv:2501.08277v2, 2025. doi:10.48550/arXiv.2501.08277.
- [3] M. Chudnovsky, D. Cizma, and N. Linial, The structure of metrizable graphs, *Discrete & Computational Geometry* **75** (2026), no. 1, 73–92. doi:10.1007/s00454-024-00685-3.
- [4] D. Cizma and N. Linial, Geodesic geometry on graphs, *Discrete & Computational Geometry* **68** (2022), no. 1, 298–347. doi:10.1007/s00454-021-00345-w.
- [5] D. Cizma and N. Linial, Irreducible non-metrizable path systems in graphs, *Journal of Graph Theory* **102** (2023), no. 1, 5–14. doi:10.1002/jgt.22854.
- [6] D. Cizma and N. Linial, Metric approximations of consistent path systems, arXiv:2601.21982v2, 2026. doi:10.48550/arXiv.2601.21982.
- [7] D. Cizma and N. Linial, On the number of path systems, *Israel Journal of Mathematics*, to appear, 2026. arXiv:2508.21379v3.
- [8] R. Ehrenborg and S. van Willigenburg, Enumerative properties of Ferrers graphs, *Discrete & Computational Geometry* **32** (2004), no. 4, 481–492. doi:10.1007/s00454-004-1135-1.
- [9] P. L. Hammer, U. N. Peled, and X. Sun, Difference graphs, *Discrete Applied Mathematics* **28** (1990), no. 1, 35–44. doi:10.1016/0166-218X(90)90092-Q.