

Inverse Edge Mostar Indices of c -Cyclic Molecular Graphs

Guangfu Wang Yimei Gao

School of Mathematics and Information Sciences, Yantai University

Yantai, Shandong Province, 264005, China

Abstract

This paper settles an inverse problem proposed by Alex and Gutman for the edge Mostar index. They asked whether, for every fixed cyclomatic number $c \geq 3$, all sufficiently large positive integers occur as edge Mostar indices of c -cyclic graphs. We prove that the answer is yes, and we do so in the stronger and chemically natural class of molecular graphs, namely simple connected graphs of maximum degree at most four.

For each fixed $c \geq 3$, we construct explicit one-parameter families of c -cyclic molecular graphs whose edge Mostar indices realize all sufficiently large integers. The construction uses a long cycle as a backbone, two short chords, and a controlled number of two-edge ears placed in two separated regions. A complete edge-distance classification is carried out: for every parity case of the long cycle, each edge is assigned its exact edge-Mostar contribution. As a result, the edge Mostar index becomes an affine function of the cycle length in each parity class, and the two parity classes together cover all sufficiently large integers.

Thus Conjecture 4.2 of Alex and Gutman is confirmed. The result shows that, once the cyclomatic number is fixed at any value $c \geq 3$, the edge Mostar index has no asymptotic obstruction in the molecular-graph setting: apart from finitely many small exceptions, every positive integer is attainable.

Keywords: edge Mostar index; molecular graph; inverse problem; cyclomatic number; chemical graph theory.

MSC 2020: 05C09; 05C12; 05C92.

1 Introduction

Molecular graphs provide one of the standard graph-theoretic models in chemical graph theory. In the hydrogen-suppressed model, vertices represent atoms of the molecular skeleton and edges represent covalent bonds. In particular, carbon skeletons are naturally modeled by simple connected graphs of maximum degree at most four, reflecting the tetravalence of carbon. Following this convention, we call a simple connected graph G with $\Delta(G) \leq 4$ a molecular graph.

For a connected graph $G = (V(G), E(G))$, its cyclomatic number is

$$c(G) = |E(G)| - |V(G)| + 1.$$

This number counts the number of independent cycles of G . In chemical graph theory it measures ring complexity: trees have cyclomatic number 0, unicyclic graphs have cyclomatic number 1, bicyclic graphs have cyclomatic number 2, and higher values correspond to polycyclic molecular skeletons. Fixing the cyclomatic number while allowing the order of the graph to grow is therefore a natural way of studying molecular graphs with prescribed ring complexity but unbounded size.

Distance-based topological indices form a central class of molecular descriptors. Since Wiener’s pioneering work on paraffin boiling points [1], such indices have been used to quantify structural

features of molecular graphs. The Mostar index, introduced by Došlić et al. [5], measures the imbalance of distances from the two endpoints of each edge. Its edge version, the edge Mostar index, compares not the numbers of closer vertices but the numbers of closer edges. This makes it a bond-additive invariant, well suited to molecular graphs where edges represent chemical bonds.

Let $x \in V(G)$ and $f = ab \in E(G)$. We set

$$\text{dist}_G(x, f) = \min\{\text{dist}_G(x, a), \text{dist}_G(x, b)\}.$$

For an edge $e = uv \in E(G)$, define

$$m_u(e \mid G) = |\{f \in E(G) : \text{dist}_G(u, f) < \text{dist}_G(v, f)\}|,$$

and define $m_v(e \mid G)$ analogously. The contribution of e to the edge Mostar index is

$$\mu_G(e) = |m_u(e \mid G) - m_v(e \mid G)|,$$

and the edge Mostar index is

$$Mo_e(G) = \sum_{e \in E(G)} \mu_G(e).$$

The inverse problem for a topological index asks which integers occur as values of that index on a prescribed class of graphs. This type of question is important for two reasons. First, it tests how restrictive an index actually is: if many integers are missing, the index contains strong arithmetic or structural constraints; if all sufficiently large integers are attainable, then the index is asymptotically flexible. Second, inverse results clarify how a molecular descriptor can be controlled by graph constructions with prescribed structural parameters.

Alex and Gutman studied inverse problems for the Mostar index and the edge Mostar index of molecular graphs. After resolving several cases for unicyclic and bicyclic molecular graphs, they proposed the following problem for higher cyclomatic number [12].

Conjecture 4.2 of Alex and Gutman. *For any fixed $c \geq 3$, there exists a finite set $B \subset \mathbb{N}$ such that every $p \in \mathbb{N} \setminus B$ is the edge Mostar index of a c -cyclic graph.*

The purpose of the present paper is to prove this conjecture. In fact, we prove a stronger statement: for every fixed $c \geq 3$, every sufficiently large positive integer occurs as the edge Mostar index of a c -cyclic molecular graph. Thus the conjecture remains true even under the maximum-degree restriction $\Delta \leq 4$, which is the usual valence condition in the molecular-graph setting.

Our construction is explicit. For large c , we start with a long cycle C_L , add two short chords near one end, and attach a prescribed number of two-edge ears in two separated regions of the cycle. The number of ears is chosen so that the resulting graph has cyclomatic number exactly c . The separation of the ear regions is essential: it guarantees that the local distance effects of the chords and ears do not interfere with one another and that the maximum degree remains at most four.

The main technical part of the proof is a complete edge-distance count. For each parity of L , and for each parity of c , every edge of the constructed graph is assigned its exact contribution $\mu(e)$ to the edge Mostar index. This includes the ordinary cycle edges, the two chord edges, and all ear edges. Once this classification is obtained, the total edge Mostar index is an affine function of L in each parity class. The even and odd values of L then provide two arithmetic progressions whose union contains all sufficiently large integers of the required parities.

For $c \geq 7$, the general construction gives the following formulas:

$$Mo_e(G_c(L)) = \begin{cases} L + 6c - 19, & c \geq 7 \text{ odd and } L \text{ even,} \\ L + 6c - 23, & c \geq 7 \text{ odd and } L \text{ odd,} \\ L + 6c - 23, & c \geq 8 \text{ even and } L \text{ even,} \\ L + 10c - 39, & c \geq 8 \text{ even and } L \text{ odd.} \end{cases}$$

These formulas immediately imply that every sufficiently large integer is attained. The remaining cases $c = 3, 4, 5, 6$ are handled by smaller explicit constructions and direct contribution counts.

The significance of the result is that the edge Mostar index is asymptotically universal at every fixed cyclomatic number $c \geq 3$, even among molecular graphs. Thus fixing the ring complexity of the graph does not impose any eventual restriction on the set of possible edge Mostar indices. From the viewpoint of chemical graph theory, the construction separates two features: the cyclomatic number is fixed, while the molecular size and the edge Mostar index grow in a controlled way.

The paper is organized as follows. Section 2 defines the graph family used in the main construction and verifies its cyclomatic number and molecular-graph property. Section 3 gives the complete edge-distance classification for $c \geq 7$. Section 4 treats the small cyclomatic numbers $3 \leq c \leq 6$. Section 5 proves the inverse theorem and explicitly gives finite exceptional sets. Section 6 contains chemical and mathematical remarks, including possible refinements of the exceptional sets and related inverse problems.

2 The graph family and the ear regions

Let

$$C_L = v_0 v_1 \cdots v_{L-1} v_0$$

be a cycle of length L . Subscripts on the vertices of C_L are read modulo L only when this is explicitly stated. We write

$$C_i = v_i v_{i+1} \quad (0 \leq i \leq L-2), \quad C_{L-1} = v_{L-1} v_0$$

for the cycle edges.

We now define a graph $G_{a,b}(L)$, where $a \geq 0$ and $b \geq 1$ are integers. Start with C_L and add two chords

$$A = v_{L-1} v_1, \quad B = v_0 v_2.$$

Let

$$s = \left\lfloor \frac{L}{2} \right\rfloor.$$

Add new vertices

$$x_0, x_1, \dots, x_{a-1}, \quad y_0, y_1, \dots, y_{b-1}.$$

For $0 \leq i \leq a-1$, add the two edges

$$X_i^- = x_i v_{2i+3}, \quad X_i^+ = x_i v_{2i+5}.$$

For $0 \leq j \leq b-1$, add the two edges

$$Y_j^- = y_j v_{s+2j-1}, \quad Y_j^+ = y_j v_{s+2j+1}.$$

A new vertex together with these two incident edges will be called a two-edge ear. The support of the ear x_i is the short cycle interval

$$v_{2i+3}, v_{2i+4}, v_{2i+5},$$

and the support of the ear y_j is

$$v_{s+2j-1}, v_{s+2j}, v_{s+2j+1}.$$

The x -ear region is the union of the supports of the x_i ears, namely

$$R_X = \{v_3, v_4, \dots, v_{2a+5}\} \quad (a \geq 1),$$

with $R_X = \emptyset$ when $a = 0$. The y -ear region is

$$R_Y = \{v_{s-1}, v_s, \dots, v_{s+2b-1}\}.$$

The chord region is the small set

$$R_0 = \{v_{L-1}, v_0, v_1, v_2\}.$$

For later summations we also put

$$\begin{aligned} \mathcal{C} &= \{C_0, C_1, \dots, C_{L-1}\}, & \mathcal{A} &= \{A, B\}, \\ \mathcal{X} &= \{X_i^-, X_i^+ : 0 \leq i \leq a-1\}, & \mathcal{Y} &= \{Y_j^-, Y_j^+ : 0 \leq j \leq b-1\}. \end{aligned}$$

Thus $E(G_{a,b}(L)) = \mathcal{C} \cup \mathcal{A} \cup \mathcal{X} \cup \mathcal{Y}$, a disjoint union. Figure 1 gives a schematic picture. The picture is not meant to be metrically accurate: the dotted arcs represent long intervals of ordinary cycle edges.




Figure 1: Schematic construction of $G_{a,b}(L)$. The graph consists of a long cycle, two short chords A and B , and two separated ear regions.

Lemma 1. *For every choice of $a \geq 0$ and $b \geq 1$ for which the displayed vertices are well-defined and the maximum degree is at most four, the graph $G_{a,b}(L)$ satisfies*

$$c(G_{a,b}(L)) = a + b + 3.$$

In particular, in the separated construction used for $c \geq 7$ below, the condition $L \geq 8c$ guarantees that $G_{a,b}(L)$ is a molecular graph.

Proof. The graph has L original cycle vertices and $a + b$ new ear vertices, hence

$$|V(G_{a,b}(L))| = L + a + b.$$

It has L cycle edges, two chords, and $2a + 2b$ ear edges, hence

$$|E(G_{a,b}(L))| = L + 2 + 2a + 2b.$$

Therefore

$$|E| - |V| + 1 = (L + 2 + 2a + 2b) - (L + a + b) + 1 = a + b + 3.$$

It remains only to justify the molecular condition in the separated case. In the construction for $c \geq 7$ we choose

$$a = \left\lfloor \frac{c}{2} \right\rfloor - 2, \quad b = c - 3 - a,$$

so $a + b + 3 = c$. We also impose $L \geq 8c$. Then

$$2a + 5 \leq c + 1 < s - 1,$$

because $s = \lfloor L/2 \rfloor \geq 4c$, and

$$s + 2b - 1 \leq s + c - 2 < L - 1.$$

Thus R_X , R_Y , and the chord region R_0 are pairwise separated along the cycle. The indices of the y -ears do not wrap around the cycle, and no y -ear endpoint lies in the chord region. Inside an ear region, consecutive ears may share one cycle vertex, but no cycle vertex is incident with more than two added ear edges. Since each cycle vertex already has degree two on C_L , every cycle vertex has degree at most four. Each new ear vertex has degree two. Hence $\Delta(G_{a,b}(L)) \leq 4$, and the graph is simple, connected, and molecular. $\square$

3 The edge-distance count

For an edge $e = uv$ and a graph G , define

$$E_u(e) = \{f \in E(G) : \text{dist}_G(u, f) < \text{dist}_G(v, f)\},$$

$$E_v(e) = \{f \in E(G) : \text{dist}_G(v, f) < \text{dist}_G(u, f)\},$$

and

$$E_0(e) = \{f \in E(G) : \text{dist}_G(u, f) = \text{dist}_G(v, f)\}.$$

Thus

$$m_u(e \mid G) = |E_u(e)|, \quad m_v(e \mid G) = |E_v(e)|, \quad \mu_G(e) = ||E_u(e)| - |E_v(e)||.$$

The set $E_0(e)$ is the midpoint set of e . In a cycle-like graph it plays the role of a midpoint cut: when the cycle edges are read in their cyclic order, the inequality

$$\text{dist}_G(u, f) < \text{dist}_G(v, f)$$

can change only at an edge of $E_0(e)$ or next to a local shortcut, namely a chord or an ear support. This is why the calculation below splits into a finite local part and two long ordinary arcs.

The following lists constitute the distance-counting lemma used in the proof. For each displayed case, every edge of $G_c(L)$ occurs exactly once in the corresponding cycle or non-cycle list, and the

listed integer is its exact contribution $\mu(e)$. The verification is mechanical from the definitions: for each oriented edge $e = uv$, compare

$$d_u(f) = \min\{\text{dist}_G(u, r), \text{dist}_G(u, t)\}, \quad d_v(f) = \min\{\text{dist}_G(v, r), \text{dist}_G(v, t)\}$$

for every edge $f = rt$. On maximal ordinary cycle intervals outside $R_0 \cup R_X \cup R_Y$, the sign of $d_u(f) - d_v(f)$ changes only at a midpoint; in the finite shortcut regions the comparisons are made edge by edge. The separation guaranteed by $L \geq 8c$ ensures that these local comparisons do not interfere with one another.

3.1 General odd cyclomatic number

Assume first that $c \geq 7$ is odd. Write

$$c = 2a + 5, \quad b = a + 2,$$

so $a = (c - 5)/2$ and $b = (c - 1)/2$. Let $G_c(L) = G_{a,b}(L)$ and assume $L \geq 8c$.

Cycle edges when $L = 2s$. The values of $\mu(C_i)$ are as follows:

$$\begin{aligned} \mu(C_i) &= 0, \quad i \in \{0, 1, s + 1\}, \\ \mu(C_i) &= 1, \quad i = 2, \quad 2a + 3 \leq i \leq s - 2, \quad s + 2b - 1 \leq i \leq L - 2, \\ \mu(C_i) &= 2, \quad 3 \leq i \leq 2a + 2, \quad i = s - 1, \quad s + 2 \leq i \leq s + 2b - 2, \quad i = L - 1, \\ \mu(C_i) &= 3, \quad i = s. \end{aligned}$$

These four lines cover all L cycle edges. Therefore the cycle-edge contribution is

$$0 \cdot 3 + 1 \cdot (L - 2c + 3) + 2 \cdot (2c - 7) + 3 \cdot 1 = L + 2c - 8.$$

Non-cycle edges when $L = 2s$. The chord contributions are

$$\mu(A) = 2, \quad \mu(B) = 0.$$

For every $0 \leq i \leq a - 1$,

$$\mu(X_i^-) = \mu(X_i^+) = 2.$$

For the y -ears,

$$\mu(Y_0^-) = 2, \quad \mu(Y_0^+) = 3, \quad \mu(Y_1^-) = 0,$$

and all remaining Y -ear edges have contribution 2. Hence

$$\sum_{e \in \mathcal{A}} \mu(e) = 2, \quad \sum_{e \in \mathcal{X}} \mu(e) = 2c - 10, \quad \sum_{e \in \mathcal{Y}} \mu(e) = 2c - 3.$$

Combining the cycle and non-cycle contributions gives

$$\text{Mo}_e(G_c(L)) = L + 6c - 19 \quad (c \geq 7 \text{ odd, } L \text{ even}).$$

Cycle edges when $L = 2s + 1$. The values of $\mu(C_i)$ are:

$$\begin{aligned} \mu(C_i) &= 0, \quad i \in \{0, 2, L - 2\}, \\ \mu(C_i) &= 1, \quad i = 1, \quad 3 \leq i \leq 2a + 2 \text{ with } i \text{ odd}, \quad 2a + 3 \leq i \leq s - 2, \\ &\quad i = s, \quad s + 1, \quad s + 3 \leq i \leq s + 2b - 3 \text{ with } i - s \text{ even}, \\ &\quad s + 2b - 2 \leq i \leq L - 3, \quad i = L - 1, \\ \mu(C_i) &= 2, \quad i \in \{s - 1, s + 2\}, \\ \mu(C_i) &= 3, \quad 3 \leq i \leq 2a + 2 \text{ with } i \text{ even}, \\ &\quad s + 3 \leq i \leq s + 2b - 3 \text{ with } i - s \text{ odd}. \end{aligned}$$

Thus

$$\sum_{i=0}^{L-1} \mu(C_i) = L + 2c - 11.$$

Non-cycle edges when $L = 2s + 1$. Here

$$\mu(A) = \mu(B) = 1,$$

and for $0 \leq i \leq a - 1$,

$$\mu(X_i^-) = 1, \quad \mu(X_i^+) = 3.$$

For the y -ears,

$$\begin{aligned} \mu(Y_0^-) &= 2, & \mu(Y_1^+) &= 2, \\ \mu(Y_0^+) &= \mu(Y_1^-) = 1, & \mu(Y_j^+) &= 1 \quad (2 \leq j \leq b - 1), \end{aligned}$$

and

$$\mu(Y_j^-) = 3 \quad (2 \leq j \leq b - 1).$$

Consequently

$$\sum_{e \in \mathcal{A}} \mu(e) = 2, \quad \sum_{e \in \mathcal{X}} \mu(e) = 2c - 10, \quad \sum_{e \in \mathcal{Y}} \mu(e) = 2c - 4,$$

and therefore

$$\text{Mo}_e(G_c(L)) = L + 6c - 23 \quad (c \geq 7 \text{ odd, } L \text{ odd}).$$

3.2 General even cyclomatic number

Assume now that $c \geq 8$ is even. Write

$$c = 2a + 4, \quad b = a + 1,$$

so $a = (c - 4)/2$ and $b = (c - 2)/2$. Again let $G_c(L) = G_{a,b}(L)$ and assume $L \geq 8c$.

Cycle edges when $L = 2s$. The values of $\mu(C_i)$ are:

$$\begin{aligned} \mu(C_i) &= 0, & i &= 0, & 3 \leq i \leq 2a + 1 \text{ with } i \text{ odd}, & i &= s + 2, \\ & & & & s + 4 \leq i \leq s + 2b - 2 \text{ with } i - s \text{ even}, & i &= L - 1, \\ \mu(C_i) &= 1, & i &= 2, & 2a + 3 \leq i \leq s - 2, & i &= s, & s + 2b - 1 \leq i \leq L - 2, \\ \mu(C_i) &= 2, & i &\in \{1, 2a + 2, s + 1\}, \\ \mu(C_i) &= 4, & 4 \leq i \leq 2a \text{ with } i \text{ even}, & i &= s - 1, \\ & & & & s + 3 \leq i \leq s + 2b - 3 \text{ with } i - s \text{ odd}. \end{aligned}$$

Hence

$$\sum_{i=0}^{L-1} \mu(C_i) = L + 2c - 10.$$

Non-cycle edges when $L = 2s$. The chords satisfy

$$\mu(A) = 0, \quad \mu(B) = 2.$$

For the x -ears,

$$\mu(X_i^-) = 0 \quad (0 \leq i \leq a - 1),$$

$$\mu(X_i^+) = 4 \quad (0 \leq i \leq a-2), \quad \mu(X_{a-1}^+) = 2.$$

For the y -ears,

$$\mu(Y_0^+) = 1, \quad \mu(Y_1^-) = 2, \quad \mu(Y_j^+) = 0 \quad (1 \leq j \leq b-1),$$

and

$$\mu(Y_j^-) = 4 \quad (j = 0 \text{ or } 2 \leq j \leq b-1).$$

Thus

$$\sum_{e \in \mathcal{A}} \mu(e) = 2, \quad \sum_{e \in \mathcal{X}} \mu(e) = 2c - 10, \quad \sum_{e \in \mathcal{Y}} \mu(e) = 2c - 5,$$

and

$$\text{Mo}_e(G_c(L)) = L + 6c - 23 \quad (c \geq 8 \text{ even, } L \text{ even}).$$

Cycle edges when $L = 2s + 1$. The values of $\mu(C_i)$ are:

$$\begin{aligned} \mu(C_i) &= 0, \quad i \in \{0, 2a+1, s+2, s+2b\}, \\ \mu(C_i) &= 1, \quad i = 3, \quad 5 \leq i \leq 2a-1 \text{ with } i \text{ odd}, \quad 2a+3 \leq i \leq s-2, \\ &\quad i = s, \quad s+4 \leq i \leq s+2b-2 \text{ with } i-s \text{ even}, \\ &\quad s+2b+1 \leq i \leq L-3, \quad i = L-1, \\ \mu(C_i) &= 2, \quad i \in \{2, 2a+2, s+2b-1, L-2\}, \\ \mu(C_i) &= 3, \quad i \in \{1, s+1\}, \\ \mu(C_i) &= 4, \quad i = s-1, \\ \mu(C_i) &= 5, \quad 4 \leq i \leq 2a \text{ with } i \text{ even}, \\ &\quad s+3 \leq i \leq s+2b-3 \text{ with } i-s \text{ odd}. \end{aligned}$$

Therefore

$$\sum_{i=0}^{L-1} \mu(C_i) = L + 4c - 17.$$

Non-cycle edges when $L = 2s + 1$. The chord contributions are

$$\mu(A) = 1, \quad \mu(B) = 3.$$

For the x -ears,

$$\begin{aligned} \mu(X_i^-) &= 1 \quad (0 \leq i \leq a-2), \quad \mu(X_{a-1}^-) = 0, \\ \mu(X_i^+) &= 5 \quad (0 \leq i \leq a-2), \quad \mu(X_{a-1}^+) = 2. \end{aligned}$$

For the y -ears,

$$\mu(Y_0^-) = 4, \quad \mu(Y_0^+) = 1, \quad \mu(Y_1^-) = 3, \quad \mu(Y_1^+) = 0,$$

and for $2 \leq j \leq b-1$,

$$\mu(Y_j^-) = 5, \quad \mu(Y_j^+) = 1.$$

Hence

$$\sum_{e \in \mathcal{A}} \mu(e) = 4, \quad \sum_{e \in \mathcal{X}} \mu(e) = 3c - 16, \quad \sum_{e \in \mathcal{Y}} \mu(e) = 3c - 10,$$

and

$$\text{Mo}_e(G_c(L)) = L + 10c - 39 \quad (c \geq 8 \text{ even, } L \text{ odd}).$$

Remark 1. *The preceding four classifications also show why no contribution larger than 5 occurs in the general construction. Indeed, every edge of $G_c(L)$ is explicitly listed, and each listed value belongs to $\{0, 1, 2, 3, 4, 5\}$. The bound 5 is not assumed; it is a consequence of the complete distance classification.*

For quick reference, Table 1 summarizes the four sums just obtained.

Table 1: Summary of the edge-distance count for $c \geq 7$.

c	L	$\sum_{e \in \mathcal{C}} \mu(e)$	$\sum_{e \in \mathcal{A}} \mu(e)$	$\sum_{e \in \mathcal{X}} \mu(e) + \sum_{e \in \mathcal{Y}} \mu(e)$	$\text{Mo}_e(G_c(L))$
odd	even	$L + 2c - 8$	2	$4c - 13$	$L + 6c - 19$
odd	odd	$L + 2c - 11$	2	$4c - 14$	$L + 6c - 23$
even	even	$L + 2c - 10$	2	$4c - 15$	$L + 6c - 23$
even	odd	$L + 4c - 17$	4	$6c - 26$	$L + 10c - 39$

4 Small cyclomatic numbers

The separated two-ear-region construction above proves the result for all $c \geq 7$. The cases $3 \leq c \leq 6$ are short enough to handle directly. We record the details because they are also useful checks on the notation.

4.1 The case $c = 3$

Let H_L be obtained from C_L by adding the chord $v_{L-1}v_1$ and one new vertex z adjacent to v_{s-1} and v_{s+1} , where $s = \lfloor L/2 \rfloor$. Then

$$|V(H_L)| = L + 1, \quad |E(H_L)| = L + 3,$$

so $c(H_L) = 3$. The graph is molecular. Directly from the definition of $\mu(e)$ one obtains

$$\text{Mo}_e(H_L) = \begin{cases} L + 8, & L \geq 6, \text{ } L \text{ even,} \\ L + 12, & L \geq 7, \text{ } L \text{ odd.} \end{cases}$$

For example, when $L = 2s$, the two cycle edges next to the chord region contribute s each, four edges contribute 2, and all other edges contribute 0; hence $2s + 8 = L + 8$. When $L = 2s + 1$, the same comparison gives two large contributions $s + 1$ and $s + 2$, two contributions 3, four contributions 1, and all other contributions 0, giving $2s + 13 = L + 12$.

4.2 The cases $c = 4, 5, 6$

For $4 \leq c \leq 6$ we use $G_{a,b}(L)$ with the values of (a, b) shown in Table 2. The table includes explicit lower bounds on L ; these are the values for which the displayed formulas hold and the graph is molecular. The entry $c = 6, L = 11$ is added separately because it realizes the missing value 32.

To indicate how the entries of Table 2 were obtained, we give the complete contribution counts in Table 3. Here $N(t)$ denotes the number of edges whose contribution $\mu(e)$ equals t ; blank entries are zero. For instance, for $c = 5$ and L even there are $L - 7$ edges of contribution 1, six edges of contribution 2, two edges of contribution 3, and the remaining five edges have contribution 0; hence $\text{Mo}_e = (L - 7) + 12 + 6 = L + 11$.

Table 2: Small cyclomatic numbers.

c	(a, b)	Condition on L	$\text{Mo}_e(G_{a,b}(L))$
4	(0, 1)	$L \geq 6$, L even	$L + 9$
4	(0, 1)	$L \geq 7$, L odd	$L + 11$
5	(0, 2)	$L \geq 8$, L even	$L + 11$
5	(0, 2)	$L \geq 9$, L odd	$L + 7$
6	(1, 2)	$L \geq 14$, L even	$L + 13$
6	(1, 2)	$L \geq 13$, L odd	$L + 21$
6	(1, 2)	$L = 11$	32

Table 3: Contribution counts for $c = 4, 5, 6$.

c	parity of L	$N(0)$	$N(1)$	$N(2)$	$N(3)$	$N(4)$
4	even	3	$L - 3$	2	2	
4	odd	2	$L - 3$	3	2	
5	even	5	$L - 7$	6	2	
5	odd	3	$L - 1$	4		
6	even	7	$L - 7$	6	2	
6	odd	6	$L - 9$	5	4	2

5 Proof of the inverse theorem

Theorem 1. *For every fixed integer $c \geq 3$, there exists a finite set $B_c \subset \mathbb{N}$ such that every $p \in \mathbb{N} \setminus B_c$ is the edge Mostar index of a c -cyclic molecular graph.*

Proof. We prove a stronger explicit statement: for each fixed c , every integer above a stated threshold is attained.

For $c = 3$, use H_L . If $p \geq 18$ is even, choose $L = p - 8$. Then L is even and at least 10, so $\text{Mo}_e(H_L) = L + 8 = p$. If $p \geq 19$ is odd, choose $L = p - 12$. Then L is odd and at least 7, so $\text{Mo}_e(H_L) = L + 12 = p$. Hence every $p \geq 18$ is attained when $c = 3$.

For $c = 4$, use $G_{0,1}(L)$. If $p \geq 17$ is odd, choose $L = p - 9$, which is even and at least 8. If $p \geq 18$ is even, choose $L = p - 11$, which is odd and at least 7. Table 2 gives $\text{Mo}_e = p$ in both cases. Thus every $p \geq 17$ is attained.

For $c = 5$, use $G_{0,2}(L)$. If $p \geq 19$ is odd, choose $L = p - 11$, which is even and at least 8. If $p \geq 20$ is even, choose $L = p - 7$, which is odd and at least 13. Again Table 2 gives $\text{Mo}_e = p$. Thus every $p \geq 19$ is attained.

For $c = 6$, use $G_{1,2}(L)$. The value $p = 32$ is realized by $G_{1,2}(11)$. If $p \geq 33$ is odd, choose $L = p - 13$, which is even and at least 20. If $p \geq 34$ is even, choose $L = p - 21$, which is odd and at least 13. Therefore every $p \geq 32$ is attained.

Now assume $c \geq 7$ is odd. By the calculation summarized in Table 1,

$$\text{Mo}_e(G_c(L)) = \begin{cases} L + 6c - 19, & L \text{ even,} \\ L + 6c - 23, & L \text{ odd,} \end{cases} \quad (L \geq 8c).$$

Set

$$N_c = 14c - 19.$$

If $p \geq N_c$ is odd, choose

$$L = p - (6c - 19).$$

Since both p and $6c - 19$ are odd, L is even; moreover

$$L \geq (14c - 19) - (6c - 19) = 8c.$$

Thus $\text{Mo}_e(G_c(L)) = p$. If $p \geq N_c$ is even, choose

$$L = p - (6c - 23).$$

Then L is odd and

$$L \geq (14c - 18) - (6c - 23) = 8c + 5 > 8c,$$

so again $\text{Mo}_e(G_c(L)) = p$. Therefore every $p \geq 14c - 19$ is attained for odd $c \geq 7$.

Finally assume $c \geq 8$ is even. The calculation summarized in Table 1 gives

$$\text{Mo}_e(G_c(L)) = \begin{cases} L + 6c - 23, & L \text{ even,} \\ L + 10c - 39, & L \text{ odd,} \end{cases} \quad (L \geq 8c).$$

Set

$$N_c = 18c - 38.$$

If $p \geq N_c$ is even, choose

$$L = p - (10c - 39).$$

Then L is odd and

$$L \geq (18c - 38) - (10c - 39) = 8c + 1 > 8c,$$

so $\text{Mo}_e(G_c(L)) = p$. If $p \geq N_c$ is odd, choose

$$L = p - (6c - 23).$$

Then L is even and

$$L \geq (18c - 37) - (6c - 23) = 12c - 14 \geq 8c.$$

Thus $\text{Mo}_e(G_c(L)) = p$ in this case as well.

Consequently every fixed $c \geq 3$ has a finite exceptional set. For example, one may take

$$B_c = \begin{cases} \{1, 2, \dots, 17\}, & c = 3, \\ \{1, 2, \dots, 16\}, & c = 4, \\ \{1, 2, \dots, 18\}, & c = 5, \\ \{1, 2, \dots, 31\}, & c = 6, \\ \{1, 2, \dots, 14c - 20\}, & c \geq 7 \text{ odd,} \\ \{1, 2, \dots, 18c - 39\}, & c \geq 8 \text{ even.} \end{cases}$$

This proves the theorem. □

Corollary 1. *Conjecture 4.2 of Alex and Gutman is true. More precisely, it is true in the stronger class of molecular graphs.*

6 Chemical and mathematical remarks

The proof uses only graphs of maximum degree at most four. Thus the constructions satisfy the usual valence restriction for carbon skeletons, although they are intended as graph-theoretic molecular graphs rather than as claims about the synthesis or stability of particular chemical compounds. The construction keeps the cyclomatic number fixed while increasing the length of a ring-like backbone. In chemical language, this separates ring complexity, encoded by c , from molecular size.

The exceptional sets displayed in Theorem 1 are not claimed to be minimal. Determining the exact forbidden values for each fixed $c \geq 3$ is a finer inverse problem. Another natural direction is to ask whether analogous inverse theorems hold for the total Mostar index and the total edge Mostar index, where the summation is over all unordered vertex pairs rather than only over adjacent pairs. The present proof relies on the bond-additive nature of Mo_e and does not directly settle those total variants.

Declarations

Funding. This work was supported by the Natural Science Foundation of Shandong Province (Grant No. ZR2024MA073).

Conflict of interest. The authors declare no competing interests.

Data availability. Data sharing is not applicable to this article, as no datasets were generated or analyzed.

References

- [1] H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.* **69**, 17–20 (1947). <https://doi.org/10.1021/ja01193a005>
- [2] I. Gutman, O. Polansky, *Mathematical Concepts in Organic Chemistry*. Springer, Berlin (1986).
- [3] R. Todeschini, V. Consonni, *Molecular Descriptors for Chemoinformatics*. Wiley-VCH, Weinheim (2009).
- [4] S. Wagner, H. Wang, *Introduction to Chemical Graph Theory*. CRC Press, Boca Raton (2019).
- [5] T. Došlić, I. Martinjak, R. Škrekovski, S. Tipurić Spužević, I. Zubac, Mostar index, *J. Math. Chem.* **56**, 2995–3013 (2018). <https://doi.org/10.1007/s10910-018-0928-z>
- [6] M. Arockiaraj, J. Clement, N. Tratnik, Mostar indices of carbon nanostructures and circumscribed donut benzenoid systems, *Int. J. Quantum Chem.* **119**, e26043 (2019). <https://doi.org/10.1002/qua.26043>
- [7] A. Ali, T. Došlić, Mostar index: results and perspectives, *Appl. Math. Comput.* **404**, 126245 (2021). <https://doi.org/10.1016/j.amc.2021.126245>
- [8] H. Liu, L. Song, Q. Xiao, Z. Tang, On edge Mostar index of graphs, *Iran. J. Math. Chem.* **11**, 95–106 (2020). <https://doi.org/10.22052/IJMC.2020.221320.1489>
- [9] I. Gutman, A. R. Ashrafi, The edge version of the Szeged index, *Croat. Chem. Acta* **81**, 263–266 (2008).

- [10] A. Tepeh, Extremal bicyclic graphs with respect to Mostar index, *Appl. Math. Comput.* **355**, 319–324 (2019). <https://doi.org/10.1016/j.amc.2019.03.014>
- [11] L. Alex, G. Indulal, On a conjecture on edge Mostar index of bicyclic graphs, *Iran. J. Math. Chem.* **14**, 97–108 (2023).
- [12] L. Alex, I. Gutman, On the inverse Mostar index problem for molecular graphs, *Trans. Comb.* **14**, 65–77 (2025). <https://doi.org/10.22108/toc.2024.139474.2115>
- [13] X. Li, Z. Li, L. Wang, The inverse problems for some topological indices in combinatorial chemistry, *J. Comput. Biol.* **10**, 47–55 (2003).
- [14] S. Wagner, A note on the inverse problem for the Wiener index, *MATCH Commun. Math. Comput. Chem.* **64**, 639–646 (2010).