

Almost complete graphs determined by Laplacian hook immanantal polynomials

Shuaijun Li^a, Guangfu Wang^{b,c} *

^a School of Mathematics and Statistics, Qinghai Minzu University,
Xining, Qinghai 810007, P.R. China

^b School of Mathematical and Information Sciences, Yantai University,
Yantai, Shandong 264000, P.R. China

^c ECOPRO, Institute for Basic Science,
55 Expo-ro, Yuseong-gu, Daejeon, 34126, Korea

Abstract: Let $\mathcal{G}_n$ be the family of simple graphs obtained from K_n by deleting at most five edges. For a fixed integer $1 \leq k \leq n$, let $\Phi_k(L(G), x)$ denote the immanantal polynomial of the Laplacian matrix associated with the hook partition $(k, 1^{n-k})$. We prove that, for $n > 7$ and $n \neq 2k - 1$, every graph in $\mathcal{G}_n$ is determined by $\Phi_k(L(G), x)$ among all simple graphs. We prove that, for $n > 7$ and $n \neq 2k - 1$, every graph in $\mathcal{G}_n$ is determined by $\Phi_k(L(G), x)$ among all simple graphs. The proof recovers the order, size, and degree-square sum from the first coefficients, and then separates the remaining candidates by explicit third- and fourth-coefficient comparisons based on the finite classification of complements with at most five edges. The case $n = 2k - 1$ is left open because the binomial differences used in these comparisons vanish.

Keywords: Laplacian matrix; Hook immanant; Hook immanantal polynomial; The k -th hook immanant partner

MSC: 05C31; 05C30; 15A15

1 Introduction

Let $G = (V(G), E(G))$ be a simple graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$. Let d_{v_i} denote the degree of vertex v_i , and let $D(G) = \text{diag}(d_{v_1}, d_{v_2}, \dots, d_{v_n})$ be the degree diagonal matrix of G . The adjacency matrix of G is an $n \times n$ matrix denoted by $A(G) = [a_{ij}]$, where $a_{ij} = 1$ if vertices v_i and v_j are adjacent, and $a_{ij} = 0$ otherwise. The Laplacian matrix of graph G is defined as $L(G) = D(G) - A(G)$.

Let n be a positive integer, and let S_n denote the symmetric group of order n . For any partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$, let χ_λ stand for the irreducible character of S_n . For any $n \times n$ matrix M , the immanant function of M associated with χ_λ is defined as

$$d_\lambda(M) = \sum_{\sigma \in S_n} \chi_\lambda(\sigma) \prod_{i=1}^n m_{i, \sigma(i)}.$$

*Corresponding author.

E-mail address: gfwang@ytu.edu.cn

When $\lambda = (k, 1^{n-k})$, $d_{(k, 1^{n-k})}(M)$ is called the hook immanant of M . In particular, the determinant $\det M$ and permanent $\text{per } M$ correspond respectively to the alternating character $\chi_{(1^n)}$ and the trivial character $\chi_{(n)}$. Bürgisser [15] proved that the computation of hook immanants is both #P-complete and VNP-complete as the polynomial width grows. As a special graph invariant, hook immanants have profound intrinsic connections with the structural properties of graphs and have been widely studied in algebraic combinatorics [2, 9]. Morris [17] uncovered the inherent relation between the hook immanant of the adjacency matrix and the number of Hamiltonian cycles in a graph. Chan and Lam [13] characterized the minimum bound of the immanants of Laplacian matrices of trees. The theories concerning matrix immanants and the immanantal properties of graph matrices have been systematically elaborated in numerous references; see [6, 7].

Let I_n denote the $n \times n$ identity matrix. The hook immanantal polynomial $\Phi_k(M, x)$ of a matrix M is defined as $d_{(k, 1^{n-k})}(xI_n - M)$, i.e.,

$$\Phi_k(M, x) = d_{(k, 1^{n-k})}(xI_n - M).$$

In particular, suppose M is a graph matrix. When $k = 1$, $\Phi_1(M, x)$ is called the characteristic polynomial of the graph matrix M ; when $k = 2$, $\Phi_2(M, x)$ is referred to as the second immanantal polynomial of M ; when $k = n$, $\Phi_n(M, x)$ is termed the permanent polynomial of M . The polynomial $\Phi_k(L(G), x) = d_{(k, 1^{n-k})}(xI_n - L(G))$ is named the k -th hook immanantal polynomial of graph G . The cases $k = 1, 2, n$ recover the characteristic, second immanantal, and permanent polynomials, respectively. The characteristic polynomials and permanent polynomials of graph matrices have been extensively investigated [22]. Morris [14] pointed out that almost all trees possess a complete set of Laplacian immanantal polynomials. Cash [3] generalized the Sachs theorem for adjacency matrices to immanantal polynomials and studied their structural properties of chemical molecular graphs. Yu and Qu [10] derived an explicit expression for the immanantal polynomial of a graph matrix M via fundamental subgraphs. In algebraic combinatorics, immanants corresponding to several special partitions have been widely researched [5, 26, 27]. For further investigations on immanantal polynomials, readers may refer to [1, 12, 19].

For a fixed k , two n -vertex graphs G and H are called k -th hook-immanantally cospectral if $\Phi_k(L(G), x) = \Phi_k(L(H), x)$. If they are non-isomorphic, they form a pair of k -th hook-immanantal partners. If equality holds for every $1 \leq k \leq n$, we say that G and H have the same complete hook-immanantal polynomial family. A graph G is said to be determined by its k -th hook immanantal polynomial if G has no k -th hook immanantal partner.

A classical open problem in spectral graph theory reads: which graphs are determined by their polynomials? This problem is notoriously difficult within the theory of graph polynomials and has attracted substantial attention from researchers in recent years. Van Dam and Haemers [8, 31] systematically investigated polynomials associated with various graph matrices and remarked that characterizing graphs which are determined by a given graph polynomial constitutes an intriguing research direction. Morris et al. [20] first initiated the study of the question: which graphs are determined by their Laplacian permanent polynomials? Liu [21] proved that both the complete graph K_n and the star graph S_n are uniquely determined by their Laplacian permanent polynomials. Liu et al. [30] verified that a family of unicyclic graphs are uniquely determined by their Laplacian characteristic polynomials. For further relevant results, the reader is referred to [23, 24].

Let $\mathcal{G}_n$ be the family of simple graphs obtained from K_n by deleting at most five edges. Zhang and Wu characterized a graph family $\mathcal{G}_n$ in [11], which consists of all graphs obtained by deleting




Figure 1: $\mathcal{G}_n$: All subgraphs obtained by deleting 1 to 5 edges from the complete graph K_n ; the displayed labels refer to the deleted-edge complements.

at most five edges from the complete graph K_n (see Figure 1). We write $G_{ij} = K_n - E(F_{ij})$, where F_{ij} is the small deleted-edge graph represented by the label G_{ij} in Figure 1. Cámara and Haemers [4] proved that every graph in $\mathcal{G}_n$ is uniquely determined by the characteristic polynomial of its adjacency matrix. Zhang and Wu [11] further showed that all graphs in $\mathcal{G}_n$ are uniquely determined by their Laplacian permanent polynomials. Zeng and Wu verified that graphs in $\mathcal{G}_n$ can be uniquely identified by their second Laplacian immanantal polynomials. Since the characteristic polynomial, permanent polynomial, and second immanantal polynomial are all special cases of hook immanantal polynomials, a natural question arises: are all graphs in $\mathcal{G}_n$ uniquely determined by their k -th hook immanantal polynomials?

We discuss the above question and derive the following theorem:

Theorem 1.1. *For any $1 \leq k \leq n$ and any n -vertex graph $G \in \mathcal{G}_n$, if $n > 7$ and $n \neq 2k - 1$, then G is uniquely determined by its k -th hook immanantal polynomial.*

The remainder of this paper is organized as follows. In Section 2, we derive several special properties of the Laplacian hook immanantal polynomials and present some inequalities for combinatorial sequences. In Section 3, we provide the proof of Theorem 1.1. In Section 4, we summarize the main conclusions of this paper and propose several open problems.

2 Preliminaries

Let G be a simple graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$. Denote by $G - e$ the graph obtained by deleting an edge e from G . Let $\mathfrak{D}(G)$ stand for the degree sequence of G , that is, $\mathfrak{D}(G) = (d(v_1), d(v_2), \dots, d(v_n))$. Let $\mathcal{C}(G)$ and $\mathcal{M}(G)$ denote the sets consisting of all cycles of length l and all l -matchings in G , respectively. Let $\mathcal{T}_j(G)$ stand for the sum of the degrees of the three vertices of the j -th triangle in $\mathcal{C}_3(G)$. If $C = v_{i_1}v_{i_2} \cdots v_{i_l}v_{i_1} \in \mathcal{C}_l(G)$, then

let $\mathfrak{D}_C(G)$ denote the degree sequence obtained by deleting $d(v_{i_1}), d(v_{i_2}), \dots, d(v_{i_l})$ from $\mathfrak{D}(G)$, If $\mathfrak{M} = \{v_{i_1}v_{i'_1}, v_{i_2}v_{i'_2}, \dots, v_{i_l}v_{i'_l}\} \in \mathcal{M}_l(G)$, then let $\mathfrak{D}_{\mathfrak{M}}(G)$ denote the degree sequence obtained by deleting $d(v_{i_1}), d(v_{i'_1}), d(v_{i_2}), d(v_{i'_2}), \dots, d(v_{i_l}), d(v_{i'_l})$ from $\mathfrak{D}(G)$, Let F_r denote the r -th elementary symmetric function, and define $F_r(G) = F_r(\mathfrak{D}(G))$. For any integer $r \geq l$, we define

$$\mathcal{C}_r^l(G) = \sum_{C \in \mathcal{C}_l(G)} F_{r-l}(\mathfrak{D}_C(G)),$$

and

$$\mathcal{M}_r^l(G) = \sum_{\mathfrak{M} \in \mathcal{M}_l(G)} F_{r-2l}(\mathfrak{D}_{\mathfrak{M}}(G)).$$

Lemma 2.1. ([32]) Let G be a graph with n vertices and m edges, and $(d(v_1), d(v_2), \dots, d(v_n))$ denote the degree sequence of G . Denote

$$\Phi_{(k, 1^{n-k})}(xI - L(G)) = \sum_{r=0}^n (-1)^r c_{(k, 1^{n-k}), r}(L(G)) x^{n-r}.$$

Then

$$\begin{aligned} c_{(k, 1^{n-k}), 0}(L(G)) &= \binom{n-1}{k-1}, \\ c_{(k, 1^{n-k}), 1}(L(G)) &= F_1(G) \binom{n-1}{k-1} = 2m \binom{n-1}{k-1}, \\ c_{(k, 1^{n-k}), 2}(L(G)) &= F_2(G) \binom{n-1}{k-1} + m \left[\binom{n-2}{k-2} - \binom{n-2}{k-1} \right] \quad (n \geq 3), \\ c_{(k, 1^{n-k}), 3}(L(G)) &= F_3(G) \binom{n-1}{k-1} + \mathcal{M}_3^1(G) \left[\binom{n-2}{k-2} - \binom{n-2}{k-1} \right] \\ &\quad - 2|\mathcal{C}_3(G)| \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] \quad (n \geq 4), \\ c_{(k, 1^{n-k}), 4}(L(G)) &= F_4(G) \binom{n-1}{k-1} + \mathcal{M}_4^1(G) \left[\binom{n-2}{k-2} - \binom{n-2}{k-1} \right] \\ &\quad - 2\mathcal{C}_4^3(G) \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] + 2|\mathcal{C}_4(G)| \left[\binom{n-5}{k-5} - \binom{n-5}{k-1} \right] \\ &\quad + \left[\binom{m}{2} - \sum_{i=1}^n \binom{d(v_i)}{2} \right] \left[\binom{n-5}{k-5} - 2\binom{n-5}{k-3} + \binom{n-5}{k-1} \right] \quad (n \geq 5), \end{aligned}$$

where $|\mathcal{C}_l(G)|$ denotes the number of cycles of length l in G . Meanwhile, if $b < 0$ or $b > a$, we stipulate that $\binom{a}{b} = 0$.

Lemma 2.2. ([21]) Let G be a simple connected graph with n vertices and m edges, and let $(d_1, d_2, \dots, d_n)$ denote its degree sequence, $S_r(G) = \sum_{i=1}^n d_i^r$. Then:

$$\begin{aligned} F_2(G) &= -\frac{1}{2} \sum_{i=1}^n d_i^2 + 2m^2, \\ F_3(G) &= \frac{1}{3} \sum_{i=1}^n d_i^3 - m \sum_{i=1}^n d_i^2 + \frac{4}{3} m^3, \\ F_4(G) &= -\frac{1}{4} \sum_{i=1}^n d_i^4 + \frac{2}{3} m \sum_{i=1}^n d_i^3 - m^2 \sum_{i=1}^n d_i^2 + \frac{1}{8} \left(\sum_{i=1}^n d_i^2 \right)^2 + \frac{2}{3} m^4, \end{aligned}$$

$$\begin{aligned}
\mathcal{M}_3^1(G) &= 2m^2 - \sum_{i=1}^n d_i^2, \\
\mathcal{M}_4^1(G) &= \sum_{i=1}^n d_i^3 - \frac{5}{2}m \sum_{i=1}^n d_i^2 + \sum_{v_i v_j \in E(G)} d_i d_j + 2m^3, \\
\mathcal{C}_4^3(G) &= 2m|\mathcal{C}_3(G)| - \sum_{C \in \mathcal{C}_3(G)} \sum_{v \in V(C)} d(v) \\
&= 2m|\mathcal{C}_3(G)| - \sum_j \mathcal{T}_j(G).
\end{aligned}$$

Lemma 2.3. *Let $G \in \mathcal{G}_n$, and let H be any simple graph. For a fixed $1 \leq k \leq n$, if $\Phi_k(L(G), x) = \Phi_k(L(H), x)$, then $H \in \mathcal{G}_n$.*

Proof. The degree of the polynomial is the number of vertices, because $c_{k,0} = \binom{n-1}{k-1} > 0$. Thus H has n vertices. Equality of the leading coefficient and of the coefficient of x^{n-1} , together with Lemma 2.1, gives

$$|E(H)| = |E(G)| = \binom{n}{2} - t, \quad 0 \leq t \leq 5.$$

Every simple n -vertex graph with this many edges is obtained from K_n by deleting t edges, so $H \in \mathcal{G}_n$. $\square$

Lemma 2.4. *For any graph G , its hook immanantal polynomial can determine the quantities: n , $m = |E(G)|$ and $\sum_{i=1}^n d_i^2$.*

Proof. The degree determines n . Put $c_0 = \binom{n-1}{k-1}$. Then

$$m = \frac{c_{k,1}(G)}{2c_0}, \quad \sum_{i=1}^n d_i^2 = 4m^2 - \frac{2}{c_0} \left(c_{k,2}(G) - m \left[\binom{n-2}{k-2} - \binom{n-2}{k-1} \right] \right).$$

This formula uses the convention $\binom{a}{b} = 0$ for $b < 0$ or $b > a$, and therefore also covers $k = 1$. $\square$

Lemma 2.5. ([29]) *When $n > 7$, the complete graph with 1, 2, 3, 4 edges removed is determined by its Laplacian characteristic polynomial.*

Lemma 2.6. ([28]) *When $n > 7$, the complete graph with 5 edges removed is determined by its Laplacian characteristic polynomial.*

Lemma 2.7. ([33]) *All graphs in $\mathcal{G}_n$ are determined by their second immanantal polynomial.*

Wu and Zhou [25] computed the sum of squares of degrees for graphs in $\mathcal{G}_n$ (see Table 1). For the convenience of subsequent proofs, we further calculated the sum of cubes of degrees, the sum of the products of the degrees of vertices on every cycle of length 3, and the sum of the products of the degrees of the two endpoints over all edges for several graphs in $\mathcal{G}_n$, as listed in Table 2 and Table 3 respectively.

Lemma 2.8. ([16]) *Let n be a positive integer. Then the binomial coefficient sequence*

$$\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n}$$

is unimodal.

Table 1: $\sum_{i=1}^n d_i^2$: The sum of squares of degrees in $\mathcal{G}_n$ (Source: [25])

Graph	$\sum_{i=1}^n (d_i)^2$	Graph	$\sum_{i=1}^n (d_i)^2$
G_{20}	$n^3 - 2n^2 - 7n + 14$	G_{30}, G_{32}	$n^3 - 2n^2 - 11n + 24$
G_{33}	$n^3 - 2n^2 - 11n + 22$	G_{40}	$n^3 - 2n^2 - 15n + 36$
G_{42}, G_{48}	$n^3 - 2n^2 - 15n + 28$	G_{44}, G_{410}	$n^3 - 2n^2 - 15n + 32$
G_{49}	$n^3 - 2n^2 - 15n + 24$	G_{52}	$n^3 - 2n^2 - 19n + 32$
G_{54}	$n^3 - 2n^2 - 19n + 50$	$G_{56}, G_{512}, G_{515}, G_{525}$	$n^3 - 2n^2 - 19n + 42$
$G_{58}, G_{511}, G_{514}, G_{517}$	$n^3 - 2n^2 - 19n + 36$	G_{510}, G_{524}	$n^3 - 2n^2 - 19n + 44$
G_{21}	$n^3 - 2n^2 - 7n + 12$	G_{31}	$n^3 - 2n^2 - 11n + 20$
G_{34}	$n^3 - 2n^2 - 11n + 18$	G_{41}, G_{45}, G_{47}	$n^3 - 2n^2 - 15n + 30$
G_{43}	$n^3 - 2n^2 - 15n + 26$	G_{46}	$n^3 - 2n^2 - 15n + 34$
G_{50}, G_{51}	$n^3 - 2n^2 - 19n + 34$	G_{59}, G_{516}	$n^3 - 2n^2 - 19n + 46$
$G_{55}, G_{518}, G_{520}, G_{523}$	$n^3 - 2n^2 - 19n + 40$	G_{521}	$n^3 - 2n^2 - 19n + 30$
$G_{53}, G_{57}, G_{513}, G_{519}, G_{522}$	$n^3 - 2n^2 - 19n + 38$		

Lemma 2.9. *Let n, k be two positive integers with $k \geq 5$. Then*

$$\begin{cases} \binom{n-4}{k-4} - \binom{n-4}{k-1} < 0, \binom{n-5}{k-5} - \binom{n-5}{k-1} < 0 & n \geq 2k, \\ \binom{n-4}{k-4} - \binom{n-4}{k-1} = 0, \binom{n-5}{k-5} - \binom{n-5}{k-1} = 0 & n = 2k - 1, \\ \binom{n-4}{k-4} - \binom{n-4}{k-1} > 0, \binom{n-5}{k-5} - \binom{n-5}{k-1} > 0 & n \leq 2k - 2. \end{cases}$$

Proof. Let $A = \binom{n-4}{k-4} - \binom{n-4}{k-1}$, $B = \binom{n-5}{k-5} - \binom{n-5}{k-1}$. The case $n = 2k - 1$ is straightforward to verify. By Lemma 2.7, the binomial coefficient sequence is unimodal. If $k - 1 \leq \frac{n-4}{2}$, i.e., $n \geq 2k + 2$, then $A < 0$. If $k - 4 \geq \frac{n-4}{2}$, i.e., $n \leq 2k - 4$, then $A > 0$. When $n = 2k - 3$: $A = \binom{2k-7}{k-4} - \binom{2k-7}{k-1} > 0$. When $n = 2k - 2$: $A = \binom{2k-4}{k-4} - \binom{2k-4}{k-1} > 0$. When $n = 2k$: $A = \binom{2k-4}{k-4} - \binom{2k-4}{k-1} < 0$. When $n = 2k + 1$: $A = \binom{2k-3}{k-4} - \binom{2k-3}{k-1} < 0$. To summarize: $A < 0$ whenever $n \geq 2k$, and $A > 0$ whenever $n \leq 2k - 2$. Similarly, we deduce: $B < 0$ for $n \geq 2k$, $B = 0$ for $n = 2k - 1$, $B > 0$ for $n \leq 2k - 2$. $\square$

Corollary 2.1. *For any positive integers a, b, c, n, k with $n \geq k$, $k \geq 5$ and $n \neq 2k - 1$, we have*

$$\frac{a(2k - n - 1)}{k - 1} \binom{n-2}{k-2} + b \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] + c \left[\binom{n-5}{k-5} - \binom{n-5}{k-1} \right] \neq 0.$$

Proof. If $n \geq 2k$, then $\frac{2k-n-1}{k-1} < 0$, $\binom{n-4}{k-4} - \binom{n-4}{k-1} < 0$, $\binom{n-5}{k-5} - \binom{n-5}{k-1} < 0$, so $\frac{a(2k-n-1)}{k-1} \binom{n-2}{k-2} + b \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] + c \left[\binom{n-5}{k-5} - \binom{n-5}{k-1} \right] < 0$. If $n \leq 2k - 2$, then $\frac{2k-n-1}{k-1} > 0$, $\binom{n-4}{k-4} - \binom{n-4}{k-1} > 0$, $\binom{n-5}{k-5} - \binom{n-5}{k-1} > 0$, so $\frac{a(2k-n-1)}{k-1} \binom{n-2}{k-2} + b \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] + c \left[\binom{n-5}{k-5} - \binom{n-5}{k-1} \right] > 0$. Therefore, $\frac{a(2k-n-1)}{k-1} \binom{n-2}{k-2} + b \left[\binom{n-4}{k-4} - \binom{n-4}{k-1} \right] + c \left[\binom{n-5}{k-5} - \binom{n-5}{k-1} \right] \neq 0$. $\square$

Lemma 2.10. *Let n, k be two positive integers with $n \geq k$ and $n \neq k - 1$. Then*

$$\binom{n-1}{k-1} \geq \binom{n-4}{k-4}.$$

Proof. The cases $k = 1, 2, 3$ are trivial to verify.

Table 2: $\sum_{i=1}^n (d_i)^3$: The sum of cubes of degrees in $\mathcal{G}_n$

Graph	$\sum_{i=1}^n (d_i)^3$	Graph	$\sum_{i=1}^n (d_i)^3$
G_{30}	$n^4 - 3n^3 - 15n^2 + 71n - 84$	$G_{511}, G_{514}, G_{517}$	$n^4 - 3n^3 - 27n^2 + 107n - 106$
G_{32}	$n^4 - 3n^3 - 15n^2 + 71n - 78$	G_{512}, G_{515}	$n^4 - 3n^3 - 27n^2 + 125n - 148$
G_{41}	$n^4 - 3n^3 - 21n^2 + 89n - 98$	$G_{513}, G_{519}, G_{522}$	$n^4 - 3n^3 - 27n^2 + 113n - 118$
G_{42}, G_{48}	$n^4 - 3n^3 - 21n^2 + 83n - 80$	G_{516}	$n^4 - 3n^3 - 27n^2 + 137n - 178$
G_{44}	$n^4 - 3n^3 - 21n^2 + 95n - 104$	G_{518}, G_{523}, G_{55}	$n^4 - 3n^3 - 27n^2 + 119n - 136$
G_{45}, G_{47}	$n^4 - 3n^3 - 21n^2 + 89n - 92$	G_{520}	$n^4 - 3n^3 - 27n^2 + 119n - 130$
G_{410}	$n^4 - 3n^3 - 21n^2 + 95n - 110$	G_{524}	$n^4 - 3n^3 - 27n^2 + 131n - 178$
G_{50}, G_{51}	$n^4 - 3n^3 - 27n^2 + 101n - 94$	G_{525}	$n^4 - 3n^3 - 27n^2 + 125n - 154$
G_{53}, G_{57}	$n^4 - 3n^3 - 27n^2 + 113n - 124$	G_{59}	$n^4 - 3n^3 - 27n^2 + 137n - 190$
G_{56}	$n^4 - 3n^3 - 27n^2 + 125n - 166$	G_{510}	$n^4 - 3n^3 - 27n^2 + 131n - 166$
G_{58}	$n^4 - 3n^3 - 27n^2 + 107n - 112$		

For $k \geq 4$, direct computation yields:

$$\binom{n-1}{k-1} = \frac{(n-1)!}{(k-1)!(n-k)!}, \quad \binom{n-4}{k-4} = \frac{(n-4)!}{(k-4)!(n-k)!}$$

Thus

$$\frac{\binom{n-1}{k-1}}{\binom{n-4}{k-4}} = \frac{(n-1)(n-2)(n-3) \cdot (n-4)!}{(k-1)(k-2)(k-3) \cdot (k-4)!} \cdot \frac{(k-4)!}{(n-4)!} = \frac{(n-1)(n-2)(n-3)}{(k-1)(k-2)(k-3)}$$

Since $n \geq k \geq 4$ and n, k are positive integers, we have

$$n-1 \geq k-1 > 0, \quad n-2 \geq k-2 > 0, \quad n-3 \geq k-3 > 0$$

Hence $\frac{(n-1)(n-2)(n-3)}{(k-1)(k-2)(k-3)} \geq 1$.

To sum up, $\binom{n-1}{k-1} \geq \binom{n-4}{k-4}$.

□

Lemma 2.11. *Let n, k be two positive integers with $n \geq k$ and $n \neq k-1$. Then*

$$\binom{n-1}{k-1} \geq \binom{n-4}{k-1}.$$

Proof. If $n \leq 4$ or $n < k+3$, then $\binom{n-4}{k-1} = 0$, and $\binom{n-1}{k-1} \geq 0$. Hence $\binom{n-1}{k-1} \geq \binom{n-4}{k-1}$.

For $n \geq k+3$ and $n > 4$, expand via the definition of binomial coefficients:

$$\binom{n-1}{k-1} = \frac{(n-1)!}{(k-1)!(n-k)!}, \quad \binom{n-4}{k-1} = \frac{(n-4)!}{(k-1)!(n-k-3)!}$$

take the ratio of the two expressions:

$$\begin{aligned} \frac{\binom{n-1}{k-1}}{\binom{n-4}{k-1}} &= \frac{(n-1)(n-2)(n-3) \cdot (n-4)!}{(n-k)(n-k-1)(n-k-2)(n-k-3)!} \cdot \frac{(n-k-3)!}{(n-4)!} \\ &= \frac{(n-1)(n-2)(n-3)}{(n-k)(n-k-1)(n-k-2)} \end{aligned}$$

Table 3: The sum of the products of the degrees of all vertices of cycles of length 3 and the sum of the products of the degrees of all edges in $\mathcal{G}_n$

Graph	$\sum_i \mathcal{T}_j(G)$	$\sum_{v_i v_j \in E(G)} d_{v_i} d_{v_j}$
G_{42}	$(1/2)n^4 - 2n^3 - (27/2)n^2 + 65n - 50$	$(1/2)n^4 - (3/2)n^3 - (21/2)n^2 + (71/2)n - 6$
G_{50}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 79n - 48$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (87/2)n + 4$
G_{53}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 89n - 79$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (95/2)n - 7$
G_{55}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 94n - 94$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (99/2)n - 13$
G_{57}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 89n - 75$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (95/2)n - 5$
G_{519}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 89n - 80$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (95/2)n - 9$
G_{511}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 84n - 61$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (91/2)n - 1$
G_{48}	$(1/2)n^4 - 2n^3 - (27/2)n^2 + 65n - 52$	$(1/2)n^4 - (3/2)n^3 - (21/2)n^2 + (71/2)n - 7$
G_{51}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 79n - 46$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (87/2)n + 5$
G_{522}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 89n - 78$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (95/2)n - 8$
G_{517}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 84n - 63$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (91/2)n - 2$
G_{523}	$(1/2)n^4 - 2n^3 - (35/2)n^2 + 94n - 96$	$(1/2)n^4 - (3/2)n^3 - (27/2)n^2 + (99/2)n - 14$

Since $n \geq k$ and n, k are positive integers, we have $\frac{(n-1)(n-2)(n-3)}{(n-k)(n-k-1)(n-k-2)} \geq 1$. Therefore, $\binom{n-1}{k-1} \geq \binom{n-4}{k-1}$.

□

3 Proof of Theorem 1.1

From Lemmas 2.4, 2.5 and 2.6, we know that Theorem 1.1 holds for any hook partition $\lambda = (k, 1^{n-k})$ with $k = 1, 2$. Therefore, to prove Theorem 1.1, it suffices to treat the case $k \geq 3$. We outline the main proof idea as follows. According to Table 3, the graph class $\mathcal{G}_n$ is partitioned into distinct groups by the sum of squares of degrees of its graphs. By Lemmas 2.1 and 2.3, any two graphs from different groups are determined by their hook immanantal polynomials, since they differ either in the number of edges or in the sum of squares of degrees. Furthermore, combining Lemmas 2.7, 2.8, Corollary 2.1 and Tables 1, 2, 3, 4, 5, we compute the 3rd and 4th coefficients of the hook immanantal polynomial for each graph in every group under different hook partitions $\lambda = (k, 1^{n-k})$. We then judge whether these graphs are determined by the k -hook immanantal polynomial via coefficient comparison. Zhang and Wu [11] presented the numbers of triangles (Table 4) and quadrilaterals (Table 5) for graphs in $\mathcal{G}_n$, which greatly facilitates the computation of the aforementioned coefficients. Consequently, proving the following lemma is sufficient to establish Theorem 1.1.

Lemma 3.1. *For any partition $\lambda = (k, 1^{n-k})$ with $k > 2$ and $n \neq 2k - 1$, the complete graph K_n and the graphs $G_{10}, G_{20}, G_{21}, G_{31}, G_{33}, G_{34}, G_{40}, G_{43}, G_{46}, G_{49}, G_{52}, G_{54}, G_{521}$ are each uniquely determined by their k -th hook immanantal polynomial.*

Proof. For the graphs listed above, any two distinct graphs differ either in the number of vertices, the number of edges, or the sum of squares of degrees. It is therefore straightforward to verify that all these graphs are uniquely determined by their k -th hook immanantal polynomial. □

Table 4: The number of $c_3(G)$ in $\mathcal{G}_n$

Graph	$\mathcal{C}_3(G)$	Graph	$\mathcal{C}_3(G)$
G_{10}	$\binom{n}{3} - n + 2$	G_{521}	$\binom{n}{3} - 5n + 10$
G_{20}	$\binom{n}{3} - 2n + 5$	G_{32}, G_{33}	$\binom{n}{3} - 3n + 8$
G_{21}	$\binom{n}{3} - 2n + 4$	G_{41}, G_{47}	$\binom{n}{3} - 4n + 11$
G_{30}	$\binom{n}{3} - 3n + 9$	G_{59}, G_{524}	$\binom{n}{3} - 5n + 17$
G_{31}	$\binom{n}{3} - 3n + 7$	G_{42}, G_{45}, G_{48}	$\binom{n}{3} - 4n + 10$
G_{34}	$\binom{n}{3} - 3n + 6$	G_{44}, G_{46}, G_{410}	$\binom{n}{3} - 4n + 12$
G_{40}	$\binom{n}{3} - 4n + 14$	G_{50}, G_{51}, G_{514}	$\binom{n}{3} - 5n + 12$
G_{43}	$\binom{n}{3} - 4n + 9$	$G_{55}, G_{512}, G_{520}, G_{523}$	$\binom{n}{3} - 5n + 15$
G_{49}	$\binom{n}{3} - 4n + 8$	$G_{58}, G_{511}, G_{513}, G_{517}$	$\binom{n}{3} - 5n + 13$
G_{52}	$\binom{n}{3} - 5n + 11$	$G_{53}, G_{57}, G_{518}, G_{519}, G_{522}$	$\binom{n}{3} - 5n + 14$
G_{54}	$\binom{n}{3} - 5n + 20$	$G_{56}, G_{510}, G_{515}, G_{516}, G_{525}$	$\binom{n}{3} - 5n + 16$

Lemma 3.2. *Let G be a simple connected graph with n vertices ($n > 7$) and m edges. For any partition $\lambda = (k, 1^{n-k})$, if $n \geq k > 2$ and $n \neq 2k - 1$, then the graphs G_{30} and G_{32} are determined by their k -th hook immanantal polynomial.*

Proof. From Table 4, Lemma 2.1 and Lemma 2.10, for graphs G_{30} and G_{32} , when $k = 3$, we have $c_{k,3}(G_{30}) - c_{k,3}(G_{32}) = (F_3(G_{30}) - F_3(G_{32}))\binom{n-1}{k-1} + \frac{(2k-n-1)(\mathcal{M}_3^1(G_{30}) - \mathcal{M}_3^1(G_{32}))}{k-1}\binom{n-2}{k-2} - 2(|\mathcal{C}_3(G_{30})| - |\mathcal{C}_3(G_{32})|)\binom{n-4}{k-1} = -2\binom{n-1}{2} + 2\binom{n-4}{2} \neq 0$. When $k \geq 4$, $c_{k,3}(G_{30}) - c_{k,3}(G_{32}) = (F_3(G_{30}) - F_3(G_{32}))\binom{n-1}{k-1} + \frac{(2k-n-1)(\mathcal{M}_3^1(G_{30}) - \mathcal{M}_3^1(G_{32}))}{k-1}\binom{n-2}{k-2} - 2(|\mathcal{C}_3(G_{30})| - |\mathcal{C}_3(G_{32})|)\binom{n-4}{k-4} - \binom{n-4}{k-1} = -2\binom{n-1}{k-1} - 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \leq -2\binom{n-4}{k-4} < 0$. Therefore, the graphs G_{30} and G_{32} are determined by their k -th hook immanantal polynomial. $\square$

Table 5: The number of $c_4(G)$ in $\mathcal{G}_n$

Graph	$\mathcal{C}_4(G)$	Graph	$\mathcal{C}_4(G)$
G_{32}	$3\binom{n}{4} - 3n^2 + 18n - 27$	G_{510}	$3\binom{n}{4} - 5n^2 + 32n - 50$
G_{33}	$3\binom{n}{4} - 3n^2 + 17n - 23$	G_{511}	$3\binom{n}{4} - 5n^2 + 28n - 26$
G_{42}	$3\binom{n}{4} - 4n^2 + 22n - 22$	G_{515}	$3\binom{n}{4} - 5n^2 + 31n - 45$
G_{44}	$3\binom{n}{4} - 4n^2 + 24n - 35$	G_{516}	$3\binom{n}{4} - 5n^2 + 33n - 55$
G_{46}	$3\binom{n}{4} - 4n^2 + 25n - 39$	G_{517}	$3\binom{n}{4} - 5n^2 + 28n - 27$
G_{47}	$3\binom{n}{4} - 4n^2 + 23n - 29$	G_{518}	$3\binom{n}{4} - 5n^2 + 30n - 37$
G_{48}	$3\binom{n}{4} - 4n^2 + 22n - 23$	G_{520}	$3\binom{n}{4} - 5n^2 + 30n - 40$
G_{410}	$3\binom{n}{4} - 4n^2 + 24n - 34$	G_{523}	$3\binom{n}{4} - 5n^2 + 30n - 39$
G_{50}	$3\binom{n}{4} - 5n^2 + 27n - 21$	G_{524}	$3\binom{n}{4} - 5n^2 + 32n - 48$
G_{51}	$3\binom{n}{4} - 5n^2 + 27n - 20$	G_{41}, G_{45}	$3\binom{n}{4} - 4n^2 + 23n - 27$
G_{53}	$3\binom{n}{4} - 5n^2 + 29n - 32$	G_{57}, G_{513}	$3\binom{n}{4} - 5n^2 + 29n - 30$
G_{55}	$3\binom{n}{4} - 5n^2 + 30n - 38$	G_{58}, G_{514}	$3\binom{n}{4} - 5n^2 + 28n - 25$
G_{56}	$3\binom{n}{4} - 5n^2 + 31n - 40$	G_{512}, G_{525}	$3\binom{n}{4} - 5n^2 + 31n - 44$
G_{59}	$3\binom{n}{4} - 5n^2 + 33n - 54$	G_{519}, G_{522}	$3\binom{n}{4} - 5n^2 + 29n - 33$

Lemma 3.3. *Let G be a simple connected graph with $n > 7$ vertices and m edges. For any partition $\lambda = (k, 1^{n-k})$, if $k > 2$ and $n \neq 2k - 1$, each of the following statements holds:*

- (i) The graphs G_{42} and G_{48} are determined by their k -th hook immanantal polynomial.
- (ii) The graphs G_{44} and G_{410} are determined by their k -th hook immanantal polynomial.
- (iii) The graphs G_{41} , G_{45} and G_{47} are determined by their k -th hook immanantal polynomial.

Proof. (i) From Tables 4, 5, Lemma 2.1 and Corollary 2.1, for graphs G_{42} and G_{48} with $n > 7$:
When $k = 3$, $c_{k,4}(G_{42}) - c_{k,4}(G_{48}) = [F_4(G_{42}) - F_4(G_{48})] \binom{n-1}{2} + \frac{(5-n)(\mathcal{M}_4^1(G_{42}) - \mathcal{M}_4^1(G_{48}))}{2} \binom{n-2}{1} - 2(\mathcal{C}_4^3(G_{42}) - \mathcal{C}_4^3(G_{48}))[-\binom{n-4}{2}] + 2(|\mathcal{C}_4(G_{42})| - |\mathcal{C}_4(G_{48})|)[- \binom{n-5}{2}] + [\binom{m}{2} - \sum_{i=1}^n ((d_{42}^{(v_i)}) - (d_{48}^{(v_i)}))][- 2 \binom{n-5}{k-1}] = \frac{(5-n)}{2} \binom{n-2}{k-2} - 4 \binom{n-4}{2} - 2 \binom{n-5}{2} < 0$, When $k = 4$, $c_{k,4}(G_{42}) - c_{k,4}(G_{48}) = \frac{(7-n)}{3} \binom{n-2}{2} + 4[1 - \binom{n-4}{3}] - 2 \binom{n-5}{k-1} = \frac{-(n-7)(7n^2-59n+138)}{6}$, This equation has no positive integer solutions for $n > 7$. When $k \geq 5$, $c_{k,4}(G_{42}) - c_{k,4}(G_{48}) = [F_4(G_{42}) - F_4(G_{48})] \binom{n-1}{k-1} + \frac{(2k-n-1)(\mathcal{M}_4^1(G_{42}) - \mathcal{M}_4^1(G_{48}))}{k-1} \binom{n-2}{k-2} - 2(\mathcal{C}_4^3(G_{42}) - \mathcal{C}_4^3(G_{48}))[\binom{n-4}{k-4} - \binom{n-4}{k-1}] + 2(|\mathcal{C}_4(G_{42})| - |\mathcal{C}_4(G_{48})|)[\binom{n-5}{k-5} - \binom{n-5}{k-1}] + [\binom{m}{2} - \sum_{i=1}^n ((d_{42}^{(v_i)}) - (d_{48}^{(v_i)}))][\binom{n-5}{k-5} - 2 \binom{n-5}{k-3} + \binom{n-5}{k-1}] = \frac{(2k-n-1)}{k-1} \binom{n-2}{k-2} + 4[\binom{n-4}{k-4} - \binom{n-4}{k-1}] + 2[\binom{n-5}{k-5} - \binom{n-5}{k-1}] \neq 0$, Therefore, the graphs G_{42} and G_{48} are determined by their k -th hook immanantal polynomial.

(ii) From Table 4 and Lemma 2.1, for graphs G_{44} and G_{410} , $c_{k,3}(G_{44}) - c_{k,3}(G_{410}) = (F_3(G_{44}) - F_3(G_{410})) \binom{n-1}{k-1} + \frac{(2k-n-1)(\mathcal{M}_3^1(G_{44}) - \mathcal{M}_3^1(G_{410}))}{k-1} \binom{n-2}{k-2} - 2(|\mathcal{C}_3(G_{44})| - |\mathcal{C}_3(G_{410})|)[\binom{n-4}{k-4} - \binom{n-4}{k-1}] = 2 \binom{n-1}{k-1} \neq 0$, Therefore, the graphs G_{44} and G_{410} are determined by their k -th hook immanantal polynomial.

(iii) From Table 4, Lemma 2.1 and Lemma 2.10, for graphs G_{41} and G_{45} : When $k = 3$, $c_{k,3}(G_{41}) - c_{k,3}(G_{45}) = -2 \binom{n-1}{2} + 2 \binom{n-4}{2} \neq 0$, When $k \geq 4$, $c_{k,3}(G_{41}) - c_{k,3}(G_{45}) = -2 \binom{n-1}{k-1} - 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \leq -2 \binom{n-4}{k-4} < 0$, Thus, the graphs G_{41} and G_{45} are determined by their k -th hook immanantal polynomial. Meanwhile, for graphs G_{41} and G_{47} , $c_{k,3}(G_{41}) - c_{k,3}(G_{47}) = -2 \binom{n-1}{k-1} \neq 0$, G_{41} and G_{47} are determined by their k -th hook immanantal polynomial. Similarly, for graphs G_{45} and G_{47} , $c_{k,3}(G_{45}) - c_{k,3}(G_{47}) = 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}]$, Since $n \neq 2k-1$, we have $2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \neq 0$, Therefore, the graphs G_{45} and G_{47} are determined by their k -th hook immanantal polynomial. □

Lemma 3.4. Let G be a simple connected graph with $n > 7$ vertices and m edges. For any partition $\lambda = (k, 1^{n-k})$, if $k > 2$ and $n \neq 2k-1$, each of the following statements holds:

- (i) The graphs G_{56} , G_{512} , G_{515} and G_{525} are determined by their k -th hook immanantal polynomial.
- (ii) The graphs G_{58} , G_{511} , G_{514} and G_{517} are determined by their k -th hook immanantal polynomial.
- (iii) The graphs G_{510} and G_{524} are determined by their k -th hook immanantal polynomial.
- (iv) The graphs G_{53} , G_{57} , G_{513} , G_{519} and G_{522} are determined by their k -th hook immanantal polynomial.
- (v) The graphs G_{55} , G_{518} , G_{520} and G_{523} are determined by their k -th hook immanantal polynomial.
- (vi) The graphs G_{59} and G_{516} are determined by their k -th hook immanantal polynomial.
- (vii) The graphs G_{50} and G_{51} are determined by their k -th hook immanantal polynomial.

Proof. (i) From Table 4, Lemma 2.1 and Lemma 2.10, for graphs G_{56} and G_{512} : When $k = 3$, $c_{k,3}(G_{56}) - c_{k,3}(G_{512}) = -6 \binom{n-1}{2} + 2 \binom{n-4}{2} \leq -4 \binom{n-1}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{56}) -$

$c_{k,3}(G_{512}) = -6\binom{n-1}{k-1} - 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \leq -4\binom{n-1}{k-1} - 2\binom{n-4}{k-4} < 0$, Thus, G_{56} and G_{512} are determined by their k -th hook immanantal polynomial. Similarly, $c_{k,3}(G_{56}) - c_{k,3}(G_{515}) = -6\binom{n-1}{k-1} \neq 0$, $c_{k,3}(G_{56}) - c_{k,3}(G_{525}) = -4\binom{n-1}{k-1} \neq 0$, so G_{56} is determined from both G_{515} and G_{525} by the k -th hook immanantal polynomial. For the previously missing pair G_{512}, G_{515} , the first differing coefficient is

$$c_{k,3}(G_{512}) - c_{k,3}(G_{515}) = \begin{cases} -2\binom{n-4}{2} < 0, & k = 3, \\ 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \neq 0, & k \geq 4, \end{cases}$$

where the second line is nonzero because $n \neq 2k - 1$. For the pair G_{512}, G_{525} , the difference is

$$c_{k,3}(G_{512}) - c_{k,3}(G_{525}) = \begin{cases} 2\binom{n-1}{2} - 2\binom{n-4}{2} \neq 0, & k = 3, \\ 2\binom{n-1}{k-1} + 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] > 0, & k \geq 4, \end{cases}$$

because $\binom{n-1}{k-1} \geq \binom{n-4}{k-1}$. Thus all six unordered pairs in this four-graph group have a nonzero coefficient certificate. For graphs G_{515} and G_{525} , $c_{k,3}(G_{515}) - c_{k,3}(G_{525}) = 2\binom{n-1}{k-1} \neq 0$, so G_{515} and G_{525} are determined by their k -th hook immanantal polynomial.

- (ii) From Tables 4, 5, Lemma 2.1, Corollary 2.1 and Lemma 2.10: For graphs G_{58} and G_{511} , $c_{k,3}(G_{58}) - c_{k,3}(G_{511}) = -2\binom{n-1}{k-1} \neq 0$. For graphs G_{58} and G_{514} : When $k = 3$, $c_{k,3}(G_{58}) - c_{k,3}(G_{514}) = 2\binom{n-1}{2} - 2\binom{n-4}{2} \neq 0$, When $k \geq 4$, $c_{k,3}(G_{58}) - c_{k,3}(G_{514}) = -2\binom{n-1}{k-1} - 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \leq -2\binom{n-4}{k-4} < 0$. For graphs G_{58} and G_{517} , $c_{k,3}(G_{58}) - c_{k,3}(G_{517}) = -2\binom{n-1}{k-1} \neq 0$. For graphs G_{511} and G_{514} : When $k = 3$, $c_{k,3}(G_{511}) - c_{k,3}(G_{514}) = 2\binom{n-4}{2} > 0$, When $k \geq 4$, $c_{k,3}(G_{511}) - c_{k,3}(G_{514}) = -2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right]$, Since $n \neq 2k - 1$, we have $-2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \neq 0$. For graphs G_{511} and G_{517} with $n > 7$: When $k = 3$, $c_{k,4}(G_{511}) - c_{k,4}(G_{517}) = \frac{(5-n)}{2}\binom{n-2}{1} - 4\binom{n-4}{2} - 2\binom{n-5}{2} < 0$, When $k = 4$, $c_{k,4}(G_{511}) - c_{k,4}(G_{517}) = \frac{(7-n)}{3}\binom{n-2}{2} + 4\left[1 - \binom{n-4}{3}\right] - 2\binom{n-5}{3} = \frac{-(n-7)(7n^2-59n+138)}{6}$, which has no positive integer solutions for $n > 7$. When $k \geq 5$, $c_{k,4}(G_{511}) - c_{k,4}(G_{517}) = \frac{(2k-n-1)}{k-1}\binom{n-2}{k-2} + 4\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] + 2\left[\binom{n-5}{k-5} - \binom{n-5}{k-1}\right] \neq 0$. For graphs G_{514} and G_{517} : When $k = 3$, $c_{k,3}(G_{514}) - c_{k,3}(G_{517}) = -2\binom{n-4}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{514}) - c_{k,3}(G_{517}) = 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right]$, Since $n \neq 2k - 1$, we have $2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \neq 0$. Thus every graph in the set $\{G_{58}, G_{511}, G_{514}, G_{517}\}$ is determined by its k -th hook immanantal polynomial.
- (iii) From Table 4, Lemma 2.1 and Lemma 2.10, for graphs G_{510} and G_{524} : When $k = 3$, $c_{k,3}(G_{510}) - c_{k,3}(G_{524}) = 4\binom{n-1}{2} - 2\binom{n-4}{2} \geq 2\binom{n-1}{2} > 0$, When $k \geq 4$, $c_{k,3}(G_{510}) - c_{k,3}(G_{524}) = 4\binom{n-1}{k-1} + 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \geq 2\binom{n-1}{k-1} + 2\binom{n-4}{k-4} > 0$. Therefore, the graphs G_{510} and G_{524} are determined by their k -th hook immanantal polynomial.
- (iv) From Tables 4, 5, Lemma 2.1, Corollary 2.1 and Lemma 2.10: For graphs G_{53} and G_{57} with $n > 7$: When $k = 3$, $c_{k,4}(G_{53}) - c_{k,4}(G_{57}) = -(5-n)\binom{n-2}{1} + 8\binom{n-4}{2} + 4\binom{n-5}{2} > 0$, When $k = 4$, $c_{k,4}(G_{53}) - c_{k,4}(G_{57}) = -\frac{2(7-n)}{3}\binom{n-2}{2} - 8\left[1 - \binom{n-4}{3}\right] + 4\binom{n-5}{3} = \frac{-(n-7)(7n^2-59n+138)}{3}$, which has no integer solutions making the difference zero for all $n > 7$. When $k \geq 5$, $c_{k,4}(G_{53}) - c_{k,4}(G_{57}) = -2\frac{(2k-n-1)}{k-1}\binom{n-2}{k-2} - 8\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] - 4\left[\binom{n-5}{k-5} - \binom{n-5}{k-1}\right] \neq 0$, Thus G_{53} and G_{57} are determined by their k -th hook immanantal polynomial. For graphs G_{53} and G_{513} : When $k = 3$, $c_{k,3}(G_{53}) - c_{k,3}(G_{513}) = -2\binom{n-1}{2} + 2\binom{n-4}{2} \neq 0$, When $k \geq 4$, $c_{k,3}(G_{53}) - c_{k,3}(G_{513}) = -2\binom{n-1}{k-1} - 2\left[\binom{n-4}{k-4} - \binom{n-4}{k-1}\right] \leq -2\binom{n-4}{k-4} < 0$, Hence G_{53} and G_{513} are determined by their k -th hook immanantal polynomial. For graphs G_{53} and G_{519} , $c_{k,3}(G_{53}) - c_{k,3}(G_{519}) = -2\binom{n-1}{k-1} \neq 0$, so G_{53} and G_{519} are determined by their k -th hook immanantal polynomial. For graphs G_{53} and G_{522} , $c_{k,3}(G_{53}) - c_{k,3}(G_{522}) = -2\binom{n-1}{k-1} \neq 0$, so G_{53} and G_{522} are

determined by their k -th hook immanantal polynomial. For graphs G_{57} and G_{513} : When $k = 3$, $c_{k,3}(G_{57}) - c_{k,3}(G_{513}) = -2\binom{n-1}{2} + 2\binom{n-4}{2} \neq 0$, When $k \geq 4$, $c_{k,3}(G_{57}) - c_{k,3}(G_{513}) = -2\binom{n-1}{k-1} - 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \leq -2\binom{n-4}{k-4} < 0$, Thus G_{57} and G_{513} are determined by their k -th hook immanantal polynomial. For graphs G_{57} and G_{519} , $c_{k,3}(G_{57}) - c_{k,3}(G_{519}) = -2\binom{n-1}{k-1} < 0$, so G_{57} and G_{519} are determined by their k -th hook immanantal polynomial. For graphs G_{57} and G_{522} , $c_{k,3}(G_{57}) - c_{k,3}(G_{522}) = -2\binom{n-1}{k-1} \neq 0$, so G_{57} and G_{522} are determined by their k -th hook immanantal polynomial. For graphs G_{513} and G_{519} : When $k = 3$, $c_{k,3}(G_{513}) - c_{k,3}(G_{519}) = -2\binom{n-4}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{513}) - c_{k,3}(G_{519}) = 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}]$, Since $n \neq 2k - 1$, we obtain $2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \neq 0$, so the graphs G_{513} and G_{519} are determined by their k -th hook immanantal polynomial. For graphs G_{513} and G_{522} : When $k = 3$, $c_{k,3}(G_{513}) - c_{k,3}(G_{522}) = -2\binom{n-4}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{513}) - c_{k,3}(G_{522}) = 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}]$, As $n \neq 2k - 1$, we have $2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \neq 0$, thus the graphs G_{513} and G_{522} are determined by their k -th hook immanantal polynomial. For graphs G_{519} and G_{522} with $n > 7$: When $k = 3$, $c_{k,4}(G_{519}) - c_{k,4}(G_{522}) = -\frac{(5-n)}{2}\binom{n-2}{1} + 4\binom{n-5}{2} > 0$, When $k = 4$, $c_{k,4}(G_{519}) - c_{k,4}(G_{522}) = -\frac{(7-n)}{3}\binom{n-2}{2} + 4\binom{n-5}{3} = \frac{(n-7)(5n^2-49n+126)}{6}$, If this difference equals zero, the above equation admits no positive integer solutions satisfying $n > 7$. When $k \geq 5$, $c_{k,4}(G_{519}) - c_{k,4}(G_{522}) = -\frac{(2k-n-1)}{k-1}\binom{n-2}{k-2} - 4[\binom{n-5}{k-5} - \binom{n-5}{k-1}] \neq 0$, Therefore, the graphs G_{519} and G_{522} are determined by their k -th hook immanantal polynomial.

- (v) From Table 4, Lemma 2.1, Corollary 2.1, Lemma 2.19 and Lemma 2.10: For graphs G_{55} and G_{518} : When $k = 3$, $c_{k,3}(G_{55}) - c_{k,3}(G_{518}) = 2\binom{n-4}{2} > 0$, When $k \geq 4$, $c_{k,3}(G_{55}) - c_{k,3}(G_{518}) = -2[\binom{n-4}{k-4} - \binom{n-4}{k-1}]$, Since $n \neq 2k - 1$, we have $-2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \neq 0$, Thus G_{55} and G_{518} are determined by their k -th hook immanantal polynomial. For graphs G_{55} and G_{520} , $c_{k,3}(G_{55}) - c_{k,3}(G_{520}) = -2\binom{n-1}{k-1} \neq 0$, so G_{55} and G_{520} are determined by their k -th hook immanantal polynomial. For graphs G_{55} and G_{523} with $n > 7$: When $k = 3$, $c_{k,4}(G_{55}) - c_{k,4}(G_{523}) = \frac{(5-n)}{2}\binom{n-2}{1} - 4\binom{n-4}{2} - 2\binom{n-5}{2} < 0$, When $k = 4$, $c_{k,4}(G_{55}) - c_{k,4}(G_{523}) = \frac{(7-n)}{3}\binom{n-2}{2} + 4[1 - \binom{n-4}{3}] - 2\binom{n-5}{3} = \frac{-(n-7)(7n^2-59n+138)}{6}$, which has no positive integer solutions for $n > 7$. When $k \geq 5$, $c_{k,4}(G_{55}) - c_{k,4}(G_{523}) = \frac{(2k-n-1)}{k-1}\binom{n-2}{k-2} + 4[\binom{n-4}{k-4} - \binom{n-4}{k-1}] + 2[\binom{n-5}{k-5} - \binom{n-5}{k-1}] \neq 0$, Hence G_{55} and G_{523} are determined by their k -th hook immanantal polynomial. For graphs G_{518} and G_{520} : When $k = 3$, $c_{k,3}(G_{518}) - c_{k,3}(G_{520}) = -2\binom{n-1}{2} - 2\binom{n-4}{2} < 0$, When $4 \leq k \leq n - 3$, $c_{k,3}(G_{518}) - c_{k,3}(G_{520}) = -2\binom{n-1}{k-1} + 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \leq -2\binom{n-4}{k-1} < 0$, When $k = n - 2$, $c_{k,3}(G_{518}) - c_{k,3}(G_{520}) = -2\binom{n-1}{n-3} + 2\binom{n-4}{n-6} \neq 0$, When $k = n - 1$, $c_{k,3}(G_{518}) - c_{k,3}(G_{520}) = -2\binom{n-1}{n-2} + 2\binom{n-4}{n-5} \neq 0$, When $k = n$, $c_{k,3}(G_{518}) - c_{k,3}(G_{520}) = 0$, $c_{k,4}(G_{518}) - c_{k,4}(G_{520}) = -1$, Therefore G_{518} and G_{520} are determined by their k -th hook immanantal polynomial. For graphs G_{518} and G_{523} : When $k = 3$, $c_{k,3}(G_{518}) - c_{k,3}(G_{523}) = -2\binom{n-4}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{518}) - c_{k,3}(G_{523}) = 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}]$, Since $n \neq 2k - 1$, we have $2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \neq 0$. Thus G_{518} and G_{523} are determined by their k -th hook immanantal polynomial. For graphs G_{520} and G_{523} , $c_{k,3}(G_{520}) - c_{k,3}(G_{523}) = 2\binom{n-1}{k-1} \neq 0$, so G_{520} and G_{523} are determined by their k -th hook immanantal polynomial.
- (vi) From Table 4, Lemma 2.1 and Lemma 2.10, for graphs G_{59} and G_{516} : When $k = 3$, $c_{k,3}(G_{59}) - c_{k,3}(G_{516}) = -4\binom{n-1}{2} - 2\binom{n-4}{2} < 0$, When $k \geq 4$, $c_{k,3}(G_{59}) - c_{k,3}(G_{516}) = -4\binom{n-1}{k-1} - 2[\binom{n-4}{k-4} - \binom{n-4}{k-1}] \leq -2\binom{n-1}{k-1} - 2\binom{n-4}{k-4} < 0$, Thus, the graphs G_{59} and G_{516} are determined by their k -th hook immanantal polynomial.
- (vii) From Table 4, Lemma 2.1 and Corollary 2.1, for graphs G_{50} and G_{51} with $n > 7$: When $k = 3$, $c_{k,4}(G_{50}) - c_{k,4}(G_{51}) = -\frac{(5-n)}{2}\binom{n-2}{1} + 4\binom{n-4}{2} + 2\binom{n-5}{2} > 0$, When $k = 4$, $c_{k,4}(G_{50}) - c_{k,4}(G_{51}) = -\frac{(7-n)}{3}\binom{n-2}{2} - 4[1 - \binom{n-4}{3}] + 2\binom{n-5}{3} = \frac{(n-7)(7n^2-59n+138)}{6}$, which has no positive

integer solutions for $n > 7$. When $k \geq 5$, $c_{k,4}(G_{50}) - c_{k,4}(G_{51}) = -1 \frac{(2k-n-1)}{k-1} \binom{n-2}{k-2} - 4[\binom{n-4}{k-4} - \binom{n-4}{k-1}] - 2[\binom{n-5}{k-5} - \binom{n-5}{k-1}] \neq 0$, Therefore, the graphs G_{50} and G_{51} are determined by their k -th hook immanantal polynomial. □

Proof of Theorem 1.1. Theorem 1.1 can be established directly from Lemmas 2.3-2.6 and Lemmas 3.1–3.4. □

4 Conclusion

In this paper, we mainly prove that for any fixed partition $(k, 1^{n-k})$ with $n \neq 2k - 1$, every subgraph obtained by deleting at most five edges from the complete graph is determined by its k -th hook immanantal polynomial of the Laplacian matrix. In fact, from Lemma 2.1 we know that when $n = 2k - 1$, the binomial differences $\binom{n-4}{k-4} - \binom{n-4}{k-1}$, $\binom{n-5}{k-5} - \binom{n-5}{k-1}$ equal zero. Hence the method used to prove Theorem 1.1 fails in this case, and new techniques need to be developed. Furthermore, many interesting problems remain for further investigation:

- (i) Are the graphs in $\mathcal{G}_n$ determined by the k -th hook immanantal polynomial when $n = 2k - 1$?
- (ii) What families of graphs are determined by the k -th hook immanantal polynomial of their graph matrix?

References

- [1] A.E. Brouwer, W.H. Haemers, Spectra of graphs, *Springer*. 2011.
- [2] C.D. Godsil, Algebraic Combinatorics, *Chapman and Hall, New York*. 1993.
- [3] G. Cash, Immanants and immanantal polynomials of chemical graphs, *J. Chem. Inform. Comput. Sci.* 43(2003) 1942–1946.
- [4] M. Cámara, W.H. Haemers, Spectral characterizations of almost complete graphs, *Discrete Appl. Math.* 76(2014) 19–23.
- [5] C.D. Godsil, Graphical Evolution, An introduction to the theory of random graphs, *SIAM Review*. 29(2012) 651–652.
- [6] B. Davide, S. Paolo, Immanant varieties, *Linear Algebra Appl.* 682(2024) 164–190.
- [7] I.P. Goulden, D.M. Jackson, Immanants of combinatorial matrices, *J. Algebra* 148(1992) 305–324.
- [8] E.R. Van, W.H. Haemers, Which graphs are determined by their spectrum?, *Linear Algebra Appl.* 373(2003) 241–272.
- [9] G.F. Royle, C.D. Godsil, Algebraic graph theory, *New York. Springer*. 2001.
- [10] G. Yu, H. Qu, The coefficients of the immanantal polynomial, *Appl. Math. Comput.* 339(2018) 38–44.
- [11] H. Zhang, T. Wu, H. Lai, Per-spectral characterizations of some edge-deleted subgraphs of a complete graph, *Linear Multilinear Algebra* 63(2015) 397–410.

- [12] M.K. Nagar, S. Sivasubramanian, Laplacian immanantal polynomials and the GTS poset on trees, *Linear Algebra Appl.* 561(2019) 1–23.
- [13] O. Chan, T. Lam, Immanant inequalities for laplacians of trees, *SIAM J. Matrix Anal Appl.* 21(2006) 129–144.
- [14] P. Botti, R. Merris, Almost all trees share a complete set of immanantal polynomials, *J. Graph Theory.* 17(1993) 467–476.
- [15] P. Bürgisser, The computational complexity of immanants, *SIAM J. Comput.* 30(2000) 1023–1040.
- [16] R.A. Brualdi, Introductory combinatorics, *North-Holland* (2009).
- [17] R. Merris, Single-hook characters and hamiltonian circuits, *Linear Multilinear Algebra* 14(1983) 21–35.
- [18] R. Merris, Immanantal invariants of graphs, *Linear Algebra Appl.* 401(2003) 67–75.
- [19] R. Merris, The second immanantal polynomial and the centroid of a graph, *SIAM J. Algebraic Discrete Methods.* 7(2005) 484–503.
- [20] R. Merris, K.R. Rebman, W. Watkins, Permanental polynomials of graphs, *Linear Algebra Appl.* 38(1981) 273–288.
- [21] S. Liu, On the (Signless) Laplacian Permanental Polynomials of Graphs, *Graphs Combin.* 35(2019) 787–803.
- [22] S. Liu, H. Zhang, On the characterizing properties of the permanental polynomials of graphs, *Linear Algebra Appl.* 438(2013) 157–172.
- [23] L. Chen, L. Sun, C. Bu, High-order spectral characterizations of graphs, *Discrete Math.* 348(2025) 114421.
- [24] F. Belardo et al., Open problems in the spectral theory of signed graphs, *Art Discrete Appl. Math.* (2019).
- [25] T. Wu, T. Zhou, The characterizing properties of (signless) laplacian permanental polynomials of almost complete graphs, *J. Math.* (2021) 9161508.
- [26] T. Wu et al., On the second immanantal polynomials of graphs, *Discrete Appl Math.* 347(2024) 114105.
- [27] T. Wu, Y. Yu, X. Gao, The second immanantal polynomials of Laplacian matrices of unicyclic graphs, *Appl. Math. Comput.* 451(2023) 128038.
- [28] T. Wu, Some edge-deleted subgraphs of complete graph are determined by their spectrums, *Adv. Appl. Discrete Math.* 11(2013) 143–160.
- [29] T. Wu, Spectral characterization of the edge-Deleted subgraphs of complete graph, *J. Math. Res. Appl.* 30(2010) 833–840.
- [30] X. Liu et al., On the spectral characterization of some unicyclic graphs, *Discrete Math.* 311(2011) 2317–2336.

- [31] E.R. Van, W.H. Haemers, Developments on spectral characterizations of graphs, *Discrete Math.* 309(2009) 576–586.
- [32] X. Dong, T. Wu, Immanantal polynomials of the linear combination matrices of graphs <https://arxiv.org/abs/2604.04489>
- [33] X. Zeng, T. Wu, J. Pan, Almost complete graphs are determined by the second immanantal polynomials, *J. Southwest China Norm. Univ. (Nat. Sci. Ed.)* 48(2023) 37–44.