Hot Wings: Quantifying the Impact of Large Appendages on the Heat Loss Potential and Migratory Decision-making of Humpback Whales

William T. Gough[1], Martin van Aswegen[1], Lewis Evans[1], Andy Szabo[1,2], Jens J. Currie[1,3] Andreas Fahlman[4,5], Fredrik Christiansen[6], Augusta Hollers[1], Frank E. Fish[7], Kristi West[8], Jana Phipps[8], Nicholas Hofmann[8], Lars Bejder[1]

Affiliations

1) Marine Mammal Research Program, Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, Kaneohe, HI 96744 USA
2) Alaska Whale Foundation, PO Box 1927, Petersburg, AK 99833 USA
3) Pacific Whale Foundation, 300 Ma'alaea Road, Suite 211, Wailuku, HI 96793 USA
4) Fundacion Oceanografic, Carrer d'Eduardo Primo Yúfera, 1B, 46013, Valencia, Spain
5) Global Diving Research SL, Sanlucar de Barrmeda, 11540, Spain
6) Department of Ecoscience, Aarhus University, 4000 Roskilde, Denmark
7) West Chester University, 700 S High St, West Chester, PA 19383 USA
8) Health and Stranding Lab, College of Tropical Agriculture and Human Resources, University of Hawai'i at Mānoa, Honolulu, HI 96822 USA

Correspondence: wgough@hawaii.edu

Acknowledgements

We would like to thank the large team of students, interns, and collaborators who made the collection of our morphometric dataset possible. We would like to thank Dr. Peter T. Madsen for his helpful feedback on early drafts of this manuscript.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: W.T.G., L.B.; Data Curation: W.T.G., M.v.A., L.E., A.S., J.J.C., A.H., F.E.F., K.L.W., J.P., N.H.; Formal Analysis: W.T.G., M.v.A., F.E.F., J.P., N.H.; Funding Acquisition: A.S., J.J.C., K.L.W., L.B.; Investigation: W.T.G., M.v.A., L.E., A.S., J.J.C., A.H., F.E.F., K.L.W., J.P., N.H., L.B.; Methodology: W.T.G., M.v.A., A.F., F.C., F.E.F., L.B.; Project Administration: L.B.; Resources: A.S., K.L.W., L.B.; Software: W.T.G., M.v.A., L.E., A.H.; Visualization: W.T.G.; Writing – Original Draft: W.T.G.; Writing – Review & Editing: all coauthors

Funding

This research was funded in part by grants from the Office of Naval Research (N00014-22-1-2721, N00014-24-1-2454, and N00014-20-1-2624), the United States Pacific Fleet Environmental Readiness Division (no. W9126G2220033), NOAA Fisheries via the Cooperative Ecosystem Studies Unit (CESU) award NA19NMF4720181, the United States Navy's Living Marine Resources Program (N39430-25-C-2006), the National Marine Sanctuary Foundation (21-10-B-334), the Department of Defense's Defense University Research Instrumentation Program (N00014-19-1-2612 and N00014-21-1-2249), the Marine Mammal Commission (MMC22-059), and Dolphin Quest (130-4010-4 and 130-6610-4). UAS-photogrammetry data from Southeast Alaska was collected with additional funding support from Lindblad Expeditions-National Geographic Conservation Fund, the North Pacific Research Board (no. 2114), and the NOAA National Marine Fisheries Service (no. 19703). Stranding response and morphometrics were supported by the NOAA National Marine Fisheries Service (NMFS) John H. Prescott Marine Mammal Rescue Assistance Grant program over many years and was conducted under NOAA/NMFS permits 18786 and 24359. Additional funding support was provided by philanthropic donations, the members and supporters of Pacific Whale Foundation, the Omidyar Ohana Foundation, Young Brothers Hawaii, Wildspace Productions, Freeborne Media, and Netflix, Inc.

Data and resource availability

Data related to the analyses in this study can be found at Data Dryad (DOI: XXXXX) or at https://github.com/wgough/XXXXXXXX. For more information, please contact the authors. All relevant data and details of resources can be found within the article and its supplementary information.