

Subgroups

1. Subgroups

Definition

Cyclic subgroups

Definition

Definition 1

Let (G, \star) denote a group. A **subgroup** $H < G$ is a subset of G such that (H, \star) is also a group.

Example 1

The set $6\mathbb{Z} \subset \mathbb{Z}$ of multiples of 6 forms a subgroup of \mathbb{Z} .

Proof: We must verify the group axioms:

- Is $+$ binary on $6\mathbb{Z}$? (Verifying this condition is sometimes called “showing $6\mathbb{Z}$ is **closed under addition**”.)

Suppose $g, h \in 6\mathbb{Z}$. Since g and h are multiples of 6 then there exist (integers) n and m such that $g = 6n$ and $h = 6m$. We must show $g + h \in 6\mathbb{Z}$:

$$\begin{aligned} g + h &= 6n + 6m \\ &= \underbrace{n + \cdots + n}_{6 \text{ times}} + \underbrace{m + \cdots + m}_{6 \text{ times}} \\ &= \underbrace{(n + m) + \cdots + (n + m)}_{6 \text{ times}} \text{ (by associativity in } G) \\ &= 6(n + m). \end{aligned}$$

Since \mathbb{Z} is a group $n + m \in \mathbb{Z}$ and hence $g + h = 6(n + m)$ is an integer multiple of 6. Therefore $g + h \in 6\mathbb{Z}$.

- Is $+$ associative in $6\mathbb{Z}$?

In general we never have to prove this property since $6\mathbb{Z} \subset \mathbb{Z}$ and therefore $+$ inherits associativity from \mathbb{Z} .

- Does $6\mathbb{Z}$ contain the identity element?

In our case we need to show the identity element is a multiple of 6. The additive identity of \mathbb{Z} is $0 = 6 \cdot 0$, so $0 \in 6\mathbb{Z}$.

- Does every element in $6\mathbb{Z}$ have an inverse in $6\mathbb{Z}$ (i.e., is $6\mathbb{Z}$ “closed under inverses”)?

Suppose $g \in 6\mathbb{Z}$, and write $g = 6n$ for some integer n . Then $g + (-g) = (6n) + (-6n) = 0 = (-6n) + (6n)$. We conclude $-g = -6n$. And, $-g = -6n = 6(-n) \in 6\mathbb{Z}$.



The following is a shortcut for proving a subset is a subgroup.

Proposition 1

Suppose $G = (G, \star)$ is a group and H is a non-empty subset of G . Then $H < G$ if $gh^{-1} \in H$ for every $g, h \in H$.

Question

Why do we suppose H is non-empty?

Exercise 1

Prove Proposition 1. Is the converse true?

Example 2 (cf. Problem 49)

Suppose H is a subset of $G = (G, \star)$ satisfying the following:

- (i) H is closed under \star .
- (ii) If $g \in H$ then $g^{-1} \in H$.

Then $H < G$.

Proof: Suppose $g, h \in H$. By (i) $h^{-1} \in H$ and by (ii) $gh^{-1} \in H$. It follows from Proposition 1 that $H < G$. □

Exercise 2 (cf. Problem 50)

Let $G = \mathbb{Z}_{12}$, as defined in Exercise ?? . Show that $H = \{0, 3, 6, 9\}$ is a subgroup of G .

Cyclic subgroups

Definition 2

Suppose G is a group and $g \in G$. Define

$$\langle g \rangle := \{g^n \mid n \in \mathbb{Z}\}$$

as the **cyclic subgroup** of G **generated by** g .

Exercise 3

For Definition 2 to make sense, we must check $\langle g \rangle$ actually is a subgroup.

As alternative notation, authors may write $\langle g \rangle$ to denote the subgroup in G generated by g (see Example 1).

Example 3

For any integer k , the subgroup $k\mathbb{Z} < \mathbb{Z}$ is cyclic.

(Recall, from Section ??, the subgroups $0\mathbb{Z} < \mathbb{Z}$ and $1\mathbb{Z} < \mathbb{Z}$.)

Question

What is $0\mathbb{Z}$? What is $1\mathbb{Z}$?

We can define subgroups with more than one generator, though we do not describe such subgroups as cyclic.

Definition 3

Let $S = \{s_1, \dots, s_k\}$ denote some set of elements in the group G . The subgroup generated by S is defined as

$$\langle S \rangle := \{s_1^{n_1} \cdots s_k^{n_k} \mid n_i \in \mathbb{Z} \text{ for all } i = 1, \dots, k\}.$$

Elements in $\langle S \rangle$ are called **words**.

Question

How would you rewrite Definition 3 in additive notation?

Exercise 4 (cf. Problem 51)

Prove $\langle S \rangle$ in Definition 3 is a subgroup of G .

Example 4 (cf. Problem 52)

In \mathbb{Z}_{12} , we list the elements of the subgroup $H = \langle 2, 3 \rangle$ by writing down **\mathbb{Z} -linear combinations** of the generators, i.e., all possible elements of the form $n_1 \cdot 2 + n_2 \cdot 3$ for $n_1, n_2 \in \mathbb{Z}$:

$$\begin{array}{llll} 1 \cdot 2 + 0 \cdot 3 = 2 & 0 \cdot 2 + 1 \cdot 3 = 3 & 1 \cdot 2 + 1 \cdot 3 = 5 & (-1) \cdot 2 + 1 \cdot 3 = 1 \\ 2 \cdot 2 + 0 \cdot 3 = 4 & 0 \cdot 2 + 3 \cdot 3 = 9 & 2 \cdot 2 + 1 \cdot 3 = 7 & \\ 3 \cdot 2 + 0 \cdot 3 = 6 & & 4 \cdot 2 + 1 \cdot 3 = 11 & \\ 4 \cdot 2 + 0 \cdot 3 = 8 & & & \\ 5 \cdot 2 + 0 \cdot 3 = 10 & & & \\ 6 \cdot 2 + 0 \cdot 3 = 0 & & & \end{array}$$

Having exhausted all possible elements, we conclude $\langle 2, 3 \rangle = \mathbb{Z}_{12}$.