

Calculus I (Math 2554)

Summer 2015

Dr. Ashley K. Wheeler

University of Arkansas

last updated: May 20, 2015

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Table of Contents I

1 Week 1: 26-29 May

- Tips for Success

§2.1 The Idea of Limits

- Book Problems

§2.2 Definition of Limits

- Determining Limits from a Graph
- Determining Limits from a Table
- One-Sided Limits
- Relationship Between One- and Two-Sided

Limits

- Book Problems

§2.3 Techniques for Computing Limits

- Limit Laws

- Limits of Polynomials and Rational Functions
- Additional (Algebra) Techniques
- Another Technique: Squeeze Theorem
- Book Problems

§2.4 Infinite Limits

- Definition of Infinite Limits
- Definition of a Vertical Asymptote
- Book Problems

§2.5 Limits at Infinity

- Horizontal Asymptotes
- Infinite Limits at Infinity
- Algebraic and Transcendental Functions
- Book Problems

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Tips for Success

- Attend class every day. **Participate**, take notes, and ask questions.
- Don't get behind on MLP homeworks. Stay on top of the book problems.
- Be sure to seek assistance (tutoring, office hours, etc.) if you are struggling.
- Don't rely on success in high school calculus to save you in college calculus.
- Find a study partner(s) to meet with on a regular basis to cover questions and study for quizzes/exams.
- REMEMBER... THE TERM STARTS TODAY! SO DOES THE EVENTUAL EARNING OF YOUR FINAL GRADE!!!

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

2.1 The Idea of Limits

Question

How would you define, and then differentiate between, the following pairs of terms?

- instantaneous velocity vs. average velocity?
- tangent line vs. secant line?

(Recall: What is a tangent line and what is a secant line?)

Example

An object is launched into the air. Its position s (in feet) at any time t (in seconds) is given by the equation:

$$s(t) = -4.9t^2 + 30t + 20.$$

-
- (a) Compute the average velocity of the object over the following time intervals: $[1, 3]$, $[1, 2]$, $[1, 1.5]$
 - (b) As your interval gets shorter, what do you notice about the average velocities? What do you think would happen if we computed the average velocity of the object over the interval $[1, 1.2]$? $[1, 1.1]$? $[1, 1.05]$?

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Example, cont.

An object is launched into the air. Its position s (in feet) at any time t (in seconds) is given by the equation:

$$s(t) = -4.9t^2 + 30t + 20.$$

-
- (c) How could you use the average velocities to estimate the instantaneous velocity at $t = 1$?
 - (d) What do the average velocities you computed in 1. represent on the graph of $s(t)$?

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Question

What happens to the relationship between **instantaneous** velocity and **average** velocity as the time interval gets shorter?

Answer: The instantaneous velocity at $t = 1$ is the limit of the average velocities as t approaches 1.

Question

What about the relationship between the **secant** lines and the **tangent** lines as the time interval gets shorter?

Answer: The slope of the tangent line at $(1, 45.1 = s(1))$ is the limit of the slopes of the secant lines as t approaches 1.

2.1 Book Problems


1-3, 7, 9, 11, 13, 17, 21

- Even though book problems aren't turned in, they're a very good way to study for quizzes and tests (wink wink wink).

2.2 Definition of Limits

Determining Limits from a Graph

Exercise



Determine the following:

- (a) $h(1)$
- (b) $h(2)$
- (c) $h(4)$
- (d) $\lim_{x \rightarrow 2} h(x)$
- (e) $\lim_{x \rightarrow 4} h(x)$
- (f) $\lim_{x \rightarrow 1} h(x)$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Question

Does $\lim_{x \rightarrow a} f(x)$ always equal $f(a)$?

(Hint: Look at the example from the previous slide!)

Determining Limits from a Table

Exercise

Suppose $f(x) = \frac{x^2 + x - 20}{x - 4}$.

(a) Create a table of values of $f(x)$ when

$$x = 3.9, 3.99, 3.999, \text{ and}$$

$$x = 4.1, 4.01, 4.001$$

(b) What can you conjecture about $\lim_{x \rightarrow 4} f(x)$?

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

One-Sided Limits

Up to this point we have been working with two-sided limits; however, for some functions it makes sense to examine one-sided limits.

Notice how in the previous example we could approach $f(x)$ from both sides as x approaches a , i.e., when $x > a$ and when $x < a$.

Definition (right-hand limit)

Suppose f is defined for all x near a with $x > a$. If $f(x)$ is arbitrarily close to L for all x sufficiently close to a with $x > a$, we write

$$\lim_{x \rightarrow a^+} f(x) = L$$

and say **the limit of $f(x)$ as x approaches a from the right equals L .**

Definition (left-hand limit)


Suppose f is defined for all x near a with $x < a$. If $f(x)$ is arbitrarily close to L for all x sufficiently close to a with $x < a$, we write

$$\lim_{x \rightarrow a^-} f(x) = L$$

and say **the limit of $f(x)$ as x approaches a from the left equals L .**

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Exercise



Determine the following:

(a) $g(2)$

(b) $\lim_{x \rightarrow 2^+} g(x)$

(c) $\lim_{x \rightarrow 2^-} g(x)$

(d) $\lim_{x \rightarrow 2} g(x)$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Relationship Between One- and Two-Sided Limits

Theorem

If f is defined for all x near a except possibly at a , then $\lim_{x \rightarrow a} f(x) = L$ if and only if **both** $\lim_{x \rightarrow a^+} f(x) = L$ **and** $\lim_{x \rightarrow a^-} f(x) = L$.

In other words, the only way for a two-sided limit to exist is if the one-sided limits equal the same number (L).

2.2 Book Problems

1-4, 10, 12, 16, 18, 20, 25

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

2.3 Techniques for Computing Limits

This section provides various laws and techniques for determining limits. These constitute **analytical** methods of finding limits. The following is an example of a very useful limit law:

Limits of Linear Functions: Let a , b , and m be real numbers. For linear functions $f(x) = mx + b$,

$$\lim_{x \rightarrow a} f(x) = f(a) = ma + b.$$

This rule says we if $f(x)$ is a linear function, then in taking the limit as $x \rightarrow a$, we can just plug in the a for x .

IMPORTANT! Using a table or a graph to compute limits, as in the previous sections, can be helpful. However, “analytical” does not include those techniques.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limit Laws

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist, c is a real number, and m, n are positive integers.

1. Sum:
$$\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

2. Difference:
$$\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$$

In other words, if we are taking a limit of two things added together or subtracted, then we can first compute each of their individual limits one at a time.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limit Laws, cont.

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist, c is a real number, and m, n are positive integers.

3. Constant Multiple: $\lim_{x \rightarrow a} (cf(x)) = c \left(\lim_{x \rightarrow a} f(x) \right)$

4. Product: $\lim_{x \rightarrow a} (f(x)g(x)) = \left(\lim_{x \rightarrow a} f(x) \right) \left(\lim_{x \rightarrow a} g(x) \right)$

The same is true for products. If one of the factors is a constant, we can just bring it outside the limit. In fact, a constant is its own limit.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limit Laws, cont.

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist, c is a real number, and m, n are positive integers.

5. Quotient:
$$\lim_{x \rightarrow a} \left(\frac{f(x)}{g(x)} \right) = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$$

(provided $\lim_{x \rightarrow a} g(x) \neq 0$)

Question

Why the caveat?

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limit Laws, cont.

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist, c is a real number, and m, n are positive integers.

6. Power: $\lim_{x \rightarrow a} (f(x))^n = \left(\lim_{x \rightarrow a} f(x) \right)^n$

7. Fractional Power: $\lim_{x \rightarrow a} (f(x))^{\frac{n}{m}} = \left(\lim_{x \rightarrow a} f(x) \right)^{\frac{n}{m}}$

(provided $f(x) \geq 0$ for x near a if m is even and $\frac{n}{m}$ is in lowest terms)

Question

Explain the caveat in 7.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limit Laws, cont.

Laws **1.-6.** hold for one-sided limits as well. But **7.** must be modified:

7. Fractional Power (one-sided limits):

- $\lim_{x \rightarrow a^+} (f(x))^{\frac{n}{m}} = \left(\lim_{x \rightarrow a^+} f(x) \right)^{\frac{n}{m}}$
(provided $f(x) \geq 0$ for x near a with $x > a$, if m is even and $\frac{n}{m}$ is in lowest terms)
- $\lim_{x \rightarrow a^-} (f(x))^{\frac{n}{m}} = \left(\lim_{x \rightarrow a^-} f(x) \right)^{\frac{n}{m}}$
(provided $f(x) \geq 0$ for x near a with $x < a$, if m is even and $\frac{n}{m}$ is in lowest terms)

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Limits of Polynomials and Rational Functions

Assume that $p(x)$ and $q(x)$ are polynomials and a is a real number.

- **Polynomials:** $\lim_{x \rightarrow a} p(x) = p(a)$
- **Rational functions:** $\lim_{x \rightarrow a} \frac{p(x)}{q(x)} = \frac{p(a)}{q(a)}$
(provided $q(a) \neq 0$)

For polynomials and rational functions we can plug in a to compute the limit, as long as we don't get zero in the denominator. Linear functions count as polynomials. A rational function is a “fraction” made of polynomials.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Exercise

Evaluate the following limits analytically.

1. $\lim_{x \rightarrow 1} \frac{4f(x)g(x)}{h(x)}$, given that

$$\lim_{x \rightarrow 1} f(x) = 5, \lim_{x \rightarrow 1} g(x) = -2, \text{ and } \lim_{x \rightarrow 1} h(x) = -4.$$

2. $\lim_{x \rightarrow 3} \frac{4x^2 + 3x - 6}{2x - 3}$

3. $\lim_{x \rightarrow 1^-} g(x)$ and $\lim_{x \rightarrow 1^+} g(x)$, given that

$$g(x) = \begin{cases} x^2 & \text{if } x \leq 1; \\ x + 2 & \text{if } x > 1. \end{cases}$$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Additional (Algebra) Techniques

When direct substitution (a.k.a. plugging in a) fails try using algebra:

- Factor and see if the denominator cancels out.

Example

$$\lim_{t \rightarrow 2} \frac{3t^2 - 7t + 2}{2 - t}$$

- Look for a common denominator.

Example

$$\lim_{h \rightarrow 0} \frac{\frac{1}{5+h} - \frac{1}{5}}{h}$$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Another Technique: Squeeze Theorem

This method for evaluating limits uses the relationship of functions with each other.

Theorem (Squeeze Theorem)

Assume $f(x) \leq g(x) \leq h(x)$ for all values of x near a , except possibly at a , and suppose

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L.$$

Then since g is always between f and h for x -values close enough to a , we must have

$$\lim_{x \rightarrow a} g(x) = L.$$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Example

(a) Draw a graph of the inequality

$$-|x| \leq x^2 \ln(x^2) \leq |x|.$$

(b) Compute $\lim_{x \rightarrow 0} x^2 \ln(x^2)$.

2.3 Book Problems

12-30 (every 3rd problem), 31, 33, 37-47 (odds), 51, 53, 61-65 odds

- In general, review your algebra techniques, since they can save you some headache.

2.4 Infinite Limits

In the next two sections, we examine limit scenarios involving infinity.
The two situations are:

- **Infinite limits:** as x (i.e., the independent variable) approaches a finite number, y (i.e., the dependent variable) becomes arbitrarily large or small

looks like: $\lim_{x \rightarrow \text{number}} f(x) = \pm\infty$

- **Limits at infinity:** as x approaches an arbitrarily large or small number, y approaches a finite number

looks like: $\lim_{x \rightarrow \pm\infty} f(x) = \text{number}$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.


Definition of Infinite Limits

Definition (positively infinite limit)

Suppose f is defined for all x near a . If $f(x)$ grows arbitrarily large for all x sufficiently close (but not equal) to a , we write

$$\lim_{x \rightarrow a} f(x) = \infty$$

and say **the limit of $f(x)$ as x approaches a is infinity.**




The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Definition (negatively infinite limit)

Suppose f is defined for all x near a . If $f(x)$ is negative and grows arbitrarily large in magnitude for all x sufficiently close (but not equal) to a , we write

$$\lim_{x \rightarrow a} f(x) = -\infty$$

and say **the limit of $f(x)$ as x approaches a is negative infinity**.



The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

The definitions work for one-sided limits, too.

Exercise

Using a graph and a table of values, given $f(x) = \frac{1}{x^2 - x}$, determine:

(a) $\lim_{x \rightarrow 0^+} f(x)$

(b) $\lim_{x \rightarrow 0^-} f(x)$

(c) $\lim_{x \rightarrow 1^+} f(x)$

(d) $\lim_{x \rightarrow 1^-} f(x)$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Definition of Vertical Asymptote

Definition

Suppose a function f satisfies at least one of the following:

- $\lim_{x \rightarrow a} f(x) = \pm\infty$,
- $\lim_{x \rightarrow a^+} f(x) = \pm\infty$
- $\lim_{x \rightarrow a^-} f(x) = \pm\infty$

Then the line $x = a$ is called a **vertical asymptote** of f .

Exercise

Given $f(x) = \frac{3x - 4}{x + 1}$, determine, analytically (meaning without using a table or a graph),

(a) $\lim_{x \rightarrow -1^+} f(x)$

(b) $\lim_{x \rightarrow -1^-} f(x)$

Remember to check for factoring –

Exercise

(a) What is/are the vertical asymptotes of

$$f(x) = \frac{3x^2 - 48}{x + 4}?$$

(b) What is $\lim_{x \rightarrow -4} f(x)$?

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

2.4 Book Problems

7-10, 15, 17-26, 36-37

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

\int 2.5 Limits at Infinity

Limits at infinity determine what is called the **end behavior** of a function.

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Horizontal Asymptotes

Definition

If $f(x)$ becomes arbitrarily close to a finite number L for all sufficiently large and positive x , then we write

$$\lim_{x \rightarrow \infty} f(x) = L.$$

The line $y = L$ is a **horizontal asymptote** of f .

The limit at negative infinity, $\lim_{x \rightarrow -\infty} f(x) = M$, is defined analogously and in this case, the horizontal asymptote is $y = M$.

Infinite Limits at Infinity

Question

Is it possible for a limit to be both an infinite limit and a limit at infinity?
(Yes.)

If $f(x)$ becomes arbitrarily large as x becomes arbitrarily large, then we write

$$\lim_{x \rightarrow \infty} f(x) = \infty.$$

(The limits $\lim_{x \rightarrow \infty} f(x) = -\infty$, $\lim_{x \rightarrow -\infty} f(x) = \infty$, and $\lim_{x \rightarrow -\infty} f(x) = -\infty$ are defined similarly.)

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Powers and Polynomials: Let n be a positive integer and let $p(x)$ be a polynomial.

- $n = \text{even number}$: $\lim_{x \rightarrow \pm\infty} x^n = \infty$
- $n = \text{odd number}$: $\lim_{x \rightarrow \infty} x^n = \infty$ and $\lim_{x \rightarrow -\infty} x^n = -\infty$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

- (again, assuming n is positive)

$$\lim_{x \rightarrow \pm\infty} \frac{1}{x^n} = \lim_{x \rightarrow \pm\infty} x^{-n} = 0$$

- For a polynomial, only look at the term with the highest exponent:

$$\lim_{x \rightarrow \pm\infty} p(x) = \lim_{x \rightarrow \pm\infty} (\text{constant}) \cdot x^n$$

The constant is called the **leading coefficient**, $\text{lc}(p)$. The highest exponent that appears in the polynomial is called the **degree**, $\deg(p)$.

Rational Functions: Suppose $f(x) = \frac{p(x)}{q(x)}$ is a rational function.

- If $\deg(p) < \deg(q)$, i.e., **the numerator has the smaller degree**, then

$$\lim_{x \rightarrow \pm\infty} f(x) = 0$$

and $y = 0$ is a horizontal asymptote of f .

- If $\deg(p) = \deg(q)$, i.e., **numerator and denominator have the same degree**, then

$$\lim_{x \rightarrow \pm\infty} f(x) = \frac{\text{lc}(p)}{\text{lc}(q)}$$

and $y = \frac{\text{lc}(p)}{\text{lc}(q)}$ is a horizontal asymptote of f .

- If $\deg(p) > \deg(q)$, (**numerator has the bigger degree**) then

$$\lim_{x \rightarrow \pm\infty} f(x) = \infty \quad \text{or} \quad -\infty$$

and f has no horizontal asymptote.

- Assuming that $f(x)$ is in **reduced form** (p and q share no common factors), vertical asymptotes occur at the zeroes of q .

(This is why it is a good idea to check for factoring and cancelling first!)

Exercise

Determine the **end behavior** of the following functions (in other words, compute both limits, as $x \rightarrow \pm\infty$, for each of the functions):

1. $f(x) = \frac{x+1}{2x^2-3}$

2. $g(x) = \frac{4x^3-3x}{2x^3+5x^2+x+2}$

3. $h(x) = \frac{6x^4-1}{4x^3+3x^2+2x+1}$

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

Algebraic and Transcendental Functions

Example

Determine the end behavior of the following functions.

1. $f(x) = \frac{4x^3}{2x^3 + \sqrt{9x^6 + 15x^4}}$ (radical signs appear)
2. $g(x) = \cos x$ (trig)
3. $h(x) = e^x$ (exponential)

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.

2.5 Book Problems

9-10, 13-35 (odds), 39, 43, 45, 53

The base for these slides was done by Shannon Dingman, later encoded in L^AT_EX by Brad Lutes.