Defining equations for matroid varieties – using the Grassmann-Cayley algebra Virtual Inspiring Talks Series

Ashley K. Wheeler

Mount Holyoke College

Fall 2020

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra

Thank-you for the invitation to speak!

Thank-you for the invitation to speak!

About me:

Thank-you for the invitation to speak!

About me:

• Biracial, 2nd generation college student

Thank-you for the invitation to speak!

- Biracial, 2nd generation college student
- Military family

Thank-you for the invitation to speak!

- Biracial, 2nd generation college student
- Military family
- Education: USAFA

Thank-you for the invitation to speak!

- Biracial, 2nd generation college student
- Military family
- Education: USAFA → KState

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra

Thank-you for the invitation to speak!

- Biracial, 2nd generation college student
- Military family
- ullet Education: USAFA o KState o Michigan

About the project:

About the project:

Joint w/ Jessica Sidman (MHC) and Will Traves (USNA).





About the project:

Joint w/ Jessica Sidman (MHC) and Will Traves (USNA).





 Paper accepted to Journal of Combinatorial Theory, Series A this year (and we've already got a citation!).

About the project:

Joint w/ Jessica Sidman (MHC) and Will Traves (USNA).





- Paper accepted to Journal of Combinatorial Theory, Series A this year (and we've already got a citation!).
- Goal: To find defining equations for matroid varieties.

Outline:

- Front matter
- Introduction: What is a matroid?
- Problem: Defining equations for matroid varieties
- Paradigm shift: Matroids as point configurations
- Tool: Grassmann-Cayley algebra
- Back matter

Outline:

- Front matter
- Introduction: What is a matroid?
- Problem: Defining equations for matroid varieties
- Paradigm shift: Matroids as point configurations
- Tool: Grassmann-Cayley algebra
- Back matter

Advice: Refer to Ravi Vakil's 3 Things when viewing this (or any other) math talk!

Introduction

What is a matroid?

Q: Which collections of 3 columns form a basis for the column space of A?

$$A = \begin{pmatrix} -1 & 1 & 2 & 3 & 4 \\ -1 & 1 & 2 & 0 & -2 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

Introduction

What is a matroid?

Q: Which collections of 3 columns form a basis for the column space of A?

$$A = \begin{pmatrix} -1 & 1 & 2 & 3 & 4 \\ -1 & 1 & 2 & 0 & -2 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}$$

How can we tell?

Could row reduce A first.

$$\begin{pmatrix} -1 & 1 & 2 & 3 & 4 \\ -1 & 1 & 2 & 0 & -2 \\ 1 & 1 & 1 & 1 \end{pmatrix} \rightarrow \operatorname{rref} A = \begin{pmatrix} 1 & 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} & 0 & -\frac{3}{2} \\ 0 & 0 & 0 & 1 & 2 \end{pmatrix}$$

Could row reduce A first.

$$\begin{pmatrix} -1 & 1 & 2 & 3 & 4 \\ -1 & 1 & 2 & 0 & -2 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \rightarrow \operatorname{rref} A = \begin{pmatrix} 1 & 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 1 & \frac{3}{2} & 0 & -\frac{3}{2} \\ 0 & 0 & 0 & 1 & 2 \end{pmatrix}$$

Preferred way: Look for non-zero **3-minors** of *A* (or of rref *A*).

Fact: Columns of an $r \times r$ matrix are linearly independent if and only if the determinant of the matrix is non-zero.

Number the columns of A 1 through 5. Here are the 3-minors, according to their column indices:

$$\{1,2,3\} o 0 \qquad \{1,2,4\} o -6 \qquad \{1,2,5\} o -12 \ \{1,3,4\} o -9 \qquad \{1,3,5\} o -18 \qquad \{1,4,5\} o -9 \ \{2,3,4\} o 3 \qquad \{2,3,5\} o -6 \qquad \{2,4,5\} o -11 \ \{3,4,5\} o 0$$

Number the columns of *A* 1 through 5. Here are the 3-minors, according to their column indices:

$$\{1,2,3\} o 0 \qquad \{1,2,4\} o -6 \qquad \{1,2,5\} o -12 \ \{1,3,4\} o -9 \qquad \{1,3,5\} o -18 \qquad \{1,4,5\} o -9 \ \{2,3,4\} o 3 \qquad \{2,3,5\} o -6 \qquad \{2,4,5\} o -11 \ \{3,4,5\} o 0$$

The **matroid** \mathcal{M}_A on A is given by the set:

$$\mathfrak{B} = \{ \text{all 3-tuples except } \{1,2,3\} \text{ and } \{3,4,5\} \} \subset \{1,\dots,5\}$$

The sets in \mathcal{B} are called **bases** for the matroid \mathcal{M}_A .

Matroids are a generalization of the collections of linearly independent columns of a matrix.

Matroids are a generalization of the collections of linearly independent columns of a matrix.

Applications: Where are matroids used?

Matroids are a generalization of the collections of linearly independent columns of a matrix.

Applications: Where are matroids used?

 Combinatorial optimization: artificial intelligence, machine learning, software engineering; example: travelling salesman problem Matroids are a generalization of the collections of linearly independent columns of a matrix.

Applications: Where are matroids used?

- Combinatorial optimization: artificial intelligence, machine learning, software engineering; example: travelling salesman problem
- Coding theory: error correcting, data compression, cryptography

Matroids are a generalization of the collections of linearly independent columns of a matrix.

Applications: Where are matroids used?

- Combinatorial optimization: artificial intelligence, machine learning, software engineering; example: travelling salesman problem
- Coding theory: error correcting, data compression, cryptography
- Network theory: particle physics, biology, social networks; example: bridges of Königsberg problem

Problem

Defining equations for matroid varieties

Q: Given a matroid on a matrix A, what other matrices have the same matroid as A? (e.g., rref A)

Problem

Defining equations for matroid varieties

Q: Given a matroid on a matrix A, what other matrices have the same matroid as A? (e.g., rref A)

Consider this generic matrix:

$$X = \begin{pmatrix} x_{11} & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & x_{22} & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & x_{33} & x_{34} & x_{35} \end{pmatrix}$$

We want all possible x-values that will give the same matroid (i.e. same columns as bases) as A.

Defining equations are a system of equations in the x's whose solution set minimally cuts out all of the matrices with matroid \mathcal{M}_A .

Defining equations are a system of equations in the x's whose solution set minimally cuts out all of the matrices with matroid \mathcal{M}_A .

Call this solution set \mathcal{V}_A , the **matroid variety** on A.











Mnëv (1985), Sturmfels (1989), Knutson, Lam, & Speyer (2013): All "hell" breaks loose when we try to find defining equations for \mathcal{V}_A .











Mnëv (1985), Sturmfels (1989), Knutson, Lam, & Speyer (2013): All "hell" breaks loose when we try to find defining equations for \mathcal{V}_A .

Why?











Mnëv (1985), Sturmfels (1989), Knutson, Lam, & Speyer (2013): All "hell" breaks loose when we try to find defining equations for \mathcal{V}_A .

Why? Two reasons:

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra











Mnëv (1985), Sturmfels (1989), Knutson, Lam, & Speyer (2013): All "hell" breaks loose when we try to find defining equations for V_A .

Why? Two reasons:

• The Zariski topology (beyond the scope of this talk).

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra











Mnëv (1985), Sturmfels (1989), Knutson, Lam, & Speyer (2013): All "hell" breaks loose when we try to find defining equations for V_A .

Why? Two reasons:

- The Zariski topology (beyond the scope of this talk).
- The equations are not obvious (as we shall see).

Paradigm shift

Matroids as point configurations

Idea: Think of the columns of A as points in the plane.

Paradigm shift

Matroids as point configurations

Idea: Think of the columns of A as points in the plane.

The columns of the matrix A are vectors in 3-space, that span lines through the origin.

If we cut these lines with a plane then the lines become points in the plane.

Paradigm shift


Matroids as point configurations

Idea: Think of the columns of A as points in the plane.


The columns of the matrix A are vectors in 3-space, that span lines through the origin.

If we cut these lines with a plane then the lines become points in the plane.

Why would we do this?



Q: What do you notice about the points in relation to the matroid \mathcal{M}_A ?



Q: What do you notice about the points in relation to the matroid \mathcal{M}_A ?

Points 1, 2, and 3 are collinear if and only if $\{1, 2, 3\}$ is not a basis for \mathcal{M}_A . We say the **bracket** [123] vanishes, or equals 0.

The bracket is shorthand for determinant:

$$[123] = 0 \iff \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{vmatrix} = 0$$

This is a defining equation in the x's!

The bracket is shorthand for determinant:

$$\begin{bmatrix} 123 \end{bmatrix} = 0 \iff \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{vmatrix} = 0$$

This is a defining equation in the x's!

Q: What is another defining equation for V_A ?

The bracket is shorthand for determinant:

$$\begin{bmatrix} 123 \end{bmatrix} = 0 \iff \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{vmatrix} = 0$$

This is a defining equation in the x's!

Q: What is another defining equation for V_A ?

Are there others?

The bracket is shorthand for determinant:

$$\begin{bmatrix} 123 \end{bmatrix} = 0 \iff \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{vmatrix} = 0$$


This is a defining equation in the x's!

Q: What is another defining equation for V_A ?

Are there others? ???

Example (Ford 2013):

 $\mathcal{M}_{pencil} = matroid on a 3 \times 7 matrix with$ *non* $bases {1, 2, 7}, {3, 4, 7}, {5, 6, 7}$



Q: Which brackets vanish?



All hell breaks loose: [134][256] - [234][156] = 0 is also a defining equation for $\mathcal{V}_{\text{pencil}}$.



All hell breaks loose: [134][256] - [234][156] = 0 is also a defining equation for \mathcal{V}_{pencil} . How could we have known this?!



All hell breaks loose: [134][256] - [234][156] = 0 is also a defining equation for $\mathcal{V}_{\text{pencil}}$. How could we have known this?!

Ans: The Grassmann-Cayley algebra.

Tool

Grassmann-Cayley algebra

The Grassmann-Cayley algebra was invented to do *synthetic* projective geometry.

We do arithmetic on the points (vectors) themselves, and obtain expressions in the brackets.

Tool

Grassmann-Cayley algebra

The Grassmann-Cayley algebra was invented to do *synthetic* projective geometry.

We do arithmetic on the points (vectors) themselves, and obtain expressions in the brackets.

There are two operations:

join (\vee) : refers to the line passing through points



The line joining 1 and 2 is $1 \lor 2$, or 12.

join (\vee) : refers to the line passing through points



The line joining 1 and 2 is $1 \lor 2$, or 12.

Three points joined makes a bracket, and three collinear points make the bracket vanish.

meet (\land) : refers to the intersection of two lines




The meet of the lines $1 \lor 2$ and $3 \lor 4$ is $(1 \lor 2) \land (3 \lor 4)$, or $12 \land 34$.

meet (\land) : refers to the intersection of two lines




The meet of the lines $1 \lor 2$ and $3 \lor 4$ is $(1 \lor 2) \land (3 \lor 4)$, or $12 \land 34$.

The meet operation uses *shuffle products*, which also produce expressions in the brackets.




We have $((1 \lor 2) \land (3 \lor 4)) \lor 5 \lor 6 = 0$, because these three points are collinear.

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra



We have $((1 \lor 2) \land (3 \lor 4)) \lor 5 \lor 6 = 0$, because these three points are collinear. Here's how to apply the shuffles:

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra





We have $((1 \lor 2) \land (3 \lor 4)) \lor 5 \lor 6 = 0$, because these three points are collinear. Here's how to apply the shuffles:

$$(12 \land 34) \lor 56 = ([134]2 - [234]1)56$$

= $[134][256] - [234][156] = 0$


Example: Pascal's theorem says if six points on a conic are joined in the way illustrated below, then the resulting intersection points (7, 8, and 9) are collinear.






We have a matroid $\mathfrak{M}_{\mathsf{Pascal}}$ corresponding to this configuration of points.

Introduction: What is a matroid?
Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra



We have a matroid $\mathfrak{M}_{\mathsf{Pascal}}$ corresponding to this configuration of points. Let's find defining equations for $\mathcal{V}_{\mathsf{Pascal}}$.

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra




We have a matroid $\mathfrak{M}_{\mathsf{Pascal}}$ corresponding to this configuration of points. Let's find defining equations for $\mathcal{V}_{\mathsf{Pascal}}$.

Q: Which brackets vanish?

Theorem (Sidman, Traves, W)

The defining equations for the matroid variety V_{Pascal} include at least one quartic, three independent cubics, and three independent quadrics in the brackets, besides the vanishing brackets [127], [238], [349], [457], [568], [169], [789].

Finding the quartic: The statement of Pascal's theorem says we have $(12 \land 45) \lor (23 \land 56) \lor (34 \land 61) = 0$.



Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra

Apply the shuffle products:

$$(12 \land 45) \lor (23 \land 56) \lor (34 \land 61)$$

= $([145]2 - [245]1) \lor ([256]3 - [356]2) \lor ([361]4 - [461]3)$

Now "foil":

$$(([145]2 - [245]1) \lor ([256]3 - [356]2)) \lor ([361]4 - [461]3)$$

$$= ([145][256]23 - [145][356]22 - [245][256]13 + [245][356]12)$$

$$\lor ([361]4 - [461]3)$$

Finish "foiling":

$$\begin{aligned} &([145][256]23 - [145][356]22 - [245][256]13 + [245][356]12) \\ &\lor ([361]4 - [461]3) \\ &= [145][256][361][234] - [145][256][461][233] \\ &- [145][356][361][224] + [145][356][461][223] \\ &- [245][256][361][134] + [245][256][461][133] \\ &+ [245][356][361][124] - [245][356][461][123] \end{aligned}$$

Finish "foiling":

$$\begin{aligned} &([145][256]23 - [145][356]22 - [245][256]13 + [245][356]12) \\ &\lor ([361]4 - [461]3) \\ &= [145][256][361][234] - [145][256][461][233] \\ &- [145][356][361][224] + [145][356][461][223] \\ &- [245][256][361][134] + [245][256][461][133] \\ &+ [245][356][361][124] - [245][356][461][123] \end{aligned}$$

Q: What happens when two numbers share a bracket?

Problem: Defining equations for matroid varieties
Paradigm shift: Matroids as point configurations
Tool: Grassmann-Cayley algebra

$$[233] \leftrightarrow \begin{vmatrix} x_{12} & x_{13} & x_{13} \\ x_{22} & x_{23} & x_{23} \\ x_{32} & x_{33} & x_{33} \end{vmatrix}$$

Q: What happens to a matrix with two repeated columns?

$$[233] \leftrightarrow \begin{vmatrix} x_{12} & x_{13} & x_{13} \\ x_{22} & x_{23} & x_{23} \\ x_{32} & x_{33} & x_{33} \end{vmatrix}$$

Q: What happens to a matrix with two repeated columns?

It means we get a bunch of cancelling...

$$[145][256][361][234] - [145][256][461][233]$$

$$- [145][356][361][224] + [145][356][461][223]$$

$$- [245][256][361][134] + [245][256][461][133]$$

$$+ [245][356][361][124] - [245][356][461][123]$$

$$= [145][256][361][234] - [245][256][361][134]$$

$$+ [245][356][361][124] - [245][356][461][123] = 0$$

$$[145][256][361][234] - [145][256][461][233]$$

$$- [145][356][361][224] + [145][356][461][223]$$

$$- [245][256][361][134] + [245][256][461][133]$$

$$+ [245][356][361][124] - [245][356][461][123]$$

$$= [145][256][361][234] - [245][256][361][134]$$

$$+ [245][356][361][124] - [245][356][461][123] = 0$$

The statement of Pascal's theorem gives us a non-obvious defining equation for $\mathcal{V}_{\mathsf{Pascal}}!$

Conclusion:

Conclusion:

• Finding defining equations for matroid varieties is HARD.

Conclusion:

- Finding defining equations for matroid varieties is HARD.
- The Grassmann-Cayley algebra lets us find some of the defining equations, by using the geometry of the matroids when thought of as point configurations.

Conclusion:

- Finding defining equations for matroid varieties is HARD.
- The Grassmann-Cayley algebra lets us find some of the defining equations, by using the geometry of the matroids when thought of as point configurations.

Further work:

Conclusion:

- Finding defining equations for matroid varieties is HARD.
- The Grassmann-Cayley algebra lets us find some of the defining equations, by using the geometry of the matroids when thought of as point configurations.

Further work:

ullet Are our equations the only defining equations for $\mathcal{V}_{\mathsf{Pascal}}$?

Conclusion:

- Finding defining equations for matroid varieties is HARD.
- The Grassmann-Cayley algebra lets us find some of the defining equations, by using the geometry of the matroids when thought of as point configurations.

Further work:

- ullet Are our equations the only defining equations for ${\cal V}_{\sf Pascal}?$
- Defining equations for matroid varieties given by other point configurations (e.g., Pappus' theorem)?

Conclusion:

- Finding defining equations for matroid varieties is HARD.
- The Grassmann-Cayley algebra lets us find some of the defining equations, by using the geometry of the matroids when thought of as point configurations.

Further work:

- ullet Are our equations the only defining equations for ${\cal V}_{\sf Pascal}?$
- Defining equations for matroid varieties given by other point configurations (e.g., Pappus' theorem)?
- Defining equations for matroid varieties given by wheel graphs?

Q & A:

background from Euclidea

Q & A:

(1) What do you love most about mathematics?

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?
- (3) How did you prepare for graduate school?

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?
- (3) How did you prepare for graduate school?
- (4) What was the most challenging part of grad school and how did you handle it?

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?
- (3) How did you prepare for graduate school?
- (4) What was the most challenging part of grad school and how did you handle it?
- (5) Do you have any general advice for students considering pursuing the doctorate?

Q & A:

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?
- (3) How did you prepare for graduate school?
- (4) What was the most challenging part of grad school and how did you handle it?
- (5) Do you have any general advice for students considering pursuing the doctorate?

Contact: awheeler@mtholyoke.edu

Q & A:

- (1) What do you love most about mathematics?
- (2) What was your favorite math topic as an undergraduate?
- (3) How did you prepare for graduate school?
- (4) What was the most challenging part of grad school and how did you handle it?
- (5) Do you have any general advice for students considering pursuing the doctorate?

Contact: awheeler@mtholyoke.edu

Thank-you!

background from Euclidea