

Assignment 4 Solution

3.2 Using the Fourier series synthesis eq. (3.95)

$$\begin{aligned} x[n] &= a_0 + a_2 e^{j2(2\pi/N)n} + a_{-2} e^{-j2(2\pi/N)n} + a_4 e^{j4(2\pi/N)n} + a_{-4} e^{-j4(2\pi/N)n} \\ &= 1 + e^{j(\pi/4)} e^{j2(2\pi/5)n} + e^{-j(\pi/4)} e^{-j2(2\pi/5)n} \\ &= 1 + 2 \cos\left(\frac{4\pi}{5}n + \frac{\pi}{4}\right) + 4 \cos\left(\frac{8\pi}{5}n + \frac{\pi}{3}\right) \\ &= 1 + 2 \sin\left(\frac{4\pi}{5}n + \frac{3\pi}{4}\right) + 4 \sin\left(\frac{8\pi}{5}n + \frac{5\pi}{6}\right) \end{aligned}$$

3.27. Using the Fourier series synthesis eq.(3.38),

$$\begin{aligned} x[n] &= a_0 + a_2 e^{j2(2\pi/N)n} + a_{-2} e^{-j2(2\pi/N)n} + a_4 e^{j4(2\pi/N)n} + a_{-4} e^{-j4(2\pi/N)n} \\ &= 2 + 2e^{j\pi/6} e^{j(4\pi/5)n} + 2e^{-j\pi/6} e^{-j(4\pi/5)n} + e^{j\pi/3} e^{j(8\pi/5)n} + e^{-j\pi/3} e^{-j(8\pi/5)n} \\ &= 2 + 4 \cos[(4\pi n/5) + \pi/6] + 2 \cos[(8\pi n/5) + \pi/3] \\ &= 2 + 4 \sin[(4\pi n/5) + 2\pi/3] + 2 \sin[(8\pi n/5) + 5\pi/6] \end{aligned}$$

3.36. We will first evaluate the frequency response of the system. Consider an input $x[n]$ of the form $e^{j\omega n}$. From the discussion in Section 3.9 we know that the response to this input will be $y[n] = H(e^{j\omega}) e^{j\omega n}$. Therefore, substituting these in the given difference equation. We get

$$H(e^{j\omega}) e^{j\omega n} - \frac{1}{4} e^{-j\omega} e^{j\omega n} H(e^{j\omega}) = e^{j\omega n}.$$

Therefore,

$$H(j\omega) = \frac{1}{1 - \frac{1}{4} e^{-j\omega}}.$$

Form eq. (3.131), we know that coefficients

$$y[n] = \sum_{k=-\infty}^{\infty} a_k H(e^{j2\pi k/N}) e^{jk(2\pi/N)n}$$

when the input is $x[n]$. $x[n]$ has the Fourier series coefficients a_k

and fundamental frequency $2/N$. therefore, the Fourier series coefficients of $y[n]$ are $a_k H(e^{j2\pi k/N})$

(a) Here, $N = \frac{8}{3}$ and the nonzero FS coefficients of $x[n]$ are $a_3 = a_{-3}^* = 1/2j$. Therefore, the nonzero FS coefficients of $y[n]$ are

$$b_3 = a_3 H(e^{j3\pi/4}) = \frac{1}{2j(1 - (1/4)e^{-j3\pi/4})}, \quad b_{-3} = a_{-3} H(e^{-j3\pi/4}) = \frac{-1}{2j(1 - (1/4)e^{j3\pi/4})}.$$

Then the FS representation of $y[n]$ is

$$y[n] = \frac{1}{2j(1 - \frac{1}{4}e^{-j3\pi/4})} e^{j\frac{3\pi}{4}n} - \frac{1}{2j(1 - \frac{1}{4}e^{j3\pi/4})} e^{-j\frac{3\pi}{4}n}$$

(b) Here, $N=8$ and the nonzero FS coefficients of $x[n]$ are $a_1 = a_{-1} = 1/2$ and $a_2 = a_{-2} = 1$. Therefore, the nonzero FS coefficients of $y[n]$ are

$$\begin{aligned} b_1 &= a_1 H(e^{j\pi/4}) = \frac{1}{2(1 - (1/4)e^{-j\pi/4})}, & b_{-1} &= a_{-1} H(e^{-j\pi/4}) = \frac{1}{2(1 - (1/4)e^{j\pi/4})}, \\ b_2 &= a_2 H(e^{j\pi/2}) = \frac{1}{(1 - (1/4)e^{-j\pi/2})}, & b_{-2} &= a_{-2} H(e^{-j\pi/2}) = \frac{1}{(1 - (1/4)e^{j\pi/2})}. \end{aligned}$$

Then the FS representation of $y[n]$ is

$$\begin{aligned} y[n] &= \frac{1}{2 - \frac{1}{2}e^{-j\pi/4}} e^{j\frac{\pi}{4}n} + \frac{1}{2 - \frac{1}{2}e^{j\pi/4}} e^{-j\frac{\pi}{4}n} \\ &\quad + \frac{1}{1 - \frac{1}{4}e^{-j\pi/2}} e^{j\frac{\pi}{2}n} + \frac{1}{1 - \frac{1}{4}e^{j\pi/2}} e^{-j\frac{\pi}{2}n} \end{aligned}$$

3.38 The frequency response of the system may be evaluated as

$$H(e^{j\omega}) = \sum_{n=-\infty}^{+\infty} h[n]e^{-jn\omega} = 1 + e^{-j\omega} + e^{-2j\omega} - e^{j\omega} - e^{2j\omega} = 1 - 2j\sin(\omega) - 2j\sin(2\omega)$$

For $x[n]$, $N=4$ and $\omega_0 = \pi/2$. the FS coefficients of input $x[n]$ are $a_k = 1/4$, for all k

Therefore, the FS coefficients of output are

$$b_k = \frac{1}{4} [1 - 2j\sin(k\frac{\pi}{2}) - 2j\sin(k\pi)] = \frac{1}{4} [1 - 2j\sin(k\frac{\pi}{2})]$$

3.50

from Table 3.2, we know that if

$$x[n] \xleftrightarrow{FS} a_k,$$

then,

$$(-1)^n x[n] = e^{(2\pi/N)(N/2)n} x[n] \xleftrightarrow{FS} a_{k-N/2}$$

In this case, $N=8$. Therefore,

$$(-1)^n x[n] \xleftrightarrow{FS} a_{k-4}$$

This implies that $x[0] = x[\pm 2] = x[\pm 4] = \dots = 0$.

We are also given that $x[1] = x[5] = \dots$ and $x[3] = x[7] = -1$. therefore, one period of $x[n]$ is as shown in Figure S3.50




Figure S3.50