

# Notes

## Assignments

- 4.5
- 4.21 (b) (g) (h)
- 4.22 (c) (e)
- 4.27

## Tutorial problems

- Basic Problems with Answers 4.8, 4.9
- Basic Problems 4.23
- Advanced Problems 4.39, 4.40

## DT Fourier Series Pair $\left(\omega_o = \frac{2\pi}{N}\right)$

$$x[n] = \sum_{k=\langle N \rangle} a_k e^{jk\omega_o n} \quad (\text{Synthesis equation})$$

$$a_k = \frac{1}{N} \sum_{n=\langle N \rangle} x[n] e^{-jk\omega_o n} \quad (\text{Analysis equation})$$

$\sum_{k=\langle N \rangle}$  = Sum over *any*  $N$  consecutive values of  $k$

$$x[n] = x[n + N]$$

$$a_{k+N} = a_k$$

## LTI System, system function and frequency response

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t} \longrightarrow \boxed{h(t)} \longrightarrow y(t) = \sum_{k=-\infty}^{\infty} H(jk\omega_0) a_k e^{jk\omega_0 t}$$

$$a_k \longrightarrow \underbrace{H(jk\omega_0) a_k}_{\text{"gain"}}$$

$$H(j\omega) = \int_{-\infty}^{+\infty} h(t) e^{-j\omega t} dt$$

$$x[n] = \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 n} \longrightarrow \boxed{h[n]} \longrightarrow y[n] = \sum_{k=-\infty}^{\infty} H(e^{jk\omega_0}) a_k e^{jk\omega_0 n}$$


$$a_k \longrightarrow \underbrace{H(e^{jk\omega_0}) a_k}_{\text{"gain"}}$$

$$H(e^{j\omega}) = \sum_{n=-\infty}^{+\infty} h[n] e^{-j\omega n}$$


The effect of the LTI system is to modify each  $a_k$  through multiplication by the value of the frequency response at  $k\omega_0$ .

## Highpass Filters

CT



DT




Remember:

$$(-1)^n = e^{j\pi n}$$

—  $\pi$  = highest frequency in DT

# Chapter 4

## The Continuous-Time Fourier Transform




# Fourier Transform

- We have shown that Fourier series are useful in analyzing periodic signals, but many (most) signals are aperiodic. Need a more general tool — *Fourier transform*.


## Fourier's own derivation of the CT Fourier transform

- $x(t)$  — an aperiodic signal
  - view it as the limit of a periodic signal as  $T \rightarrow \infty$





## So, on the derivation of FT ...



For simplicity, assume  $x(t)$  has a finite duration, and introduce a periodic  $\tilde{x}(t)$

$$\tilde{x}(t) = \begin{cases} x(t) & -\frac{T}{2} < t < \frac{T}{2} \\ \text{periodic} & |t| > \frac{T}{2} \end{cases}$$

As  $T \rightarrow \infty$ ,  $x(t) = \tilde{x}(t)$  for all  $t$



## Derivation (cont.): Analysis equation


$$\tilde{x}(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t} \quad \omega_o = \frac{2\pi}{T}$$

$$a_k = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} \tilde{x}(t) e^{-jk\omega_0 t} dt = \frac{1}{T} \int_{-\frac{T}{2}}^{\frac{T}{2}} x(t) e^{-jk\omega_0 t} dt$$



$\tilde{x}(t) = x(t)$  in this interval

$$= \frac{1}{T} \int_{-\infty}^{+\infty} x(t) e^{-jk\omega_0 t} dt \quad (1)$$



If we define

$$X(j\omega) = \int_{-\infty}^{+\infty} x(t) e^{-j\omega t} dt$$

then, Eq. (1)  $\Rightarrow$

$$a_k = \frac{1}{T} X(jk\omega_0) = \frac{1}{T} X(j\omega) \big|_{\omega=k\omega_0}$$

# Derivation (cont.): **Synthesis equation**

Thus, for  $-\frac{T}{2} < t < \frac{T}{2}$

$$x(t) = \tilde{x}(t) = \sum_{k=-\infty}^{+\infty} \underbrace{\frac{1}{T} X(jk\omega_0)}_{a_k} e^{jk\omega_0 t}$$

$$= \frac{1}{2\pi} \sum_{k=-\infty}^{+\infty} \omega_0 X(jk\omega_0) e^{jk\omega_0 t}$$

$\Downarrow$

As  $T \rightarrow \infty$ ,  $\omega_0 \rightarrow 0$ ,  $\sum \omega_0 \rightarrow \int d\omega$ , and  $k\omega_0 = \omega$ , we get the CT **FT** pair

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(j\omega) e^{j\omega t} d\omega \quad \text{Synthesis equation}$$

— "sum" of  $e^{j\omega t}$

$$X(j\omega) = \int_{-\infty}^{+\infty} x(t) e^{-j\omega t} dt \quad \text{Analysis equation}$$

# The CT Fourier Transform Pair

$$X(j\omega) = \int_{-\infty}^{+\infty} x(t)e^{-j\omega t} dt \quad \text{— } FT$$

*Fourier Transform*

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(j\omega)e^{j\omega t} d\omega \quad \text{— Inverse } FT$$

*Inverse Fourier Transform*

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

$$\mathcal{F}(x(t)) = X(j\omega)$$

$$x(t) = \mathcal{F}^{-1}(X(j\omega))$$

# CT Fourier Transform Pair

$$X(j\omega) = \int_{-\infty}^{+\infty} x(t)e^{-j\omega t} dt$$

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(j\omega)e^{j\omega t} d\omega$$

# CT Fourier Series Pair

$$a_k = \frac{1}{T} \int_T x(t)e^{-jk\omega_0 t} dt$$

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t}$$

Harmonically related



# For what kinds of signals can we do FT?

It works also even if  $x(t)$  is infinite duration, but satisfies:

a) Finite energy  $\int_{-\infty}^{+\infty} |x(t)|^2 dt < \infty$

In this case, there is *zero* energy in the error

$$e(t) = x(t) - \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega \quad \text{Then} \quad \int_{-\infty}^{\infty} |e(t)|^2 dt = 0$$

## b) Dirichlet conditions

1) absolutely integrable  $\int_{-\infty}^{\infty} |x(t)| dt < \infty$


2) finite number of maxima and minima within any finite interval

3) finite number of discontinuities with finite values within any finite interval

(i)  $\frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega = x(t)$  at points of continuity

(ii)  $\frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega =$  midpoint at discontinuity

(iii) Gibb' s phenomenon



## Example 4.3 Impulse function

$$(a) \quad x(t) = \delta(t)$$

$$X(j\omega) = \int_{-\infty}^{+\infty} \delta(t) e^{-j\omega t} dt = ?$$

$\Downarrow$


$$\delta(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{j\omega t} d\omega \quad \text{— Synthesis equation for } \delta(t)$$

$$(b) \quad x(t) = \delta(t - t_0)$$

$$\begin{aligned} X(j\omega) &= \int_{-\infty}^{+\infty} \delta(t - t_0) e^{-j\omega t} dt \\ &= e^{-j\omega t_0} \quad \text{— Linear phase shift in } \omega \end{aligned}$$

### Example 4.4 A square pulse in the time-domain


$$X(j\omega) = \int_{-T_1}^{T_1} e^{-j\omega t} dt = \frac{2 \sin \omega T_1}{\omega}$$



Note the inverse relation between the two widths  $\Rightarrow$  **Uncertainty principle**

$$X(j\omega) = \int_{-\infty}^{+\infty} x(t) e^{-j\omega t} dt$$

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(j\omega) e^{j\omega t} d\omega$$



Useful facts about CTFT's


$$X(0) = \int_{-\infty}^{+\infty} x(t) dt$$

$$x(0) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(j\omega) d\omega$$





## Example 4.5 A square pulse in the frequency domain

$$X(j\omega) = \begin{cases} 1, & |\omega| < W \\ 0, & |\omega| > W \end{cases}$$



$$x(t) = \frac{1}{2\pi} \int_{-W}^W e^{j\omega t} d\omega = \frac{\sin(Wt)}{\pi t}$$



How about  $X(j\omega) = \delta(\omega)$  ?

# CT Fourier Transforms of **Periodic** Signals

Suppose

$$X(j\omega) = \delta(\omega - \omega_0)$$


$\Downarrow$

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \delta(\omega - \omega_0) e^{j\omega t} d\omega = \frac{1}{2\pi} e^{j\omega_0 t} \quad \text{— periodic in } t \text{ with frequency } \omega_0$$

That is

$$e^{j\omega_0 t} \longleftrightarrow 2\pi\delta(\omega - \omega_0)$$


— All the energy is concentrated in one frequency —  $\omega_0$



# Fourier Transform for Periodic Signals – Unified Framework

More generally, if  $x(t) = x(t+T)$ , then

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t} \longleftrightarrow X(j\omega) = \sum_{k=-\infty}^{+\infty} 2\pi a_k \delta(\omega - k\omega_0) \quad \text{Discrete spectra}$$




## Example 4.8

$$x(t) = \sum_{n=-\infty}^{+\infty} \delta(t - nT) \quad \text{— sampling function}$$


$$x(t) \xleftrightarrow{\text{FS}} a_k = \frac{1}{T} \int_{-T/2}^{T/2} x(t) e^{-jk\omega_o t} dt = \frac{1}{T}$$

$$\Downarrow x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_o t} = \sum_{k=-\infty}^{+\infty} \frac{1}{T} e^{jk\omega_o t}$$

$$X(j\omega) = \sum_{k=-\infty}^{+\infty} \frac{2\pi}{T} \delta(\omega - \underbrace{\frac{k2\pi}{T}}_{k\omega_o})$$



Same function in  
the frequency-domain!



Note in this case, periodic  
in both time domain (with  
a period  $T$ ) and frequency  
domain (with a period  
 $2\pi/T$ )

# Properties of the CT Fourier Transform

## 1) Linearity

$$x(t) \xleftrightarrow{F} X(j\omega), \quad y(t) \xleftrightarrow{F} Y(j\omega)$$

$$ax(t) + by(t) \xleftrightarrow{F} aX(j\omega) + bY(j\omega)$$

## 2) Time Shifting

$$x(t - t_0) \longleftrightarrow e^{-j\omega t_0} X(j\omega)$$

Proof: 
$$\int_{-\infty}^{\infty} x(\underbrace{t - t_0}_{t'}) e^{-j\omega t} dt = e^{-j\omega t_0} \underbrace{\int_{-\infty}^{\infty} x(t') e^{-j\omega t'} dt'}_{X(j\omega)}$$

FT magnitude unchanged

$$|e^{-j\omega_0 t} X(j\omega)| = |X(j\omega)|$$

Linear change in FT phase

$$\angle(e^{-j\omega_0 t} X(j\omega)) = \angle X(j\omega) - \omega t_0$$

# CTFT Properties (cont.)

## 3) Conjugation & Conjugate Symmetry

### - Conjugation

$$x^*(t) \xleftrightarrow{F} X^*(-j\omega)$$

### - Conjugate Symmetry

$$x(t) \text{ real} \longleftrightarrow X(-j\omega) = X^*(j\omega)$$

$$|X(-j\omega)| = |X(j\omega)|$$

*Even*

Or

$$\text{Re}\{X(-j\omega)\} = \text{Re}\{X(j\omega)\}$$

*Even*

$$\angle X(-j\omega) = -\angle X(j\omega)$$

*Odd*

$$\text{Im}\{X(-j\omega)\} = -\text{Im}\{X(j\omega)\}$$

*Odd*

When  $x(t)$  is real (all the physically measurable signals are *real*), the negative frequency components do *not* carry any additional information from the positive frequency components.  $\omega \geq 0$  will be sufficient.

## CT Fourier Series Property

- Conjugate Symmetry

$$x(t) \text{ real} \Rightarrow a_{-k} = a_k^*$$

Proof:

$$a_{-k} = \frac{1}{T} \int_T x(t) e^{jk\omega_0 t} dt = \left[ \frac{1}{T} \int_T x^*(t) e^{-jk\omega_0 t} dt \right]^* = a_k^*$$

$$\therefore a_k = \text{Re}\{a_k\} + j\text{Im}\{a_k\}$$


$$\therefore \boxed{\text{Re}\{a_{-k}\} + j\text{Im}\{a_{-k}\}} = \boxed{\text{Re}\{a_k\} - j\text{Im}\{a_k\}}$$

$$\therefore \text{Re}\{a_k\} \text{ is even, } \text{Im}\{a_k\} \text{ is odd}$$

# CTFT Properties (cont.)

## 4) Time/Frequency Scaling $x(at) \longleftrightarrow \frac{1}{|a|} X\left(j\frac{\omega}{a}\right)$

E.g.  $a > 1 \rightarrow at > t$   
compressed in time  $\leftrightarrow$   
stretched in frequency





# CTFT Properties (cont.)

## 4) Time/Frequency Scaling

$$x(at) \longleftrightarrow \frac{1}{|a|} X\left(j\frac{\omega}{a}\right)$$

*E.g.  $a > 1 \rightarrow at > t$   
compressed in time  $\leftrightarrow$   
stretched in frequency*

$$\Downarrow a = -1$$

$$x(-t) \longleftrightarrow X(-j\omega)$$

Time reversal

$$\Downarrow$$

a)  $x(t)$  real and even

$$x(t) = x(-t) = x^*(t)$$

$$\Rightarrow X(j\omega) = X(-j\omega) = X^*(-j\omega) \text{ — Real \& even}$$

b)  $x(t)$  real and odd

$$x(t) = -x(-t) = x^*(t)$$

$$\Rightarrow X(j\omega) = -X(-j\omega) = X^*(-j\omega) \text{ — Purely imaginary \& odd}$$

$$c) \quad X(j\omega) = \operatorname{Re}\{X(j\omega)\} + j\operatorname{Im}\{X(j\omega)\}$$



For real

$$x(t) = \operatorname{Ev}\{x(t)\} + \operatorname{Od}\{x(t)\}$$

# CTFT Properties (cont.)

## 5) Differentiation/Integration

$$\frac{dx(t)}{dt} \xleftrightarrow{F} j\omega X(j\omega)$$

$$\int_{-\infty}^t x(\tau) d\tau \xleftrightarrow{F} \frac{1}{j\omega} X(j\omega) + \pi X(j0) \delta(\omega)$$

$\uparrow$   
 DC term


### Example:

What is the Fourier transform for unit step function  $u(t)$ ?


# CTFT Properties (cont.)

## 6) Duality

$$x_1(t) = \begin{cases} 1, & |t| < T_1 \\ 0, & |t| > T_1 \end{cases}$$




$\mathcal{F}$




$$X_1(j\omega) = \frac{2 \sin \omega T_1}{\omega}$$

$$x_2(t) = \frac{\sin Wt}{\pi t}$$



$\mathcal{F}$




$$X_2(j\omega) = \begin{cases} 1, & |\omega| < W \\ 0, & |\omega| > W \end{cases}$$

- Time/frequency domains are kind of “symmetric”.
- If there are characteristics of a function of time that have implications with regard to the Fourier transform, then the same characteristics associated with a function of frequency will have *dual* implications in the time domain.

### Example:

$$\{\delta(t - t_k), -\infty < t_k < \infty\} \quad \{2\pi\delta(\omega - \omega_k), -\infty < \omega_k < \infty\}$$



## CTFT Properties (cont.)

### 7) Parseval's Relation

$$\underbrace{\int_{-\infty}^{+\infty} |x(t)|^2 dt}_{\text{Total energy in the time-domain}} = \underbrace{\frac{1}{2\pi} \int_{-\infty}^{+\infty} |X(j\omega)|^2 d\omega}_{\text{Total energy in the frequency-domain}}$$

Total energy in the  
time-domain

Total energy in the  
frequency-domain

$$\frac{1}{2\pi} |X(j\omega)|^2$$

— spectral density

# Table 4.2

## Basic Fourier Transform Pairs

Signal	Fourier transform	Fourier series coefficients (if periodic)
$\sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t}$	$2\pi \sum_{k=-\infty}^{+\infty} a_k \delta(\omega - k\omega_0)$	$a_k$
$e^{j\omega_0 t}$	$2\pi \delta(\omega - \omega_0)$	$a_1 = 1$ $a_k = 0$ , otherwise
$\cos \omega_0 t$	$\pi[\delta(\omega - \omega_0) + \delta(\omega + \omega_0)]$	$a_1 = a_{-1} = \frac{1}{2}$ $a_k = 0$ , otherwise
$\sin \omega_0 t$	$\frac{\pi}{j}[\delta(\omega - \omega_0) - \delta(\omega + \omega_0)]$	$a_1 = -a_{-1} = \frac{1}{2j}$ $a_k = 0$ , otherwise
$x(t) = 1$	$2\pi \delta(\omega)$	$a_0 = 1$ , $a_k = 0$ , $k \neq 0$ (this is the Fourier series representation for any choice of $T > 0$ )
Periodic square wave		
$x(t) = \begin{cases} 1, &  t  < T_1 \\ 0, & T_1 <  t  \leq \frac{T}{2} \end{cases}$ and $x(t + T) = x(t)$	$\sum_{k=-\infty}^{+\infty} \frac{2 \sin k\omega_0 T_1}{k} \delta(\omega - k\omega_0)$	$\frac{\omega_0 T_1}{\pi} \operatorname{sinc}\left(\frac{k\omega_0 T_1}{\pi}\right) = \frac{\sin k\omega_0 T_1}{k\pi}$
$\sum_{n=-\infty}^{+\infty} \delta(t - nT)$	$\frac{2\pi}{T} \sum_{k=-\infty}^{+\infty} \delta\left(\omega - \frac{2\pi k}{T}\right)$	$a_k = \frac{1}{T}$ for all $k$
$x(t) \begin{cases} 1, &  t  < T_1 \\ 0, &  t  > T_1 \end{cases}$	$\frac{2 \sin \omega T_1}{\omega}$	—
$\frac{\sin Wt}{\pi t}$	$X(j\omega) = \begin{cases} 1, &  \omega  < W \\ 0, &  \omega  > W \end{cases}$	—
$\delta(t)$	1	—
$u(t)$	$\frac{1}{j\omega} + \pi \delta(\omega)$	—
$\delta(t - t_0)$	$e^{-j\omega t_0}$	—
$e^{-at} u(t), \operatorname{Re}\{a\} > 0$	$\frac{1}{a + j\omega}$	—
$te^{-at} u(t), \operatorname{Re}\{a\} > 0$	$\frac{1}{(a + j\omega)^2}$	—
$\frac{t^{n-1}}{(n-1)!} e^{-at} u(t), \operatorname{Re}\{a\} > 0$	$\frac{1}{(a + j\omega)^n}$	—

# Summary

- **Understand CT Fourier transform**
  - ◆ **Synthesis and analysis equations**
  - ◆ **Difference with CT Fourier series**
  - ◆ **Fourier transform for periodic signal**
  - ◆ **Properties of CT Fourier transform**