

Generic Degree Three in $JC(3)$: Collision Geometry and Galois Group S_3

Chloe van der Vlugt

August 24, 2026

Abstract

We study possible counterexamples to the complex three-dimensional Jacobian conjecture. For a polynomial map $F: \mathbb{A}_{\mathbb{C}}^n \rightarrow \mathbb{A}_{\mathbb{C}}^n$, let R_F be the scheme of pairs (x, y) satisfying $F(x) = F(y)$, and let C_F be its coordinate ring. The pairs (x, x) form a closed diagonal in R_F , whose defining ideal in C_F we denote by $\text{Obs}(F)$. The coordinate functions of F generate

$$B = \mathbb{C}[F_1, \dots, F_n] \subseteq A = \mathbb{C}[x_1, \dots, x_n],$$

and hence a function-field extension $K = \text{Frac}(B) \subseteq L = \text{Frac}(A)$. We prove that every three-dimensional Keller map with $[L : K] = 3$ is a counterexample to $JC(3)$.

For every such map, the generic collision algebra decomposes as

$$K \otimes_B C_F \simeq_K L \times N,$$

where N is the normal closure of L/K . Consequently, the normal closure of the function-field extension attached to every generic-degree-three counterexample to $JC(3)$ has Galois group S_3 .

Licensed under [CC BY 4.0](#).

Contents

1	Introduction	2
2	General construction	3
2.1	The two ideals and their obstruction	3
2.2	The fiber-product interpretation	4
2.3	Generic fibers and marked-root decomposition	5
2.4	Automorphism detection	6
3	The cubic S_3 obstruction in dimension three	8
3.1	The cubic extension and its normal closure	8
4	Further directions	11
5	Conclusion	12

1 Introduction

The complex Jacobian conjecture $JC(n)$ asks whether every polynomial map

$$F = (F_1, \dots, F_n): \mathbb{A}_{\mathbb{C}}^n \longrightarrow \mathbb{A}_{\mathbb{C}}^n, \quad \det JF \in \mathbb{C}^\times,$$

has a polynomial inverse. Such an F is called a Keller map. The Jacobian condition makes F an étale morphism; the conjecture asks whether this local condition forces global invertibility.

The equations

$$F_i(x) - F_i(y) = 0 \quad (1 \leq i \leq n)$$

define a closed collision scheme

$$R_F \subseteq \mathbb{A}_{\mathbb{C}}^n \times \mathbb{A}_{\mathbb{C}}^n.$$

Its complex points are the pairs (x, y) with $F(x) = F(y)$; equivalently, R_F is the fiber square of F . It contains the diagonal Δ , consisting of the pairs (x, x) . If C_F is the coordinate ring of R_F , restriction to the diagonal gives

$$\bar{\mu}_F: C_F \longrightarrow A, \quad A = \mathbb{C}[x_1, \dots, x_n],$$

and we write $\text{Obs}(F) = \ker(\bar{\mu}_F)$. By Ax–Grothendieck,

$$\text{Obs}(F) = 0 \iff R_F = \Delta \iff F \text{ is a polynomial automorphism.}$$

A possible counterexample must therefore remain globally noninvertible despite being étale. We organize this failure by generic degree. On coordinate rings, F induces

$$F^*: \mathbb{C}[t_1, \dots, t_n] \longrightarrow A, \quad t_i \longmapsto F_i.$$

Let

$$B = \text{im}(F^*) = \mathbb{C}[F_1, \dots, F_n] \subseteq A, \quad K = \text{Frac}(B), \quad L = \text{Frac}(A).$$

For a Keller map, the F_i are algebraically independent, so F^* identifies the target coordinate ring with B . Thus $K = \mathbb{C}(F_1, \dots, F_n) \subseteq L = \mathbb{C}(x_1, \dots, x_n)$ is the induced finite separable extension of function fields, and $[L : K]$ is the generic degree. If L/K is normal, the classical Galois case of the Jacobian conjecture implies that F is a polynomial automorphism. Since extensions of degree one and separable extensions of degree two are normal, neither degree can occur for a counterexample. Generic degree three is therefore the first possible counterexample degree.

We henceforth study polynomial maps $F: \mathbb{A}_{\mathbb{C}}^3 \rightarrow \mathbb{A}_{\mathbb{C}}^3$ with generic degree $[L : K] = 3$. Here $JC(3)$ denotes the conjecture in three variables, while $[L : K] = 3$ is the separate generic-degree condition. Since polynomial automorphisms have generic degree one, every map with $[L : K] = 3$ is already nonautomorphic. In this generic-degree-three case,

$$F \text{ is a counterexample to } JC(3) \iff F \text{ is Keller.}$$

The residual decomposition itself is field-theoretic and does not use the Keller condition. For a map of generic degree three, the generic collision algebra has a diagonal factor L and a quadratic residual factor D_α :

$$K \otimes_B C_F \simeq_K L \times D_\alpha.$$

The quadratic residual factor is split in the normal branch and is a field in the nonnormal branch. In the latter case it is L -isomorphic to the normal closure N . Galois rigidity excludes the normal branch for Keller maps, so every generic-degree-three counterexample to $JC(3)$ has

$$K \otimes_B C_F \simeq_K L \times N, \quad \text{Gal}(N/K) \cong S_3.$$

Section 2 develops the dimension-independent collision obstruction, generic base change, marked-root decomposition, and automorphism criterion. Section 3 specializes these constructions to cubic extensions. Section 4 isolates the normalization and boundary questions leading to the planar sequel. The closing low-degree corollary records the precise handoff to that paper.

2 General construction

For a fixed dimension n , where can $JC(n)$ fail? To answer this, we attach to an arbitrary polynomial map the scheme of pairs with the same image and the ideal defining its diagonal. A Keller map is a counterexample to $JC(n)$ precisely when this ideal is nonzero. The construction itself is dimension-independent and does not require the Keller condition.

2.1 The two ideals and their obstruction

Throughout the paper, the base field is $\mathbb{C}$. Let

$$F = (F_1, \dots, F_n): X = \mathbb{A}_{\mathbb{C}}^n \longrightarrow \mathbb{A}_{\mathbb{C}}^n$$

be a polynomial map, and let

$$A = \mathbb{C}[x_1, \dots, x_n], \quad S = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n].$$

The map F is Keller if

$$\det JF \in \mathbb{C}^\times.$$

Define

$$I_R(F) = (F_i(x) - F_i(y))_{i=1}^n, \quad I_\Delta = (x_i - y_i)_{i=1}^n, \quad C_F := S/I_R(F), \quad R_F := \text{Spec } C_F.$$

For a fixed map F , we abbreviate $I_R(F)$ to I_R .

Proposition 2.1 (Canonical containment). *For every polynomial map F ,*

$$I_R(F) \subseteq I_\Delta.$$

Proof. For $P \in \mathbb{C}[x_1, \dots, x_n]$, the difference $P(x) - P(y)$ belongs to $(x_1 - y_1, \dots, x_n - y_n)$. This follows by induction on P , or by telescoping each monomial one coordinate at a time. Apply the statement to each coordinate F_i . $\square$

Let $q_F: S \rightarrow C_F$ be the quotient map. Diagonal substitution $S \rightarrow A$, given by $s(x, y) \mapsto s(x, x)$, has kernel I_Δ . Since $I_R \subseteq I_\Delta$, it descends uniquely to a surjection

$$\bar{\mu}_F: C_F \rightarrow A, \quad [s]_{I_R} \mapsto s(x, x).$$

We define the ideal of the diagonal in R_F by

$$\text{Obs}(F) := q_F(I_\Delta) = I_\Delta/I_R \subseteq C_F.$$

Proposition 2.2 (The diagonal kernel). *As ideals of C_F ,*

$$\ker(\bar{\mu}_F) = \text{Obs}(F) = I_\Delta/I_R.$$

Equivalently,

$$\ker(\bar{\mu}_F) = \text{Obs}(F) = 0 \iff I_R = I_\Delta.$$

Proof. The construction of $\bar{\mu}_F$ identifies its kernel with $I_\Delta/I_R = \text{Obs}(F)$. Since $I_R \subseteq I_\Delta$,

$$I_\Delta/I_R = 0 \iff I_R = I_\Delta.$$

□

2.2 The fiber-product interpretation

Let

$$A = \mathbb{C}[x_1, \dots, x_n], \quad B = \mathbb{C}[F_1, \dots, F_n] \subseteq A.$$

Let $X = \text{Spec } A$, let $Y = \text{Spec } B$, and let

$$f_F: X \longrightarrow Y$$

be the morphism induced by $B \hookrightarrow A$. The homomorphisms $\iota_x, \iota_y: A \rightrightarrows C_F$, defined by $\iota_x(a) = [a(x)]$ and $\iota_y(a) = [a(y)]$, agree on B because $[F_i(x)] = [F_i(y)]$ in C_F . The equality $\iota_x|_B = \iota_y|_B$ defines the B -algebra structure on C_F .

Theorem 2.3 (Fiber-product interpretation). *There is a canonical isomorphism*

$$R_F \cong X \times_Y X$$

under which $\text{Spec}(\bar{\mu}_F): X \rightarrow R_F$ is the relative diagonal $\Delta_{X/Y}$. Dually, there is a canonical B -algebra isomorphism

$$C_F \xrightarrow{\sim}_B A \otimes_B A$$

under which $\bar{\mu}_F$ is tensor multiplication $a \otimes a' \mapsto aa'$.

Proof. For every commutative $\mathbb{C}$ -algebra T , maps from either C_F or $A \otimes_B A$ to T are pairs $f, g: A \rightrightarrows T$ agreeing on B . In the presentation of C_F , this agreement is imposed by the relations $F_i(x) = F_i(y)$, since the F_i generate B . The common universal property gives the algebra isomorphism, and applying Spec gives the scheme isomorphism. Tensor multiplication corresponds to $\bar{\mu}_F$, so its dual is the relative diagonal. □

On complex points,

$$R_F(\mathbb{C}) = \{(u, v) : F(u) = F(v)\}.$$

Inside $X \times X = \text{Spec } S$, the ideals I_R and I_Δ define $X \times_Y X$ and its diagonal. Thus

$$I_R = I_\Delta \iff X \times_Y X = \Delta_{X/Y}(X)$$

as closed subschemes. This is the geometric form of obstruction vanishing.

2.3 Generic fibers and marked-root decomposition

Let F be a polynomial map. Retain A , and set

$$B = \mathbb{C}[F_1, \dots, F_n] \subseteq A.$$

Let

$$K = \text{Frac}(B), \quad L = \text{Frac}(A).$$

Proposition 2.4 (Keller generic geometry). *If F is Keller, then its coordinate functions are algebraically independent, the morphism*

$$f_F: \text{Spec } A \longrightarrow \text{Spec } B$$

is étale and quasi-finite, and L/K is a finite separable extension. Consequently, every Keller map, and in particular every complex Jacobian-conjecture counterexample, has a finite separable generic field extension.

Proof. The identity

$$dF_1 \wedge \dots \wedge dF_n = \det(JF) dx_1 \wedge \dots \wedge dx_n \neq 0$$

gives algebraic independence, so $B \cong \mathbb{C}[u_1, \dots, u_n]$. As a B -algebra, A has the presentation

$$A \cong \mathbb{C}[u_1, \dots, u_n, X_1, \dots, X_n] / (F_i(X) - u_i)_{i=1}^n.$$

Its relative Jacobian determinant is the unit $\det(JF)$. The Jacobian criterion therefore makes f_F étale, hence quasi-finite. Since $B \subseteq A$, the morphism is dominant, so quasi-finiteness gives $[L : K] < \infty$. Its generic point is étale, and therefore L/K is separable. $\square$

For the remainder of this subsection, assume that F is dominant and generically finite. Let

$$\theta: K \otimes_B A \longrightarrow L, \quad \lambda \otimes a \longmapsto \lambda a.$$

Let

$$\mu_L: L \otimes_K L \longrightarrow L, \quad \ell \otimes \ell' \longmapsto \ell \ell',$$

denote tensor multiplication.

Theorem 2.5 (Generic collision bridge). *The map θ is an isomorphism. Consequently, there is a K -algebra isomorphism*

$$\Phi_F: K \otimes_B C_F \xrightarrow{\sim} L \otimes_K L,$$

and the following square commutes:

$$\begin{array}{ccc} K \otimes_B C_F & \xrightarrow{1 \otimes \bar{\mu}_F} & K \otimes_B A \\ \Phi_F \downarrow & & \downarrow \theta \sim \\ L \otimes_K L & \xrightarrow{\mu_L} & L. \end{array}$$

Proof. Let $U = B \setminus \{0\}$. Since $K = U^{-1}B$, localization gives

$$K \otimes_B A \cong U^{-1}A \subseteq L,$$

and under this identification θ is the inclusion. Since L/K is finite, each x_i is algebraic over K ; hence

$$U^{-1}A = K[x_1, \dots, x_n]$$

is a finite-dimensional domain over K , and therefore a field. It contains A , so it contains $\text{Frac}(A) = L$. Thus $U^{-1}A = L$, and θ is an isomorphism.

Using Theorem 2.3 and standard tensor base change, define Φ_F by the composite

$$K \otimes_B C_F \cong K \otimes_B (A \otimes_B A) \cong (K \otimes_B A) \otimes_K (K \otimes_B A) \xrightarrow{\theta \otimes \theta} L \otimes_K L.$$

Under this composite, both routes around the square send $\lambda \otimes (a \otimes a')$ to $\lambda a a'$. Hence the square commutes. $\square$

Suppose now that L/K is separable and has a power basis $L = K(\alpha)$. Let $m_\alpha(T) \in K[T]$ be the minimal polynomial of α , and let $h_\alpha(T) \in L[T]$ be determined by

$$m_\alpha(T) = (T - \alpha)h_\alpha(T) \quad \text{in } L[T].$$

Set

$$D_\alpha := L[T]/(h_\alpha).$$

Theorem 2.6 (Marked-root tensor decomposition). *Give $L \otimes_K L$ its L -algebra structure through the left tensor factor. There is an L -algebra isomorphism*

$$\Phi_\alpha: L \otimes_K L \xrightarrow{\sim}_L L \times D_\alpha,$$

under which μ_L is the first-coordinate map.

Proof. View m_α in $L[T]$. Separability gives $h_\alpha(\alpha) = m'_\alpha(\alpha) \neq 0$, so $T - \alpha$ and h_α are coprime. The power-basis isomorphism and the Chinese remainder theorem give the composite

$$L \otimes_K L \xrightarrow{\sim}_L L[T]/(m_\alpha) \xrightarrow{\sim}_L L[T]/(T - \alpha) \times L[T]/(h_\alpha) \xrightarrow{\sim}_L L \times D_\alpha.$$

The first coordinate of the composite is evaluation at $T = \alpha$, which is precisely μ_L . $\square$

Together with Theorem 2.5, this gives

$$K \otimes_B C_F \simeq_K L \times D_\alpha,$$

with the base-changed diagonal evaluation equal to the first-coordinate map.

2.4 Automorphism detection

Theorem 2.7 (Automorphism criterion). *For an arbitrary polynomial map $F: \mathbb{A}_{\mathbb{C}}^n \rightarrow \mathbb{A}_{\mathbb{C}}^n$, the following are equivalent:*

- (1) F is a polynomial automorphism;
- (2) F is injective on $\mathbb{A}^n(\mathbb{C})$;
- (3) $I_R(F) = I_\Delta$;

(4) $\text{Obs}(F) = 0$.

Proof. Suppose first that F has polynomial inverse $G = (G_1, \dots, G_n)$. Then

$$x_i - y_i = G_i(F(x)) - G_i(F(y)) \in I_R$$

for every i , so $I_\Delta \subseteq I_R$. Proposition 2.1 gives equality. Thus (1) implies (3).

Suppose $I_R = I_\Delta$, and let $\text{ev}_{(a,b)}: S \rightarrow \mathbb{C}$ be evaluation at (a, b) . If $F(a) = F(b)$, then

$$I_R \subseteq \ker(\text{ev}_{(a,b)}).$$

Therefore $x_i - y_i \in I_\Delta = I_R$ gives $a_i - b_i = 0$ for every i . Thus $a = b$, proving (3) $\Rightarrow$ (2). The Ax–Grothendieck theorem [Ax69] gives (2) $\Rightarrow$ (1). Finally, Proposition 2.2 gives (3) $\iff$ (4). $\square$

Applied to Keller maps, Theorem 2.7 says that $JC(n)$ is equivalent to universal vanishing of $\text{Obs}(F)$, or equivalently to $I_R(F) = I_\Delta$.

We will also use the classical Galois case of the conjecture. It is the step that rules out the normal cubic branch under the Keller hypothesis.

Theorem 2.8 (Classical Galois rigidity). *Let*

$$F: \mathbb{A}_{\mathbb{C}}^n \longrightarrow \mathbb{A}_{\mathbb{C}}^n, \quad \det JF \in \mathbb{C}^\times,$$

and put

$$K = \mathbb{C}(F_1, \dots, F_n), \quad L = \mathbb{C}(x_1, \dots, x_n).$$

If L/K is normal, then F is a polynomial automorphism and $L = K$.

Proof. The Keller condition makes L/K finite and separable. Thus normality makes it Galois, and the conclusion is the classical Galois case of the Jacobian conjecture due to Campbell, with algebraic treatments by Razar and Wright [Cam73, Raz79, Wri81]. $\square$

Lean correspondence. `CollisionIdeals/General/Collision/Diagonal.lean`,
`CollisionIdeals/General/Collision/Ideal.lean`,
`CollisionIdeals/General/Collision/CollisionDiagonal.lean`,
`CollisionIdeals/General/FiberProduct/Polynomial.lean`,
`CollisionIdeals/General/GenericFiber/FunctionField.lean`,
`CollisionIdeals/General/GenericFiber/CollisionDiagonal.lean`,
`CollisionIdeals/General/GenericFiber/MarkedRootDecomposition.lean`,
`CollisionIdeals/General/Keller/Interfaces.lean`,
`CollisionIdeals/General/Automorphism/Equivalences.lean`, and
`CollisionIdeals/General/Automorphism/GaloisRigidity.lean`. The polynomial fiber-product module uses the abstract target polynomial ring; the equivalence over the actual image algebra $B = \mathbb{C}[F_1, \dots, F_n]$ and its generic base change are owned by the generic-function-field module. The Lean form of Theorem 2.5 takes surjectivity of θ as a parameter; the proof above derives it from generic finiteness. The predicates `KellerEtaleBridge` and `KellerFlatBridge` expose parts of Proposition 2.4 as theorem targets; Lean does not yet derive them, generic finiteness, or separability from `IsKeller`. Ax–Grothendieck also remains an explicit parameter of the Lean automorphism equivalence.

3 The cubic S_3 obstruction in dimension three

We now specialize the dimension-independent constructions of Section 2 to generic degree three. After the diagonal is exhibited as the first factor, the complementary generic collision algebra has degree two over L . For a map of generic degree three it is either split, corresponding to a normal cubic extension, or connected, corresponding to a nonnormal cubic extension. Degree three already rules out polynomial automorphy, so the counterexamples to $J\mathcal{C}(3)$ in generic degree three are precisely the Keller maps. For a Keller map, however, the split branch is impossible: it makes L/K normal and hence, by Theorem 2.8, forces F to be a polynomial automorphism, contradicting $[L : K] = 3$. Hence only the connected branch remains; its residual factor is the normal closure, with Galois group S_3 .

3.1 The cubic extension and its normal closure

Let

$$F = (F_1, F_2, F_3) : \mathbb{A}_{\mathbb{C}}^3 \longrightarrow \mathbb{A}_{\mathbb{C}}^3$$

be dominant and generically finite. Let

$$A = \mathbb{C}[x_1, x_2, x_3], \quad B = \mathbb{C}[F_1, F_2, F_3] \subseteq A, \quad K = \text{Frac}(B), \quad L = \text{Frac}(A).$$

Assume that F has generic degree three:

$$[L : K] = 3.$$

Because K has characteristic zero, L/K is separable. Choose a primitive element α [The26e], so that $L = K(\alpha)$ with power basis $1, \alpha, \alpha^2$. Retain the notation of Theorem 2.6: if $m_\alpha \in K[T]$ is the minimal polynomial of α , then

$$m_\alpha(T) = (T - \alpha)h_\alpha(T) \quad \text{in } L[T].$$

The residual factor in the marked-root decomposition is

$$D_\alpha = L[T]/(h_\alpha).$$

Since $[L : K] = 3$, it has dimension two over L . Let N/K be the normal closure of L/K , containing L through the marked inclusion. Since L/K is finite separable, N/K is Galois [The26c]. Let

$$G = \text{Gal}(N/K), \quad H = \text{Gal}(N/L) = \{g \in G : g|_L = \text{id}_L\},$$

where H fixes the marked copy of L in N . Restriction of automorphisms gives a bijection

$$G/H \xrightarrow{\sim} \text{Hom}_K(L, N), \quad gH \longmapsto g|_L.$$

Because the conjugates of L generate its normal closure N , an element of G fixing every conjugate embedding fixes N . Hence the coset action is faithful and

$$\text{core}_G(H) = \bigcap_{g \in G} gHg^{-1} = 1.$$

The residual collision algebra detects which cubic branch occurs. We include the short argument in order to fix its marked identification with the normal closure.

Proposition 3.1 (Connected cubic residual factor). *The following conditions are equivalent:*

- (i) $\text{Spec } D_\alpha$ is connected;
- (ii) D_α is a field;
- (iii) h_α is irreducible over L ;
- (iv) L/K is nonnormal.

When they hold, there is an L -algebra isomorphism

$$\varphi_\alpha: D_\alpha \xrightarrow{\sim}_L N.$$

Consequently,

$$\Psi_\alpha: L \otimes_K L \xrightarrow{\sim}_L L \times N,$$

under which μ_L is the first-coordinate map.

Proof. Write

$$m_\alpha(T) = T^3 + a_2T^2 + a_1T + a_0.$$

The quadratic h_α is separable, so D_α is finite étale over L ; finite étale algebras over a field are finite products of finite separable field extensions [The26a]. If h_α is irreducible, then D_α is a field and hence has connected spectrum. If it is reducible, it is a product of two distinct linear factors; the Chinese remainder theorem then gives $D_\alpha \simeq_L L \times L$, whose spectrum is disconnected. This proves the equivalence of (i), (ii), and (iii).

If h_α is reducible, it has a root $\beta \in L$. Separability gives $\beta \neq \alpha$, and the third root is

$$\gamma = -a_2 - \alpha - \beta \in L.$$

Thus m_α would split over L . Since $L = K(\alpha)$, the standard generator criterion for normality [The26d] shows that L/K would be normal. Conversely, if L/K is normal, then m_α splits over L , so h_α is reducible. Hence (iii) and (iv) are equivalent.

Assume these equivalent conditions, and let β denote the image of T in D_α . This field contains the three roots

$$\alpha, \quad \beta, \quad -a_2 - \alpha - \beta$$

of m_α , and is generated over L by β . It is therefore the splitting field of m_α , which is the normal closure N of L/K [The26f]. Choose the resulting L -algebra isomorphism

$$\varphi_\alpha: D_\alpha \xrightarrow{\sim}_L N.$$

Let

$$\Phi_\alpha: L \otimes_K L \xrightarrow{\sim}_L L \times D_\alpha$$

be the marked-root decomposition of Theorem 2.6. Under this isomorphism, μ_L is the first-coordinate map. Composing its second factor with φ_α , let

$$\Psi_\alpha = (\text{id}_L \times \varphi_\alpha) \circ \Phi_\alpha.$$

Since $\text{id}_L \times \varphi_\alpha$ leaves the first factor unchanged, the same description of μ_L holds under Ψ_α . $\square$

Corollary 3.2 (The generic-degree-three counterexample locus). *The map F is not a polynomial automorphism. Equivalently, on this generic-degree-three locus,*

$$F \text{ is a counterexample to } JC(3) \iff F \text{ is Keller.}$$

When these equivalent conditions hold, L/K is nonnormal and $\text{Spec } D_\alpha$ is connected.

Proof. If F were a polynomial automorphism, its coordinate functions would generate A , so $B = A$ and $L = K$, contrary to $[L : K] = 3$. The displayed equivalence now follows from the definition of a counterexample to $JC(3)$.

Assume these conditions, so that F is Keller. If L/K were normal, Theorem 2.8 would give $L = K$, again contradicting $[L : K] = 3$. Proposition 3.1 now identifies nonnormality with connectedness of $\text{Spec } D_\alpha$. $\square$

Theorem 3.3 (Generic cubic S_3 collision). *Assume that $\text{Spec } D_\alpha$ is connected. There is a K -algebra isomorphism*

$$\Psi_F: K \otimes_B C_F \xrightarrow{\sim}_K L \times N$$

under which, after the identification $K \otimes_B A \simeq_K L$ of Theorem 2.5, the base-changed diagonal evaluation

$$1 \otimes \bar{\mu}_F: K \otimes_B C_F \longrightarrow K \otimes_B A$$

is the first-coordinate map. In particular,

$$\Psi_F(\ker(1 \otimes \bar{\mu}_F)) = 0 \times N, \quad \text{Gal}(N/K) \cong S_3,$$

and

$$\text{Obs}(F) \neq 0, \quad I_R \subsetneq I_\Delta.$$

By Corollary 3.2, the connectedness hypothesis holds whenever F is Keller; in generic degree three these Keller maps are precisely the counterexamples to $JC(3)$.

Proof. Proposition 3.1 gives an L -algebra isomorphism

$$\Psi_\alpha: L \otimes_K L \xrightarrow{\sim}_L L \times N$$

under which μ_L is the first-coordinate map. Theorem 2.5 gives a K -algebra isomorphism

$$\Phi_F: K \otimes_B C_F \xrightarrow{\sim}_K L \otimes_K L$$

which carries the base-changed diagonal evaluation to μ_L . After restricting Ψ_α from L to K , let

$$\Psi_F = \Psi_\alpha \circ \Phi_F.$$

The asserted description of diagonal evaluation follows, and hence

$$\Psi_F(\ker(1 \otimes \bar{\mu}_F)) = 0 \times N \neq 0.$$

If $\bar{\mu}_F: C_F \rightarrow A$ were injective, flat base change along the localization $B \rightarrow K$ would make $1 \otimes \bar{\mu}_F$ injective. This would contradict the displayed kernel. Thus $\ker(\bar{\mu}_F) \neq 0$, so Proposition 2.2 gives $\text{Obs}(F) \neq 0$, and Proposition 2.1 gives $I_R \subsetneq I_\Delta$.

For the Galois group, Proposition 3.1 identifies N with the degree-two field D_α over L . The fixed-field degree formula therefore gives

$$|H| = [N : L] = 2, \quad [G : H] = [L : K] = 3.$$

The normal-closure identities recalled above give $\text{core}_G(H) = 1$. Thus the action on the three cosets of H embeds G into S_3 . But

$$|G| = [G : H]|H| = 3 \cdot 2 = 6,$$

so this embedding is an isomorphism $G \cong S_3$. We use here the fixed-field degree formula from the fundamental theorem of Galois theory [The26b]. $\square$

Remark (Ordered roots). Let

$$m_\alpha(T) = T^3 + a_2T^2 + a_1T + a_0$$

have roots $\alpha_1, \alpha_2, \alpha_3$ in N . For distinct i, j, k ,

$$\alpha_k = -a_2 - \alpha_i - \alpha_j, \quad K(\alpha_i, \alpha_j) = K(\alpha_1, \alpha_2, \alpha_3) = N.$$

Thus the residual factor, which marks two distinct roots, is already the ordered-root normal closure.

Lean correspondence. The cubic formalization is collected in `CollisionIdeals/ComplexThree/Cubic.lean`, using the generic collision descent in `CollisionIdeals/General/GenericFiber/DiagonalObstruction.lean` and the normal-closure API in `CollisionIdeals/General/Galois/NormalClosure.lean`. From supplied separability, generic-source surjectivity, a cubic power basis, marked normal-closure data, and nonnormality, Lean constructs $D_\alpha \simeq_L N$, proves $H \neq 1$ and $\text{Gal}(N/K) \cong S_3$, and descends the residual factor to $\text{Obs}(F) \neq 0$. The Keller hypothesis is then used only to package the counterexample statement. General inputs from the literature are isolated as theorem interfaces rather than reproved in Lean; in particular, classical Keller–Galois rigidity is recorded as `ComplexKellerGaloisRigidity`. Detailed object-level parity is maintained in `SEMANTIC-PARITY.md`.

4 Further directions

The normal closure also carries a dimension-independent boundary picture, although none of it is needed for the cubic collision theorem. Let L/K be finite separable, let N/K be its normal closure, and let Y be a normal integral model with function field K . Write

$$G = \text{Gal}(N/K), \quad H = \text{Gal}(N/L), \quad Z = \text{Norm}_N(Y) \longrightarrow \overline{X} = \text{Norm}_L(Y) \longrightarrow Y.$$

Suppose that a normal model X of L is an open subscheme of $\overline{X}$ over Y , and that $X \rightarrow Y$ is étale.

Proposition 4.1 (Hidden inertia on a conjugate sheet). *Let $E \in Z^{(1)}$ be ramified over Y , with divisorial valuation w_E , restriction $v = w_E|_K$, and inertia group I_E . Then some $g \in G$ satisfies*

$$e((g^{-1}w_E)|_{L/v}) = [I_E : I_E \cap gHg^{-1}] > 1.$$

For every such g , the center of $(g^{-1}w_E)|_L$ on $\overline{X}$ lies in $\overline{X} \setminus X$.

Proof. Because N is the normal closure of L/K , the action of G on G/H is faithful, so $\bigcap_{g \in G} gHg^{-1} = 1$. Since $I_E \neq 1$, it is not contained in every conjugate of H ; the usual inertia formula gives the displayed index for a suitable g . If the corresponding center belonged to X , the étale morphism $X \rightarrow Y$ would force its ramification index to be one, a contradiction. $\square$

On the generic-degree-three counterexample locus, $G \cong S_3$; hence every ramified divisor of the normal closure is visible on some conjugate sheet, necessarily at its boundary. Paper II develops this boundary geometry for planar Keller maps toward the complementary goal $\text{Obs}(F) = 0$, equivalently $R_F = \Delta$.

5 Conclusion

The collision construction encodes injectivity by the ideal $\text{Obs}(F) \subseteq C_F$. This ideal is the kernel of diagonal evaluation, and its vanishing is equivalent to the scheme-theoretic equality $R_F = \Delta$. After passage to the generic fiber, that diagonal becomes the multiplication factor of $L \otimes_K L$, while the remaining factors record the other embeddings of L .

In generic degree three, the diagonal persists as the first factor beside a quadratic residual algebra, and nonautomorphy is automatic. Thus the counterexample condition is equivalent to the Keller condition. On this locus the classical Galois case rules out the split, normal branch; hence the residual algebra is connected, it is the normal closure N , its Galois group is S_3 , and

$$K \otimes_B C_F \simeq_K L \times N.$$

Its nonzero residual factor gives $\text{Obs}(F) \neq 0$. Hence every map in the generic-degree-three counterexample locus has the normal-closure factor N , with $\text{Gal}(N/K) \cong S_3$. Counterexamples of other generic degrees lie outside the present analysis. The normalization principle of Section 4 explains how the same conjugate sheet geometry leads to boundary questions, which are reserved for the planar sequel.

Motivating example. The explicit three-dimensional polynomial counterexample announced by Levent Alpöge [Alp26] and independently formalized in Isabelle/HOL by Ramos, Hulak, and de Queiroz [RHdQ26] motivated the cubic S_3 comparison. The formal verification establishes a constant nonzero Jacobian determinant and distinct rational points with the same image. That particular map is not used in the arguments above: the dimension-three result describes the structure of the generic-degree-three counterexample locus.

The formalization uses Lean 4 and Mathlib [dMU21, The20].

Corollary 5.1 (First possible generic degree). *If F is Keller and $[L : K] \leq 2$, then F is a polynomial automorphism. Consequently, every complex Jacobian-conjecture counterexample has generic degree at least three.*

Proof. Proposition 2.4 makes L/K finite and separable. In degree one it is trivial, and in degree two it is Galois. Theorem 2.8 applies in either case. $\square$

This low-degree boundary is the handoff to the planar sequel. Paper II asks for the stronger conclusion that every planar Keller map has generic degree one—equivalently, by Galois rigidity and the automorphism criterion, that its collision scheme collapses to the diagonal.

Acknowledgment. OpenAI Codex (GPT-5.6 Sol) assisted with the Lean formalization and manuscript. The author reviewed and takes responsibility for all statements, proofs, citations, and formalization code.

References

- [Alp26] Levent Alpöge. Post announcing an explicit counterexample to the Jacobian conjecture. https://x.com/_alpoge_/status/2079028340955197566, 2026. X post, accessed July 27, 2026.
- [Ax69] James Ax. Injective endomorphisms of varieties and schemes. *Pacific Journal of Mathematics*, 31(1):1–7, 1969.
- [Cam73] L. Andrew Campbell. A condition for a polynomial map to be invertible. *Mathematische Annalen*, 205:243–248, 1973.
- [dMU21] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction—CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635. Springer, 2021.
- [Raz79] Michael J. Razar. Polynomial maps with constant Jacobian. *Israel Journal of Mathematics*, 32(2–3):97–106, 1979.
- [RHdQ26] Arthur Freitas Ramos, David Barros Hulak, and Ruy Jose Guerra Barretto de Queiroz. Formal verification of an explicit counterexample to the jacobian conjecture. https://isa-afp.org/entries/Jacobian_Counterexample.html, 2026. Archive of Formal Proofs, Isabelle/HOL formalization.
- [The20] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs*, pages 367–381. ACM, 2020.
- [The26a] The Stacks Project Authors. Étale algebras over a field. <https://stacks.math.columbia.edu/tag/00U3>, 2026. Lemma 10.143.4, Tag 00U3.
- [The26b] The Stacks Project Authors. Fundamental theorem of Galois theory. <https://stacks.math.columbia.edu/tag/09DW>, 2026. Theorem 9.21.7, Tag 09DW.
- [The26c] The Stacks Project Authors. Normal closures of finite separable extensions. <https://stacks.math.columbia.edu/tag/0EXM>, 2026. Lemma 9.21.5, Tag 0EXM.
- [The26d] The Stacks Project Authors. Normal extensions: a generator criterion. <https://stacks.math.columbia.edu/tag/0BR3>, 2026. Lemma 9.15.6, Tag 0BR3.
- [The26e] The Stacks Project Authors. Primitive elements. <https://stacks.math.columbia.edu/tag/030N>, 2026. Lemma 9.19.1, Tag 030N.
- [The26f] The Stacks Project Authors. Splitting fields and the normal closure. <https://stacks.math.columbia.edu/tag/09DT>, 2026. Lemma 9.16.3, Tag 09DT.
- [Wri81] David Wright. On the Jacobian conjecture. *Illinois Journal of Mathematics*, 25(3):423–440, 1981.