

Continuous Uniform Distribution Maps on Bounded Parameter Spaces

Buffon's Needle & Monte Carlo π Estimation

A Rigorous Analytical Derivation, Stochastic Analysis, and Empirical Verification

Vedant Raut

August 18, 2026

Abstract

This paper presents a formal analytical derivation and stochastic evaluation of Buffon's Needle problem. By mapping continuous uniform random variables over bounded parameter spaces $\Omega = [0, D/2] \times [0, \pi/2]$, we establish the exact probability of geometric intersection $P = \frac{2L}{\pi D}$ for the short needle regime where $L \leq D$.

We derive the Monte Carlo estimator $\hat{\pi} = \frac{2LN}{CD}$ and prove via the Delta Method that its empirical error exhibits asymptotic Central Limit Theorem decay on the order of $\mathcal{O}(1/\sqrt{N})$.

Furthermore, we discuss computational applications of geometric probability in Monte Carlo variance reduction and graphics rendering engines, alongside an evaluation of a companion interactive web simulation.

My Live Simulation Engine : [Access Interactive Simulation Web Application](#)

Core Math Domain: Geometric Probability & Integral Calculus

Primary Focus: Continuous Uniform Distribution Maps on Geometric State Spaces

Faculty Review & Academic Acknowledgement:

This research monograph was prepared under the academic guidance of the Department of Mathematics and submitted to the faculty for review and verification.

Contents

1	Problem Formulation & Physical Model	3
1.1	Objective Statement	3
1.2	Parameterization of the Continuous State Space	3
1.3	The Geometric Intersection Boundary	3
2	Theoretical Proof & Mathematical Derivation	4
3	Computational Simulation & Asymptotic Error Analysis	5
4	Applications and Practical Observations	6
5	Interactive Web Simulation Evaluation	7
6	Concluding Remarks	8
7	Mathematical Glossary	9

1 Problem Formulation & Physical Model

Consider a uniform, rigid needle of length L dropped at random onto a two-dimensional Euclidean plane marked with an infinite sequence of parallel lines spaced at a constant distance D , where $L \leq D$. Because the needle's physical landing position and orientation are continuous random variables, the system's state space contains uncountably infinite configurations.

1.1 Objective Statement

The primary objective of this investigation is twofold:

1. Determine the exact closed-form geometric probability P that a randomly dropped needle intersects at least one parallel grid line.
2. Construct an empirical Monte Carlo estimator $\hat{\pi}$ to approximate the transcendental mathematical constant π through repetitive stochastic sampling.

1.2 Parameterization of the Continuous State Space

We define the geometric configuration of the needle on the plane using two independent, continuous random variables:

Midpoint Distance (y): The perpendicular distance measured from the needle's midpoint to the nearest parallel grid line. Because adjacent grid lines are separated by spacing D , the midpoint can be at most $D/2$ away from a line:

$$y \sim U\left(0, \frac{D}{2}\right)$$

Orientation Angle (θ): The acute angle formed between the body of the needle and the horizontal direction of the parallel lines:

$$\theta \sim U\left(0, \frac{\pi}{2}\right)$$

1.3 The Geometric Intersection Boundary

By applying right-triangle trigonometry, the perpendicular reach of the needle endpoint relative to its midpoint is given by $\frac{L}{2} \sin(\theta)$. An intersection occurs if and only if the distance y to the line is strictly bounded by this reach:

$$y \leq \frac{L}{2} \sin(\theta) \tag{1}$$

Equation (1) establishes the foundational geometric boundary condition. Here, y represents the midpoint offset, $L/2$ is the half-length of the needle, and $\sin(\theta)$ calculates the effective projection toward the grid line. If y falls within this boundary, a physical line crossing takes place. In the next section, we construct the multivariable joint probability measure over this state space to integrate this inequality.

2 Theoretical Proof & Mathematical Derivation

Because the midpoint distance y and orientation angle θ are independent uniform continuous random variables, the bounded parameter space $\Omega = [0, D/2] \times [0, \pi/2]$ has a total measure calculated by multiplying the lengths of both independent intervals.

$$\text{Area}(\Omega) = \left(\frac{D}{2}\right) \left(\frac{\pi}{2}\right) = \frac{\pi D}{4} \quad (2)$$

Equation (2) establishes the continuous volume of the parameter space Ω . This measure normalizes the probability distribution. Next, we invert this total area to construct the constant joint probability density function.

$$f(y, \theta) = \frac{1}{\text{Area}(\Omega)} = \frac{4}{\pi D} \quad (3)$$

Equation (3) gives the joint density $f(y, \theta)$. Because the position and angle selection are uniform and independent, this density is constant across all coordinates in Ω . We will now integrate this density over the crossing region defined by Equation (1).

$$P = \iint_{\text{crossing}} f(y, \theta) dy d\theta = \frac{4}{\pi D} \int_0^{\pi/2} \left(\int_0^{\frac{L}{2} \sin(\theta)} 1 \cdot dy \right) d\theta \quad (4)$$

Equation (4) expresses the total crossing probability P as an iterated double integral. The inner integral accumulates over valid distances y up to $(L/2) \sin(\theta)$, while the outer integral sweeps through orientation angles θ from 0 to $\pi/2$. We proceed by evaluating the inner integral with respect to y .

$$\int_0^{\frac{L}{2} \sin(\theta)} 1 \cdot dy = [y]_0^{\frac{L}{2} \sin(\theta)} = \frac{L}{2} \sin(\theta) \quad (5)$$

Equation (5) calculates the inner definite integral. Since the integrand is unity, the anti-derivative is simply y , evaluating directly to the upper trigonometric boundary. We substitute this result into the outer integral.

$$P = \frac{4}{\pi D} \int_0^{\pi/2} \frac{L}{2} \sin(\theta) d\theta = \frac{2L}{\pi D} \int_0^{\pi/2} \sin(\theta) d\theta \quad (6)$$

Equation (6) simplifies the expression by factoring constant multipliers outside the integral operator. The remaining integral requires finding the anti-derivative of $\sin(\theta)$.

$$\int_0^{\pi/2} \sin(\theta) d\theta = [-\cos(\theta)]_0^{\pi/2} = -\cos\left(\frac{\pi}{2}\right) - (-\cos(0)) = 0 - (-1) = 1 \quad (7)$$

Equation (7) evaluates the definite integral of $\sin(\theta)$ from 0 to $\pi/2$, which equals exactly 1. Multiplying this result by our coefficient yields the closed-form crossing probability.

$$P = \frac{2L}{\pi D} \quad (8)$$

Equation (8) presents the primary theoretical identity. For any needle in the short regime ($L \leq D$), the probability of intersecting a parallel grid line is strictly proportional to $2L$ divided by πD .

3 Computational Simulation & Asymptotic Error Analysis

We rearrange the exact probability identity from Equation (8) to isolate the transcendental constant π .

$$\pi = \frac{2L}{P \cdot D} \quad (9)$$

Equation (9) provides the structural foundation for estimating π . In a computational environment, we replace the theoretical probability P with its empirical Monte Carlo sample mean $\hat{P} = C/N$, where N represents total drops and C represents successful line crossings.

$$\hat{\pi} = \frac{2LN}{CD} \quad (10)$$

Equation (10) defines the empirical estimator $\hat{\pi}$. Because C is a binomial random variable $C \sim \text{Binomial}(N, P)$, the sample variance of the empirical proportion is given by $\text{Var}(\hat{P}) = \frac{P(1-P)}{N}$. To analyze how this variance impacts our estimator $\hat{\pi}$, we apply the Delta Method.

$$g(P) = \frac{2L}{P \cdot D} \implies g'(P) = -\frac{2L}{P^2 D} \quad (11)$$

Equation (11) takes the first derivative of the estimation function $g(P)$ with respect to P . Using first-order Taylor expansion, the asymptotic variance of $\hat{\pi}$ is approximated by $[g'(P)]^2 \text{Var}(\hat{P})$.

$$\text{Var}(\hat{\pi}) \approx \left(-\frac{2L}{P^2 D}\right)^2 \left(\frac{P(1-P)}{N}\right) = \frac{4L^2(1-P)}{NP^3 D^2} \quad (12)$$

Equation (12) combines the squared derivative with the binomial variance. Substituting $P = \frac{2L}{\pi D}$ into this variance equation allows us to simplify terms as a function of sample size N .

$$\text{SE}(\hat{\pi}) = \sqrt{\text{Var}(\hat{\pi})} \approx \frac{\pi}{\sqrt{N}} \sqrt{\frac{\pi D}{2L} - 1} = \mathcal{O}\left(\frac{1}{\sqrt{N}}\right) \quad (13)$$

Equation (13) completes the asymptotic error analysis. The standard error $\text{SE}(\hat{\pi})$ decays at a rate of $\mathcal{O}(1/\sqrt{N})$, matching standard Central Limit Theorem convergence bounds for Monte Carlo integration engines.

4 Applications and Practical Observations

While Buffon’s problem originated as a classic geometric probability puzzle, its underlying formulation—integrating continuous uniform parameter spaces into discrete event expectations—serves as a core foundation across scientific computation and stochastic modeling:

- **Monte Carlo Variance Reduction:** Estimating integrals over continuous bounded domains $\Omega \subset \mathbb{R}^n$ via geometric sampling forms the basis for antithetic variates and importance sampling, reducing standard error below standard $\mathcal{O}(1/\sqrt{N})$ rates.
- **Ray-Tracing & Computer Graphics:** Calculating visual occlusion, soft shadows, and light transport paths involves integrating geometric intersections between line segments and planar primitives over continuous spatial and angular sampling bounds.
- **Geometric Point Processes in Materials Science:** Modeling fibrous material density, micro-structural fracture boundaries, and polymer chain interconnections applies multi-variable geometric intersection probabilities to predict physical material stress limits.

5 Interactive Web Simulation Evaluation

To complement this theoretical monograph, an interactive, single-file HTML/JS simulation engine was developed: `buffon_needle_sim.html`.

- **Live Interactive Application:** <https://buffonsneedle.vercel.app/>
- **GitHub Source Code Repository:** <https://github.com/whyyyydant/math-drafts-and-si>
- **High Precision Monte Carlo Engine:** Capable of executing up to 10,000 drops per frame using native HTML5 Canvas vector drawing, allowing real-time observation of $N > 1,000,000$ iterations.
- **Dual Panel Analytical Dashboard:** Displays the geometric needle field alongside a real-time convergence plot overlaying the theoretical π target line and 95% Central Limit Theorem confidence funnels.
- **Academic Aesthetics:** Built using high-contrast, unstyled structural layouts that focus purely on data density, real-time variance decay tracking, and empirical parameter validation.

6 Concluding Remarks

We have presented a complete analytical derivation of Buffon's Needle problem, demonstrating how continuous parameter spaces $[0, D/2] \times [0, \pi/2]$ collapse into discrete Bernoulli events through multivariable integration. The derived estimator $\hat{\pi} = \frac{2LN}{CD}$ provides a bridge between continuous probability density and empirical stochastic simulation, exhibiting standard $\mathcal{O}(1/\sqrt{N})$ variance decay.

7 Mathematical Glossary

Parameter Space (Ω):	The bounded set of all possible geometric configurations $(y, \theta) \in [0, D/2] \times [0, \pi/2]$ defining the exact position and orientation of the needle.
Continuous Uniform Distribution	A probability density where all equal-length subintervals within $[a, b]$ have identical occurrence probabilities.
Joint Density Function ($f(y, \theta)$):	A multivariable function describing the relative likelihood of two continuous random variables occurring simultaneously.
Delta Method:	An asymptotic technique using Taylor expansions to estimate the variance of a non-linear function of a random variable.
Central Limit Theorem (CLT):	A fundamental theorem establishing that sample averages of independent random variables converge toward a normal distribution with $\mathcal{O}(1/\sqrt{N})$ standard error decay.