

Chapter 23 Convolutions and the Unit Step Function.

23.1 Find  $f(x) * g(x)$  when  $f(x) = e^{3x}$  and  $g(x) = e^{2x}$ .

The title of the chapter is the clue that convolution and unit step functions are related. In Wolfram Alpha, the tasked convolution can be accomplished with the entry:

`!! Convolve[e^(3 t) UnitStep[t], e^(2t) UnitStep[t], t, x] !!`

Exact result

$$e^{2x} (e^x - 1) \theta(x)$$

This leaves the step function in the answer. It is not in the answer in the text, therefore it is necessary to employ the rationalization that inserting the step function for proper functioning of W|A and retracting it when desired are equivalent.

text answer =  $e^{3x} - e^{2x}$

23.3 Find  $f(x) * g(x)$  when  $f(x) = x$  and  $g(x) = x^2$ .

Continuing the off-label use of the function Convolve. The desired output can be obtained by using:

`!! Convolve[t UnitStep[t], t^2 UnitStep[t], t, x] !!`

Exact result

$$\frac{x^4}{12} \theta(x)$$

The unit step function appears as  $\theta$  again.

text answer =  $\frac{1}{12} x^4$

23.4 Find  $\mathcal{L}^{-1} \left\{ \frac{1}{s^2 - 5s + 6} \right\}$  by convolutions.

Once into the landscape of Laplace, the treatment of convolutions shifts. (Is that a pun?) Using Wolfram Alpha it is very straightforward.

`!! simplify[inverselaplaceform[1/(s^2 - 5 s + 6)]] !!`

Result

$$e^{2t} (e^t - 1)$$

text answer =  $e^{3x} - e^{2x}$

23.5 Find  $\mathcal{L}^{-1} \left\{ \frac{6}{s^2 - 1} \right\}$  by convolutions.

And similarly,

`!! simplify[inverselaplace transform[6/(s^2 - 1)]] !!`

Result

$6 \sinh(t)$

For some reason the text did not simplify the solution as shown above. W|A can verify,

`!! (exp(x)-exp(-x))/2 == sinh(x) !!`

with the answer of 'True'.

23.6 Find  $\mathcal{L}^{-1} \left\{ \frac{1}{s(s^2 + 4)} \right\}$  by convolutions.

And another


`!! simplify[inverselaplace transform[1/(s (s^2 + 4))]] !!`

Result

$\frac{\sin^2(t)}{2}$

Not in the same form as text answer, so a plot will be made to consider the two variations.

```
In [41]: 1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 %config InlineBackend.figure_formats = ['svg']
5
6 #x = np.arange(0, 3., 0.005)
7 x = np.linspace(-2,3,300)
8 y3 = (1/2)*(np.sin(x))*(np.sin(x))
9 y4 = (1/4)*(1 - np.cos(2*x))
10
11
12 plt.grid(True, linestyle='dotted', color='gray', linewidth=0.9)
13 plt.xlabel("t")
14 plt.ylabel("f(t)")
15 plt.title("Outline of Differential Equations Prob 23.6")
16 plt.rcParams['figure.figsize'] = [9, 7.5]
17
18
19 ax = plt.gca()
20 ax.axhline(y=0, color='#993399', linewidth=0.8)
21 ax.axvline(x=0, color='#993399', linewidth=0.8)
22 ratio = 0.95
23 xleft, xright = ax.get_xlim()
24 ybottom, ytop = ax.get_ylim()
25 ax.set_aspect(abs((xright-xleft)/(ybottom-ytop))*ratio)
26
27
28 #plt.text(965, -2.5e43, "-np.log(abs((np.cos(x/2))**2 - (np.sin(x/2))**2)))
29 #plt.text(965, 2.25e43, "SOLN: y = 3*np.exp(x**2) + 1/2", size=10,\
30 #       # bbox=dict(boxstyle="square", ec=('8C564B'),fc=(1., 1., 1),))
31 plt.ylim(-1,1.5)
32 plt.plot(x, y3, linewidth = 3)
33 plt.plot(x, y4, linewidth = 0.9, color = 'w')
34 plt.show()
35
36
```



23.7 Find  $\mathcal{L}^{-1} \left\{ \frac{1}{(s-1)^2} \right\}$  by convolutions.

And yet another

```
!! simplify[inverselaplace transform[1/((s - 1)^2)]] !!
```

Result

$e^t t$

text answer =  $x e^x$

23.9 Graph the function  $f(x) = u(x - 2) - u(x - 3)$ .

This calls for a plot of unit step functions. Seeing that


$$u(x - 2) = \begin{cases} 0 & x < 2 \\ 1 & x \geq 2 \end{cases} \quad \text{and} \quad u(x - 3) = \begin{cases} 0 & x < 3 \\ 1 & x \geq 3 \end{cases}$$

it follows that

$$f(x) = u(x - 2) - u(x - 3) = \begin{cases} 0 - 0 = 0 & x < 2 \\ 1 - 0 = 1 & 2 \leq x < 3 \\ 1 - 1 = 0 & x \geq 3 \end{cases}$$


The following plot is a primitive attempt to express the above function graphically. But it works. Following the first plot is a second, which does not apply here but is merely a pointless demo.

```
In [13]: 1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 %config InlineBackend.figure_formats = ['svg']
5
6 x = [2, 3]
7 y = [1, 1]
8 a = [-10, 2]
9 b = [0, 0]
10 z = [3, 5]
11
12 ax = plt.gca()
13 plt.grid(True, linestyle='dotted', color='gray', linewidth=0.9)
14 plt.rcParams['figure.figsize'] = [3, 3]
15 ax.set_title('Outline of Differential Equations Prob 23.9')
16
17 plt.step(x, y)
18 plt.step(a, b, color='#1F77B4')
19 plt.step(z, b, color='#1F77B4')
20
21 ratio = 0.1
22 xleft, xright = ax.get_xlim()
23 ybottom, ytop = ax.get_ylim()
24 ax.set_aspect(abs((xright-xleft)/(ybottom-ytop))*ratio)
25
26 xpts = np.array([2, 3])
27 ypts = np.array([0, 1])
28 wpts = np.array([2, 3])
29 zpts = np.array([1, 0])
30 plt.plot(xpts, ypts, markersize=7, color='#1F77B4', marker='o', \
31          linestyle='none', mfc='none', markeredgewidth=0.5)
32 plt.plot(wpts, zpts, markersize=7, color='#1F77B4', linestyle='none', \
33          marker='o', mfc='#1F77B4', markeredgewidth=0.5)
34
35 plt.ylim(-1,2)
36 plt.xlim(-1,4)
37 plt.show()
38
39
```



```
In [63]: 1 import matplotlib.pyplot as plt
2
3 y = range(1, 10)
4 xmin = range(9)
5 xmax = range(1, 10)
6
7 ax = plt.gca()
8 plt.grid(True, linestyle='dotted', color='gray', linewidth=0.9)
9 plt.rcParams['figure.figsize'] = [5, 3.5]
10
11 ratio = 1.0
12 xleft, xright = ax.get_xlim()
13 ybottom, ytop = ax.get_ylim()
14 ax.set_aspect(abs((xright-xleft)/(ybottom-ytop))*ratio)
15
```

```
16 plt.ylim(-1,10)
17 ax.hlines(y, xmin, xmax)
18 plt.show()
19
```




23.10 Graph the function  $f(x) = 5 - 5u(x - 8)$  for  $x \geq 0$ .

This calls for a plot of unit step functions. The function  $f(x)$  is

$$f(x) = 5 - 5u(x - 8) = \begin{cases} 5 & x < 8 \\ 0 & x \geq 8 \end{cases}$$

A rough imitation of a stepwise plot is shown.

```
In [16]: 1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 %config InlineBackend.figure_formats = ['svg']
5
6 x = [-10, 8]
7 y = [5, 5]
8 a = [8, 12]
9 b = [0, 0]
10 #z = [3, 5]
11
12 ax = plt.gca()
13 plt.grid(True, linestyle='dotted', color='gray', linewidth=0.9)
14 plt.rcParams['figure.figsize'] = [5, 5]
15 ax.set_title('Outline of Differential Equations Prob 23.10')
16
17 plt.step(x, y)
18 plt.step(a, b, color='#1F77B4')
19 #plt.step(z, b, color='#1F77B4')
20
21 ratio = 0.2
22 xleft, xright = ax.get_xlim()
23 ybottom, ytop = ax.get_ylim()
24 ax.set_aspect(abs((xright-xleft)/(ybottom-ytop))*ratio)
25
26 xpts = np.array([-5, 8])
27 ypts = np.array([5, 5])
28 wpts = np.array([8, 13])
29 zpts = np.array([0, 0])
30 plt.plot(xpts, ypts, markersize=7, color='#1F77B4', marker='o', \
31         linestyle='none', mfc='none', markeredgewidth=0.5)
32 plt.plot(wpts, zpts, markersize=7, color='#1F77B4', linestyle='none', \
33         marker='o', mfc='#1F77B4', markeredgewidth=0.5)
34
35 plt.ylim(-1,12)
36 plt.xlim(-1,12)
37 plt.show()
38
39
```



23.11 Find the Laplace transform of  $f(x) = \begin{cases} 0 & x < 4 \\ (x - 4)^2 & x \geq 4 \end{cases}$ .

This is the first problem in this set going from Regularland to Laplaceland. The problem can be entered into Wolfram Alpha:

|| laplace transform of Piecewise[{{0, x < 4}, {(x - 4)^2, x >= 4}}] ||

In Wolfram Alpha the result is expressed as:

Result

$\frac{2 e^{-4 p}}{p^3}$

which is equivalent to the text answer.

23.14 Find the Laplace transform of  $f(x) = \begin{cases} 0 & x < 4 \\ x^2 & x \geq 4 \end{cases}$

This problem can be entered into Wolfram Alpha:

!! laplace transform of Piecewise[{{0, x < 4}, {x^2, x >= 4}}] !!

In Wolfram Alpha the result is expressed as:

Result

$$\frac{2e^{-4p}(8p^2 + 4p + 1)}{p^3}$$

which is equivalent to the text answer.

In [ ]:

In [ ]: