

Note: In this problem set, expressions in green cells match corresponding expressions in the text answers.


4 - 18 Temperature $T(x,y)$ in plates

Find the temperature distribution $T(x,y)$ and the complex potential in the given thin metal plate whose faces are insulated and whose edges are kept at the indicated temperatures or are insulated as shown.

5. Sector

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```



According to example 3 on p. 768 the answer will be in the form

```
cnr[x_, y_] = a  $\theta$  + b
```

```
b + a  $\theta$ 
```

Looking at the geometry of the figure, the T_1 leg has angle $-\frac{\pi}{4}$, and the T_2 leg has angle $\frac{\pi}{4}$, implying that the boundary conditions are seen in

$$a \left(-\frac{\pi}{4} \right) + b = T_2, \text{ and } a \left(\frac{\pi}{4} \right) + b = T_1$$

because of the Arg values of the two T lines, $-\frac{\pi}{4}$ and $\frac{\pi}{4}$, which suggests

```
Solve[-a  $\frac{\pi}{4}$  + b == -20 && a  $\frac{\pi}{4}$  + b == 20, {a, b}]
```

```
{ {a  $\rightarrow$   $\frac{80}{\pi}$ , b  $\rightarrow$  0} }
```

putting this back into the starting equation

```
Simplify[cnr[x, y] /. {a  $\rightarrow$   $\frac{80}{\pi}$ , b  $\rightarrow$  0}]
```

$$\frac{80 \theta}{\pi}$$

According to example 3 on p. 760, $\theta = \text{ArcTan}\left[\frac{y}{x}\right]$

This is not the complete answer. I need the harmonic conjugate of this expression in order to get the complex potential, which is equal to $\Phi[x,y] + i \Psi[x,y]$. This is simple as pie using the identity discussed in problem 15 below and consists of

$$\frac{80 \theta}{\pi} = 80 \text{Arg}[z]$$

and the complex potential is

$$\Phi + i \Psi = -\frac{i 80}{\pi} \text{Log}[z]$$

But maybe I would rather find a different harmonic conjugate to go with my Φ function.

Then with help from utube's MathSorcerer in https://www.youtube.com/watch?v=tWX8YwKfd_k I look for v such that $f = u + i v$ is analytic. First I need the partials of u :

$$u[x, y] = \frac{80}{\pi} \text{ArcTan}\left[\frac{y}{x}\right]$$

$$\frac{80 \text{ArcTan}\left[\frac{y}{x}\right]}{\pi}$$

$$D[u[x, y], x]$$

$$= -\frac{80 y}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)}$$

$$D[u[x, y], y]$$

$$= \frac{80}{\pi x \left(1 + \frac{y^2}{x^2}\right)}$$

Since I'm trying to build this to be analytic, I use Cauchy-Riemann, $D[v[x,y],y] = D[u[x,y],x]$ and $-D[v[x,y],x] = D[u[x,y],y]$. Using the first of the pair of C-R,

$$D[v[x, y], y] = D[u[x, y], x] = -\frac{80 y}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)}$$

So to find the aspect of v which satisfies the y branch I can integrate this partial derivative with respect to y

$$\int -\frac{80 y}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)} dy$$

$$= -\frac{40 \text{Log}[x^2 + y^2]}{\pi}$$

And because I integrated with respect to dy , I need to add an unknown function of x ,

getting

$$-\frac{40 \operatorname{Log}[x^2 + y^2]}{\pi} + g[x]$$

$$g[x] - \frac{40 \operatorname{Log}[x^2 + y^2]}{\pi}$$

as a candidate v function with symbolic x hang-on function. At this point I can differentiate the last expression with respect to x to look for the x aspect of v

$$D\left[g[x] - \frac{40 \operatorname{Log}[x^2 + y^2]}{\pi}, x\right]$$

$$-\frac{80 x}{\pi (x^2 + y^2)} + g'[x]$$

and the above quantity can be set equal to the partial of v with respect to x which I already have (which is equal to the negative of the partial derivative of u with respect to y), thus

$$-D[v[x, y], x] = \frac{-80}{\pi x \left(1 + \frac{y^2}{x^2}\right)} = -\frac{80 x}{\pi (x^2 + y^2)} + g'[x]$$

Leaving the negative signs in now for consistency

$$\text{Solve}\left[-\frac{80}{\pi x \left(1 + \frac{y^2}{x^2}\right)} == -\frac{80 x}{\pi (x^2 + y^2)} + g'[x], g'[x]\right]$$

$$\{\{g'[x] \rightarrow 0\}\}$$

Here I assert that integrating at this point produces $g[x]$ equal to C , and I decide to set $C=0$.

And from here I should be able to build the v function from

$$v[x_, y_] = \text{Simplify}\left[-\frac{40 \operatorname{Log}[x^2 + y^2]}{\pi}\right]$$

$$-\frac{40 \operatorname{Log}[x^2 + y^2]}{\pi}$$

The above expression being only v , the entire complex potential should be equal to

$$\text{FullSimplify}\left[\frac{80}{\pi} \operatorname{ArcTan}\left[\frac{y}{x}\right] + i \left(\frac{40 \operatorname{Log}[x^2 + y^2]}{\pi}\right)\right]$$

$$\frac{80 \operatorname{ArcTan}\left[\frac{y}{x}\right] + 40 i \operatorname{Log}[x^2 + y^2]}{\pi}$$

$$\text{PossibleZeroQ}\left[\frac{80 \operatorname{ArcTan}\left[\frac{y}{x}\right] + 40 i \operatorname{Log}[x^2 + y^2]}{\pi} - \left(\frac{-80 i}{\pi} \operatorname{Log}[x + i y]\right)\right]$$

False

Oh well, I didn't expect to come up with the same Ψ as the text. To defend my answer (which is less elegant looking than the text's) I need to verify that Φ and Ψ are analytic.

```
PossibleZeroQ[D[u[x, y], x] - D[v[x, y], y]]
```

```
True
```

```
PossibleZeroQ[D[u[x, y], y] + D[v[x, y], x]]
```


```
True
```

So according to numbered line (2) on p. 760, Φ and Ψ together make up a complex potential, the thing I was looking for. The yellow cell above matches (in intent though not in content) the text answer. Using the mechanical process of the math sorcerer, I am likely to come up with rather rough looking though hopefully defensible results.

7. Corner

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```



According to example 3 on p. 768 the answer will be in the form

```
tee[x_, y_] = a  $\theta$  + b
```

```
b + a  $\theta$ 
```

Looking at the geometry of the figure, the T_1 leg has angle 0, and the T_2 leg has angle $\frac{\pi}{2}$, implying that the boundary conditions are seen in

$$a \frac{\pi}{2} + b = T_2, \text{ and } a(0) + b = T_1$$

because of the Arg values of the two T lines, $\frac{\pi}{2}$ and 0, which suggests

```
Solve[a  $\frac{\pi}{2}$  + b == T2 && b == T1, {a, b}]
```

```
{ {a → -  $\frac{2(T_1 - T_2)}{\pi}$ , b → T1 } }
```

putting this back into the starting equation

$$\text{Simplify}[\text{tee}[\mathbf{x}, \mathbf{y}] /. \{a \rightarrow -\frac{2 (T_1 - T_2)}{\pi}, b \rightarrow T_1\}]$$

$$T_1 - \frac{2 \theta (T_1 - T_2)}{\pi}$$

substituting $\text{ArcTan}[\frac{y}{x}]$ for θ and rearranging gives the expression

$$T_1 + \frac{2}{\pi} (T_2 - T_1) \text{ArcTan}\left[\frac{y}{x}\right]$$

Though it matches the text, this is not the complete answer. I need the harmonic conjugate of this expression in order to get the complex potential, which is equal to $\Phi[x,y] + i \Psi[x,y]$. This is just as easy as in problem 5.

$$\Phi + i \Psi = T_1 + \frac{-i 2}{\pi} (T_2 - T_1) \text{Log}[z]$$

There is something to consider about the above expression. In the first green cell it is seen that the ArcTan, or Arg component is separate from the T_1 component. The T_1 component will be the Φ and the component containing the Arg device will be the Ψ . So when the $-i$ is applied, it is only applied to the component with the Arg.

Now I will crank through the process of generating a complex potential mechanically. So with another helping hand from utube's MathSorcerer in https://www.youtube.com/watch?v=tWX8YwKfd_k I look for v such that $f = u + i v$ is analytic. First I need the partials of u :

$$u[\mathbf{x}_-, \mathbf{y}_-] = T_1 + \frac{2}{\pi} (T_2 - T_1) \text{ArcTan}\left[\frac{y}{x}\right]$$

$$T_1 + \frac{2 \text{ArcTan}\left[\frac{y}{x}\right] (-T_1 + T_2)}{\pi}$$

$$D[u[\mathbf{x}, \mathbf{y}], \mathbf{x}]$$

$$- \frac{2 y (-T_1 + T_2)}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)}$$

$$D[u[\mathbf{x}, \mathbf{y}], \mathbf{y}]$$

$$\frac{2 (-T_1 + T_2)}{\pi x \left(1 + \frac{y^2}{x^2}\right)}$$

Since I'm trying to build this to be analytic, I use Cauchy-Riemann, $D[v[x,y],y] = D[u[x,y],x]$ and $-D[v[x,y],x] = D[u[x,y],y]$. Using the first of the pair of C-R,

$$D[v[x, y], y] = -\frac{2 y (-T_1 + T_2)}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)}$$

So to find the aspect of v which satisfies the y branch I can integrate this partial derivative with respect to y

$$\int -\frac{2 y (-T_1 + T_2)}{\pi x^2 \left(1 + \frac{y^2}{x^2}\right)} dy$$

$$\frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi}$$

And because I integrated with respect to dy, I need to add an unknown function of x, getting

$$\frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi} + g[x]$$

$$g[x] + \frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi}$$

as a candidate v function with symbolic x hang-on function. At this point I can differentiate the last expression with respect to x to look for the x aspect of v

$$D\left[g[x] + \frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi}, x\right]$$

$$\frac{2 x (T_1 - T_2)}{\pi (x^2 + y^2)} + g'[x]$$

and the above quantity can be set equal to the partial of v with respect to x which I already have (which is equal to the negative of the partial derivative of u with respect to y), thus

$$-D[v[x, y], x] = -\frac{2 (-T_1 + T_2)}{\pi x \left(1 + \frac{y^2}{x^2}\right)} = \frac{2 x (T_1 - T_2)}{\pi (x^2 + y^2)} + g'[x]$$

Leaving the negative sign in for consistency

$$\text{Solve}\left[-\frac{2 (-T_1 + T_2)}{\pi x \left(1 + \frac{y^2}{x^2}\right)} == \frac{2 x (T_1 - T_2)}{\pi (x^2 + y^2)} + g'[x], g'[x]\right]$$

$$\{\{g'[x] \rightarrow 0\}\}$$

Here I assert that integrating at this point produces g[x] equal to simply C, and I decide to set C=0.

And from here I should be able to build the v function from

$$v[x_, y_] = \text{FullSimplify}\left[\frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi}\right]$$

$$\frac{\text{Log}[x^2 + y^2] (T_1 - T_2)}{\pi}$$

And the entire complex potential should be equal to

$$T_1 + \frac{2}{\pi} (T_2 - T_1) \operatorname{ArcTan}\left[\frac{y}{x}\right] + i \left(\frac{\operatorname{Log}[x^2 + y^2] (T_1 - T_2)}{\pi} \right)$$

$$T_1 + \frac{i \operatorname{Log}[x^2 + y^2] (T_1 - T_2)}{\pi} + \frac{2 \operatorname{ArcTan}\left[\frac{y}{x}\right] (-T_1 + T_2)}{\pi}$$

Again the Ψ is much different from the text. To defend my answer I need to verify that Φ and Ψ are analytic.

```
PossibleZeroQ[D[u[x, y], x] - D[v[x, y], y]]
```

```
True
```

```
PossibleZeroQ[D[u[x, y], y] + D[v[x, y], x]]
```

```
True
```


So according to numbered line (2) on p. 760, Φ and Ψ together make up a complex potential, the thing I was looking for. The yellow cell above matches (in intent though not in content) the text answer.

9. Upper half-plane

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```

```
innerbw = RGBColor[.97, .97, .994];
```



This problem is basically the same as example 3 on p. 760, applying a heat perspective instead of an electrostatic perspective. From there, the solution equation looks like

$$\Phi[x, y] = a + b \operatorname{Arg}[z]$$

Note here that the algebraic a and b in the above expression are unrelated to the points in the sketch.

There are three ϕ functions, according to location, call them Φ_1 , Φ_2 , and Φ_3 , separated by the points a and b . The two quantities Φ_1 and Φ_3 both have the π angle, the same as Φ_2 . However, because they have zero temperature, their contributions disappear, leaving only Φ_2 , which has both magnitude and angle. So the equation for ϕ reduces to simply

$$0 + b \operatorname{Arg}[z] = \Phi_2 = \frac{T_1}{\pi} \theta = \frac{T_1}{\pi} \operatorname{ArcTan}\left[\frac{y}{x}\right]$$

The angle $\frac{y}{x}$ has an interpretation here, because in this problem the x interval is subdivided. It is necessary to get rid of everything that does not describe T_1 , which is all x beyond the T_1 segment. To do this can look like the following:

$$\frac{T_1}{\pi} \left(\text{ArcTan} \left[\frac{y}{x-b} \right] - \text{ArcTan} \left[\frac{y}{x-a} \right] \right)$$

Now heading toward the complex potential form. The part in parentheses above could be treated like in problem 15 below, since it is the Arg,

$$\begin{aligned} \Phi + i\Psi &= \frac{i T_1}{\pi} \left(\text{Log} \left[\frac{y}{x-b} \right] - \text{Log} \left[\frac{y}{x-a} \right] \right) = \\ &= \frac{i T_1}{\pi} \left(\text{Log} \left[\frac{y}{x-b} \right] + \text{Log} \left[\frac{x-a}{y} \right] \right) = \frac{i T_1}{\pi} \left(\text{Log} \left[\frac{x-a}{x-b} \right] \right) \end{aligned}$$


When substituting $i \text{ Log}$ for Arg it is necessary to remember the minus sign.

11. Upper half-plane

```
Clear["Global`*"]

kru = RGBColor[0.392, 0.823, 0.98];
innerbw = RGBColor[.97, .97, .994];

Graphics[{{Line[{{1.15, 0}, {1.15, 1}}]}, {Line[{{0, 0}, {2.3, 0}}]},
  {kru, Opacity[0.6], Polygon[{{0, 0}, {2.3, 0}, {2.2, 0.15}, {2.1, 0.2},
    {2, 0.23}, {1.7, 0.55}, {1, 0.6}, {0.7, 0.66}, {0.5, 0.7}, {0, 0}}]},
  {Text[Style["T=100°C", Medium], {1.15, -0.12}]},
  {Text[Style["T=0", Medium], {0.3, -0.12}]},
  {Text[Style["x", Medium], {2.3, -0.1}]},
  {Text[Style["y", Medium], {1.05, 0.95}]},
  {EdgeForm[Directive[Black]], innerbw, Disk[{0.6, 0}, 0.04]},
  {EdgeForm[Directive[Black]], innerbw, Disk[{1.69, 0}, 0.04]},
  {Text[Style["T=0", Medium], {1.92, -0.12}]},
  {Text[Style["-1", Medium], {0.61, .12}]},
  {Text[Style["1", Medium], {1.7, .12}]}], ImageSize -> 200]
```



This is apparently like problem 9, except that now the points a and b are assigned specific values. Again I look to example 3 on p. 760, applying a heat perspective instead of an electrostatic perspective. From there, the solution equation is

$$\Phi[x, y] = a + b \operatorname{Arg}[z]$$

Here there are no labels on the sketch to confuse with the variables in the equation above.

There are again three phi functions, according to location, call them Φ_1 , Φ_2 , and Φ_3 , separated by the points $\{-1, 0\}$ and $\{1, 0\}$. The two quantities Φ_1 and Φ_3 both have the π angle, the same as Φ_2 . However, because they have zero temperature, their contributions disappear, leaving only Φ_2 , which has both magnitude and angle. So the equation for phi reduces to simply

$$0 + b \operatorname{Arg}[z] = \Phi_2 = \frac{T_1}{\pi} \theta = \frac{T_1}{\pi} \operatorname{Arg}[z] = \frac{100}{\pi} \operatorname{Arg}[z]$$

The axis is clearly labeled x, but since y is equal to zero, the expression above is still true. And, applying the point location elimination,

$$\Phi = \frac{100}{\pi} (\operatorname{Arg}[z - 1] - \operatorname{Arg}[z - (-1)]) = \frac{100}{\pi} (\operatorname{Arg}[z - 1] - \operatorname{Arg}[z + 1])$$


Now it is time to push for the complex potential expression, and as in problem 15, and retaining the z nomenclature to agree with text,

$$\Phi + i\Psi = \frac{i 100}{\pi} (-\operatorname{Log}[z - 1] + \operatorname{Log}[z + 1]) = \frac{i 100}{\pi} \left(\frac{\operatorname{Log}[z + 1]}{\operatorname{Log}[z - 1]} \right)$$

13. Corner

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```



I can tell from the text answer that this one will require a mapping, and the text answer suggests using $w=z^2$.

Setting up a list of test points

```
sx = {{0, 0}, {0, 1}, {1, 0}, {1, 1}}
{{0, 0}, {0, 1}, {1, 0}, {1, 1}}
```

And a point translation function independent of the plot


```
gp[{x_, y_}] = {N[Re[(x + I y)^2]], N[Im[(x + I y)^2]]}
{Re[(x + (0. + 1. I) y)^2], Im[(x + (0. + 1. I) y)^2]}
```

to get sample points for direct plotting

```
Thread[gp[sx]]
{{0., 0.}, {-1., 0.}, {1., 0.}, {0., 2.}}
```

and then showing the plot

```
d2 = DiscretizeRegion@ImplicitRegion[0 ≤ x ≤ 1 ∧ 0 ≤ y ≤ 1, {x, y}];
ParametricPlot[ReIm[(x + I y)^2], {x, y} ∈ d2,
  PlotRange → {{-2, 2}, {-1, 2.5}}, Frame → True,
  ImageSize → 200, AspectRatio → Automatic,
  Epilog → {{Red, PointSize[0.025], Point[{0, 0]}},
    {Text[Style[A, Medium], {0, -0.7}]}, {Green, PointSize[0.025],
    Point[{-1, 0}]}, {Text[Style[B, Medium], {-1, -0.7}]},
    {Blue, PointSize[0.025], Point[{1, 0}]},
    {Text[Style[C, Medium], {1, -0.7}]},
    {Text[Style[D, Medium], {0, 2.2}]}]}
```



From working other problems I know that any intervals associated with zero temp will disappear, so I don't include these in the plot. Φ_1 is from B to A on w-plane, Φ_2 is from A to C. Both functions have an angle component of π , and the same temperature, 100 °C. I can see that the answer will be of the form

$$\Phi_1 + \Phi_2 = \frac{100}{\pi} (\text{Arg}[BA]) + \frac{100}{\pi} (\text{Arg}[AC])$$


The $\frac{100}{\pi}$ part does not need re-translation back to the z-plane. And the z^2 mapping function is simple enough that it can be expressed in the answer. Now looking at the Arg function, I see it follows the boundary of the $w=z^2$ curve, and is offset on each side by the locations of B and C. The Arg expression will be affected by B and C in a similar way to the way an expression for a circle is affected by the coordinate of its center. And the mirror image of the function curves indicates a collision in sign, which will show up as

$$\Phi_1 + \Phi_2 = \frac{100}{\pi} (\text{Arg}[z^2 - 1]) - \frac{100}{\pi} (\text{Arg}[z^2 + 1]) = \frac{100}{\pi} (\text{Arg}[z^2 - 1] - \text{Arg}[z^2 + 1])$$

15. Sector

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```



Starting with the statement that a potential in an angular region with sides at constant temperature has the form

$$T = a \text{Arg}[z] + b$$

$$b + a \text{Arg}[z]$$

As stated in the text, $\text{Arg}[z] = \theta = \text{Im}[\text{Log}[z]]$ is a harmonic function. The coefficients a and b are boundary conditions determined with the initial conditions. On the horizontal axis $\text{Arg}[z] = 0$, which makes it easy to calculate b since $T = 0 + b = -20$. For the other leg, $\text{Arg}[z] = \frac{\pi}{4}$ is straightforward because b has already been calculated

$$\text{Solve}\left[a \frac{\pi}{4} - 20 = 60, a\right]$$

$$\left\{\left\{a \rightarrow \frac{320}{\pi}\right\}\right\}$$

$$\text{Tf} = T /. \left\{a \rightarrow \frac{320}{\pi}, b \rightarrow -20\right\}$$

$$-20 + \frac{320 \text{Arg}[z]}{\pi}$$

The above cell matches the text answer. But it remains to find the complex potential.

Pulling out an oldie but goodie from numbered line (2) on p. 637, $\text{Ln}[z] = \ln[\text{Abs}[z]] + i \text{Arg}[z]$, (with $z \neq 0$).

Since in *Mathematica* each complex z is treated and reported as a principal value, the text's nomenclature is used in highlighted yellow above. Numbered line (3) on p. 637 should be shown as well, $\ln[z] = \text{Ln}[z] \pm 2n\pi i$. In other words, as the text uses the term, $\ln[z]$, (or in this case $\ln[\text{Abs}[z]]$), has an infinite number of values, including when n equals zero, mean-

ing the term $\ln[\text{Abs}[z]]$ is ignorable.

Looking at the identity in numbered line (2), its prominent member is $\text{Arg}[z]$, which is modified by coefficient i . In the $\Phi + i\Psi$ which I am building, the $\text{Arg}[z]$ will reside in the Ψ , so I make use of $-i \text{Log}[z] = \text{Arg}[z]$.

So the complex potential can be assigned to the value

$$-20 - \frac{320i}{\pi} \text{Log}[z]$$


and since z is understood and agreed by Mathematica to be the principal value, the answer is compatible with the text.

17. First quadrant of the z -plane with y -axis kept at 100°C , the segment $0 < x < 1$ of the x -axis insulated and the x -axis for $x > 1$ kept at 200°C . *Hint.* Use example 4.

```
Clear["Global`*"]
```

```
kru = RGBColor[0.392, 0.823, 0.98];
```

```
innerbw = RGBColor[.97, .97, .994];
```



The problem hints that example 4 may be useful. Example 4 uses the ArcSin function to map a heated environment onto the w -plane. First step is to create a list of sample points

```
sx = {{0, 0}, {1, 0}, {1.5, 0}, {0, 1}}
```

```
{{0, 0}, {1, 0}, {1.5, 0}, {0, 1}}
```

and to define an independent function to plot the sample points

```
gp[{x_, y_}] = {N[Re[ArcSin[(x + I y)]]], N[Im[ArcSin[(x + I y)]]]}
```

```
{Re[ArcSin[x + (0. + 1. i) y]], Im[ArcSin[x + (0. + 1. i) y]]}
```

then to display the sample points (uh-oh, look at the third point below)

```
Thread[gp[sx]]
```


```
{{0., 0.}, {1.5708, 0.}, {1.5708, -0.962424}, {0., 0.881374}}
```

then to plot the ArcSin function, which looks pretty ragged with its drooping flagstaff.

```

d2 = DiscretizeRegion@ImplicitRegion[0 < x ≤ 1.5 ∧ 0 < y ≤ 20, {x, y}];
ParametricPlot[ReIm[ArcSin[(x + I y)]], {x, y} ∈ d2,
  PlotRange → {{-1, 3}, {-1.5, 4}}, Frame → False,
  Axes → True, ImageSize → 200, AspectRatio → Automatic,
  Epilog → {{Gray, Rectangle[{0, -0.1}, {1.57, 0}]},
    {Red, PointSize[0.025], Point[{0, 0}]},
    {Text[Style[A, Medium], {0, -0.3}]}, {Green, PointSize[0.025],
    Point[{0, 0.88}]}, {Text[Style[D, Medium], {-0.2, 0.88}]},
    {Blue, PointSize[0.025], Point[{1.57, 0}]},
    {Text[Style[B, Medium], {1.68, -0.2}]}, {Black, PointSize[0.025],
    Point[{1.57, -0.96}]}, {Text[Style[C, Medium], {1.7, -1.06}]}}]

```



This does not look good. The x-axis beyond point B is being mapped negatively down the flagstaff. This is not what example 4 led me to expect. How can this possibly work?

At the Digital Library of Mathematical Functions (<https://dlmf.nist.gov/4.23#E16>) I found this:


$$\arcsin = -i \operatorname{Log} \left[\left(1 - (x + i y)^2 \right)^{0.5} + i (x + i y) \right]$$

which I guess means that I have to take some care if I want to invert the sine function in the complex domain. So I will re-do the plot using this new information.

```

d2 = DiscretizeRegion@ImplicitRegion[0 < x ≤ 80 ∧ 0 < y ≤ 100, {x, y}];
sx = {{0.001, 0.001}, {1, 0}, {1.5, 0.001}, {0.001, 1}}
gq[{x_, y_}] = {N[Re[-i Log[(1 - (x + I y)^2)^0.5 + i (x + I y)]]],
  N[Im[-i Log[(1 - (x + I y)^2)^0.5 + i (x + I y)]]]}
Thread[gq[sx]]
ParametricPlot[ReIm[-i Log[(1 - (x + I y)^2)^0.5 + i (x + I y)]]],
  {x, y} ∈ d2, PlotRange → {{-3, 2}, {-1, 6}}, Frame → False,
  Axes → True, ImageSize → 200, AspectRatio → Automatic,
  Epilog → {{Gray, Rectangle[{0, -0.1}, {1.57, 0}]},
    {Red, PointSize[0.025], Point[{0.001, 0.001}]},
    {Text[Style[A, Medium], {0, -0.3}]}, {Green, PointSize[0.025],
    Point[{0, 0.88}]}, {Text[Style[D, Medium], {-0.2, 0.88}]},
    {Blue, PointSize[0.025], Point[{1.57, 0}]},
    {Text[Style[B, Medium], {1.68, -0.2}]}, {Black, PointSize[0.025],
    Point[{1.57, 0.96}]}, {Text[Style[C, Medium], {1.25, 1.06}]}}]
{{0.001, 0.001}, {1, 0}, {1.5, 0.001}, {0.001, 1}}
{Im[Log[(1. - 1. (x + (0. + 1. i) y)^2)^0.5 + (0. + 1. i) (x + (0. + 1. i) y)]]],
  -1. Re[Log[(1. - 1. (x + (0. + 1. i) y)^2)^0.5 + (0. + 1. i) (x + (0. + 1. i) y)]]}
{{0.001, 0.001}, {1.5708, 0.},
  {1.5699, 0.962424}, {0.000707107, 0.881374}}

```



Okay, this looks better. Since the test points are the same as before except for the sign of the v value of point C, I will use the first instantiation of the ArcSine plot, and consider this one a visual correction. Just for clarity, D-A is at 100°C, A-B is insulated, and B-C is at 200°C. Since they are parallel in the w -plane, it is like a calculation for parallel plates. I don't think there can be two function coefficient terms, and A is located at zero, which I think zaps it, leaving the field open for B. So the calculation would be

$$\text{Solve}\left[\mathbf{a} \cdot \mathbf{0} + \mathbf{b} == 100 \ \&\& \ \mathbf{a} \frac{\pi}{2} + \mathbf{b} == 200, \{\mathbf{a}, \mathbf{b}\}\right]$$

$$\left\{\left\{\mathbf{a} \rightarrow \frac{200}{\pi}, \mathbf{b} \rightarrow 100\right\}\right\}$$

Under the reasoning I just used, the $\frac{\pi}{2}$ in the above set refers to the position of B, not to the angle of B-C with the u-axis (also equal to $\frac{\pi}{2}$). The equation for Φ then would be

$$100 + \frac{200}{\pi} \text{Arg}\left[\frac{\mathbf{y}}{\mathbf{x}}\right]$$

in the w-plane. But the solution needs to be referred back to the z-plane where it started, so the simple Arg has to be embroidered to express the mapping, I think by writing

$$100 + \frac{200}{\pi} \text{ArcSin}[\mathbf{z}]$$

In the text answer this expression is

$$\text{Re}[\mathbf{F}[\mathbf{z}]] = 100 + \frac{200}{\pi} \text{Re}[\text{ArcSin}[\mathbf{z}]]$$

And I assume it is written this way to make clear that although it has two parts, it is not a complex potential.