

Some extracurricular practice junk up at the top of this section. For the first actual problem, see the orange cell below, numbered 1.

```
grating = DSolve[ $x^2 y''[x] + 1.5 x y'[x] - 0.5 y[x] == 0$ ,  $y[x]$ ,  $x$ ]
```

```
{ { $y[x] \rightarrow x^{0.5+0. \pm} C[1] + x^{-1.+0. \pm} C[2]$  } }
```

```
grid = Simplify[grating]
```

```
{ { $y[x] \rightarrow \frac{x^{1.5} C[1] + C[2]}{x^{1.}}$  } }
```

```
Apart[grid]
```

```
{ { $y[x] \rightarrow x^{0.5} C[1] + \frac{C[2]}{x^{1.}}$  } }
```

Cauchy-Euler equation. The auxiliary equation is  $m^2 + 0.5 m - 0.5 == 0$

```
Solve[ $m^2 + 0.5 m - 0.5 == 0$ ,  $m$ ]
```

```
{ { $m \rightarrow -1.$  }, { $m \rightarrow 0.5$  } }
```

Numbered line (1) on p. 71 goes like this:  $x^2 y'' + a x y' + b y = 0$

Numbered line (2) on p. 71 goes like this:  $m^2 + (a - 1) m + b = 0$

1. Double root. Verify directly by substitution that  $x^{(1-1)/2} \text{Log}[x]$  is a solution of (1) if (2) has a double root, but  $x^{m_1} \text{Log}[x]$  and  $x^{m_2} \text{Log}[x]$  are not solutions of (1) if the roots  $m_1$  and  $m_2$  of (2) are different.

```
ClearAll["Global`*"]
```

Solving (2) might be recommended.

```
Solve[ $m^2 + (a - 1) m + b == 0$ ,  $m$ ]
```

```
{ { $m \rightarrow \frac{1}{2} \left( 1 - a - \sqrt{1 - 2 a + a^2 - 4 b} \right)$  }, { $m \rightarrow \frac{1}{2} \left( 1 - a + \sqrt{1 - 2 a + a^2 - 4 b} \right)$  } }
```

The above lists the two possible roots to (2). A double root would mean they're equal.

```
Solve[ $\left( -\sqrt{(1 - a)^2 - 4 b} \right) == \left( \sqrt{(1 - a)^2 - 4 b} \right)$ ,  $a$ ]
```

```
{ { $a \rightarrow 1 - 2 \sqrt{b}$  }, { $a \rightarrow 1 + 2 \sqrt{b}$  } }
```

So it looks like (2) has a double root under two circumstances. And the relationship between a and b has also been established (call it 'Rel'). Now it is time to look at the problem's proposed sol'n.

```
y[x_] :=  $x^{\frac{(1-a)}{2}} \text{Log}[x]$ 
```

```
cran = FullSimplify [ $x^2 y''[x] + a x y'[x] + b y[x]$ ]  
-  $\frac{1}{4} ((-1 + a)^2 - 4 b) x^{\frac{1}{2} - \frac{a}{2}} \text{Log}[x]$ 
```

```
Simplify [cran /.  $a \rightarrow (1 - 2 \sqrt{b})$ ]  
0
```

```
Simplify [cran /.  $a \rightarrow (1 + 2 \sqrt{b})$ ]  
0
```

I think I have done what the first part of the problem asked: if (2) has a double root, a certain relationship ‘Rel’ exists between the coefficients  $a$  and  $b$  in the double root. The proposed sol’n only incorporates one of the coefficients, but the candidate equation does have both, and after substituting the proposed sol’n into the candidate equation and analyzing the resulting wreckage, the relationship ‘Rel’ is found to be present. In other words, invoking ‘Rel’ proves the proposed sol’n to be an actual sol’n.

For the second part of the problem. If the roots of (2) are not equal, it would imply that  $m_1 = \frac{1}{2} \left( 1 - a - \sqrt{(1 - a)^2 - 4b} \right) \neq m_2 = \frac{1}{2} \left( 1 - a + \sqrt{(1 - a)^2 - 4b} \right)$ . It seems obvious that the exclamation point cannot be removed by squaring both sides! However, I don’t see what is wrong with multiplying both sides (of the pertinent sub-expressions) by the same quantity, e.g.  $\sqrt{(1 - a)^2 - 4b}$ .

```
-  $((1 - a)^2 - 4b) \neq ((1 - a)^2 - 4b)$   
-  $(1 - a)^2 + 4b \neq (1 - a)^2 - 4b$ 
```

```
 $8b \neq 2(1 - a)^2$ 
```

```
 $4b \neq (1 - a)^2$ 
```

But in the expansion of **cran** something very much like this came up. What if I look at

```
Simplify [ $(1 - a)^2 == (a - 1)^2$ ]
```

```
True
```

Let me bring **cran** back.

```
cran
```

```
-  $\frac{1}{4} ((-1 + a)^2 - 4 b) x^{\frac{1}{2} - \frac{a}{2}} \text{Log}[x]$ 
```

I contend that having the expression inside parentheses above, i.e.  $((-1 + a)^2 - 4b)$ , equal to zero is necessary for **cran** to go to zero, and, backtracking a few steps, this only happens if the roots are equal. This line of reasoning does not manipulate the specific proposed sol’ns,

however. Incidentally, there is no text answer for this problem.

$$3. \ 5x^2y'' + 23xy' + 16.2y = 0$$

```
ClearAll["Global`*"]
```

```
doxy = {5 x^2 y''[x] + 23 x y'[x] + 16.2 y[x] == 0}
```

```
plat = DSolve[doxy, y, x]
```

```
{16.2 y[x] + 23 x y'[x] + 5 x^2 y''[x] == 0}
```

```
{ {y -> Function[{x},  $\frac{C[1]}{x^{1.8}} + \frac{1.8 C[2] \text{Log}[x]}{x^{1.8}}$  ] } }
```

```
gerz = plat[[1, 1, 2, 2]]
```

```
 $\frac{C[1]}{x^{1.8}} + \frac{1.8 C[2] \text{Log}[x]}{x^{1.8}}$ 
```

```
derz = gerz /. {C[1] -> c1, C[2] -> c2}
```

$$\frac{c1}{x^{1.8}} + \frac{1.8 c2 \text{Log}[x]}{x^{1.8}}$$

The above answer matches the text. Here 1.8\*c2 equals the text answer constant  $c_2$ .

```
lint[x_, c1_, c2_] := derz
```

```
TableForm[Table[{x, c1, c2, derz}, {x, 4}, {c1, -1, 1}, {c2, -3, 3, .25}],  
TableHeadings -> {{}, {"c1=-1", "c1=0", "c1=1"}}]
```


	c1=-1				c1=0				c1=1			
1	-1	-3.	-1.		1	0	-3.	0.	1	1	-3.	1.
1	-1	-2.75	-1.		1	0	-2.75	0.	1	1	-2.75	1.
1	-1	-2.5	-1.		1	0	-2.5	0.	1	1	-2.5	1.
1	-1	-2.25	-1.		1	0	-2.25	0.	1	1	-2.25	1.
1	-1	-2.	-1.		1	0	-2.	0.	1	1	-2.	1.
1	-1	-1.75	-1.		1	0	-1.75	0.	1	1	-1.75	1.
1	-1	-1.5	-1.		1	0	-1.5	0.	1	1	-1.5	1.
1	-1	-1.25	-1.		1	0	-1.25	0.	1	1	-1.25	1.
1	-1	-1.	-1.		1	0	-1.	0.	1	1	-1.	1.
1	-1	-0.75	-1.		1	0	-0.75	0.	1	1	-0.75	1.
1	-1	-0.5	-1.		1	0	-0.5	0.	1	1	-0.5	1.
1	-1	-0.25	-1.		1	0	-0.25	0.	1	1	-0.25	1.
1	-1	0.	-1.		1	0	0.	0.	1	1	0.	1.
1	-1	0.25	-1.		1	0	0.25	0.	1	1	0.25	1.
1	-1	0.5	-1.		1	0	0.5	0.	1	1	0.5	1.
1	-1	0.75	-1.		1	0	0.75	0.	1	1	0.75	1.
1	-1	1.	-1.		1	0	1.	0.	1	1	1.	1.
1	-1	1.25	-1.		1	0	1.25	0.	1	1	1.25	1.
1	-1	1.5	-1.		1	0	1.5	0.	1	1	1.5	1.
1	-1	1.75	-1.		1	0	1.75	0.	1	1	1.75	1.
1	-1	2.	-1.		1	0	2.	0.	1	1	2.	1.
1	-1	2.25	-1.		1	0	2.25	0.	1	1	2.25	1.
1	-1	2.5	-1.		1	0	2.5	0.	1	1	2.5	1.
1	-1	2.75	-1.		1	0	2.75	0.	1	1	2.75	1.
1	-1	3.	-1.		1	0	3.	0.	1	1	3.	1.

2	-1	-3.	-1.36207	2	0	-3.	-1.07489	2	1	-3.	-0.787
2	-1	-2.75	-1.27249	2	0	-2.75	-0.985319	2	1	-2.75	-0.698
2	-1	-2.5	-1.18292	2	0	-2.5	-0.895744	2	1	-2.5	-0.608
2	-1	-2.25	-1.09334	2	0	-2.25	-0.80617	2	1	-2.25	-0.518
2	-1	-2.	-1.00377	2	0	-2.	-0.716595	2	1	-2.	-0.429
2	-1	-1.75	-0.914195	2	0	-1.75	-0.627021	2	1	-1.75	-0.339
2	-1	-1.5	-0.824621	2	0	-1.5	-0.537446	2	1	-1.5	-0.250
2	-1	-1.25	-0.735047	2	0	-1.25	-0.447872	2	1	-1.25	-0.160
2	-1	-1.	-0.645472	2	0	-1.	-0.358298	2	1	-1.	-0.071
2	-1	-0.75	-0.555898	2	0	-0.75	-0.268723	2	1	-0.75	0.0184
2	-1	-0.5	-0.466323	2	0	-0.5	-0.179149	2	1	-0.5	0.1080
2	-1	-0.25	-0.376749	2	0	-0.25	-0.0895744	2	1	-0.25	0.1976
2	-1	0.	-0.287175	2	0	0.	0.	2	1	0.	0.2871
2	-1	0.25	-0.1976	2	0	0.25	0.0895744	2	1	0.25	0.3767
2	-1	0.5	-0.108026	2	0	0.5	0.179149	2	1	0.5	0.4663
2	-1	0.75	-0.0184513	2	0	0.75	0.268723	2	1	0.75	0.5558
2	-1	1.	0.0711231	2	0	1.	0.358298	2	1	1.	0.6454
2	-1	1.25	0.160697	2	0	1.25	0.447872	2	1	1.25	0.7350
2	-1	1.5	0.250272	2	0	1.5	0.537446	2	1	1.5	0.8246
2	-1	1.75	0.339846	2	0	1.75	0.627021	2	1	1.75	0.9141
2	-1	2.	0.429421	2	0	2.	0.716595	2	1	2.	1.0037
2	-1	2.25	0.518995	2	0	2.25	0.80617	2	1	2.25	1.0933
2	-1	2.5	0.60857	2	0	2.5	0.895744	2	1	2.5	1.1829
2	-1	2.75	0.698144	2	0	2.75	0.985319	2	1	2.75	1.2724
2	-1	3.	0.787718	2	0	3.	1.07489	2	1	3.	1.3620
3	-1	-3.	-0.95956	3	0	-3.	-0.821145	3	1	-3.	-0.682
3	-1	-2.75	-0.891131	3	0	-2.75	-0.752716	3	1	-2.75	-0.614
3	-1	-2.5	-0.822702	3	0	-2.5	-0.684288	3	1	-2.5	-0.545
3	-1	-2.25	-0.754273	3	0	-2.25	-0.615859	3	1	-2.25	-0.477
3	-1	-2.	-0.685845	3	0	-2.	-0.54743	3	1	-2.	-0.409
3	-1	-1.75	-0.617416	3	0	-1.75	-0.479001	3	1	-1.75	-0.340
3	-1	-1.5	-0.548987	3	0	-1.5	-0.410573	3	1	-1.5	-0.272
3	-1	-1.25	-0.480558	3	0	-1.25	-0.342144	3	1	-1.25	-0.203
3	-1	-1.	-0.41213	3	0	-1.	-0.273715	3	1	-1.	-0.135
3	-1	-0.75	-0.343701	3	0	-0.75	-0.205286	3	1	-0.75	-0.066
3	-1	-0.5	-0.275272	3	0	-0.5	-0.136858	3	1	-0.5	0.0015
3	-1	-0.25	-0.206843	3	0	-0.25	-0.0684288	3	1	-0.25	0.0699
3	-1	0.	-0.138415	3	0	0.	0.	3	1	0.	0.1384
3	-1	0.25	-0.0699858	3	0	0.25	0.0684288	3	1	0.25	0.2068
3	-1	0.5	-0.00155702	3	0	0.5	0.136858	3	1	0.5	0.2752
3	-1	0.75	0.0668717	3	0	0.75	0.205286	3	1	0.75	0.3437
3	-1	1.	0.135301	3	0	1.	0.273715	3	1	1.	0.4121
3	-1	1.25	0.203729	3	0	1.25	0.342144	3	1	1.25	0.4805
3	-1	1.5	0.272158	3	0	1.5	0.410573	3	1	1.5	0.5489
3	-1	1.75	0.340587	3	0	1.75	0.479001	3	1	1.75	0.6174
3	-1	2.	0.409016	3	0	2.	0.54743	3	1	2.	0.6858
3	-1	2.25	0.477444	3	0	2.25	0.615859	3	1	2.25	0.7542
3	-1	2.5	0.545873	3	0	2.5	0.684288	3	1	2.5	0.8227
3	-1	2.75	0.614302	3	0	2.75	0.752716	3	1	2.75	0.8911
3	-1	3.	0.682731	3	0	3.	0.821145	3	1	3.	0.9595

4	-1	-3.	-0.699833	4	0	-3.	-0.617364	4	1	-3.	-0.534
4	-1	-2.75	-0.648386	4	0	-2.75	-0.565917	4	1	-2.75	-0.483
4	-1	-2.5	-0.596939	4	0	-2.5	-0.51447	4	1	-2.5	-0.432
4	-1	-2.25	-0.545492	4	0	-2.25	-0.463023	4	1	-2.25	-0.380
4	-1	-2.	-0.494045	4	0	-2.	-0.411576	4	1	-2.	-0.329
4	-1	-1.75	-0.442598	4	0	-1.75	-0.360129	4	1	-1.75	-0.277
4	-1	-1.5	-0.391151	4	0	-1.5	-0.308682	4	1	-1.5	-0.226
4	-1	-1.25	-0.339704	4	0	-1.25	-0.257235	4	1	-1.25	-0.174
4	-1	-1.	-0.288257	4	0	-1.	-0.205788	4	1	-1.	-0.123
4	-1	-0.75	-0.23681	4	0	-0.75	-0.154341	4	1	-0.75	-0.071
4	-1	-0.5	-0.185363	4	0	-0.5	-0.102894	4	1	-0.5	-0.020
4	-1	-0.25	-0.133916	4	0	-0.25	-0.051447	4	1	-0.25	0.0310
4	-1	0.	-0.0824692	4	0	0.	0.	4	1	0.	0.0824
4	-1	0.25	-0.0310223	4	0	0.25	0.051447	4	1	0.25	0.1339
4	-1	0.5	0.0204247	4	0	0.5	0.102894	4	1	0.5	0.1853
4	-1	0.75	0.0718717	4	0	0.75	0.154341	4	1	0.75	0.2368
4	-1	1.	0.123319	4	0	1.	0.205788	4	1	1.	0.2882
4	-1	1.25	0.174766	4	0	1.25	0.257235	4	1	1.25	0.3397
4	-1	1.5	0.226213	4	0	1.5	0.308682	4	1	1.5	0.3911
4	-1	1.75	0.27766	4	0	1.75	0.360129	4	1	1.75	0.4425
4	-1	2.	0.329107	4	0	2.	0.411576	4	1	2.	0.4940
4	-1	2.25	0.380554	4	0	2.25	0.463023	4	1	2.25	0.5454
4	-1	2.5	0.432001	4	0	2.5	0.51447	4	1	2.5	0.5969
4	-1	2.75	0.483448	4	0	2.75	0.565917	4	1	2.75	0.6483
4	-1	3.	0.534895	4	0	3.	0.617364	4	1	3.	0.6998

According to above table, the general sol'n is a specific sol'n only when  $c_1 = 0$ . There are a number of instances shown for various values of  $c_2$ . The plot below shows these functions, each of which is a particular sol'n (when  $x = 1$ ).

```
plot3 =
Plot[Evaluate[Table[lint[x, c1, c2], {c1, 0, 0}, {c2, -3, 3, .25}]],
{x, 0, 2}, PlotRange → {-2, 2}, PlotStyle → Thickness[0.003]]
```



What happened? The sol'n found by **DSolve** matches the text's answer. However, the original DE only vanishes when  $x = 1$  &  $c_1 = 0$ .

$$5. \quad 4x^2 y'' + 5y = 0$$

```
ClearAll["Global`*"]
```

```
tier = {4 x^2 y''[x] + 5 y[x] == 0}
{5 y[x] + 4 x^2 y''[x] == 0}
```

```
cins = DSolve[tier, y, x]
```

```
{ {y -> Function[{x}, Sqrt[x] C[2] Cos[Log[x]] + Sqrt[x] C[1] Sin[Log[x]]] } }
```

The above cell matches the text answer.

```
Simplify[tier /. cins]
{{True}}
```

In the above case the sol'n checks out with Mathematica as well as matching the text's answer.

7.  $(x^2 D^2 - 4 x D + 6 I) y = C$

```
ClearAll["Global`*"]
```

```
hol = {x^2 y''[x] - 4 x y'[x] + 6 y[x] == c}
dea = DSolve[hol, y, x]
{6 y[x] - 4 x y'[x] + x^2 y''[x] == c}
```

```
{ {y -> Function[{x}, c/6 + x^2 C[1] + x^3 C[2]] } }
```

```
Simplify[hol /. dea]
{{True}}
```

The above answer matches the text's, with the original rhs c constant equaling zero in the text answer.

9.  $(x^2 D = 0.2 x D + 0.36 I) y = 0$

```
ClearAll["Global`*"]
```

```
reg = {x^2 y''[x] - 0.2 x y'[x] + 0.36 y[x] == 0}
dirk = DSolve[reg, y, x]
{0.36 y[x] - 0.2 x y'[x] + x^2 y''[x] == 0}
```

```
{ {y -> Function[{x}, x^0.6 C[1] + 0.6 x^0.6 C[2] Log[x]] } }
```

```
Chop[Simplify[reg /. dirk], 10^-16]
{{True}}
```

The above answer matches the text's. However, the checking step does not work cleanly.

The default tolerance for **Chop** is  $10^{-10}$ , and in this case  $10^{-16}$  will work. But an exact answer would be preferred.

$$11. (x^2 D^2 - 3 x D + 10 I) y = 0$$

```
ClearAll["Global`*"]
```

```
deam = {x^2 y''[x] - 3 x y'[x] + 10 y[x] == 0}
```

```
vre = DSolve[deam, y, x]
```

```
{10 y[x] - 3 x y'[x] + x^2 y''[x] == 0}
```

```
{ {y -> Function[{x}, x^2 C[2] Cos[√6 Log[x]] + x^2 C[1] Sin[√6 Log[x]]] ] }
```

```
Simplify[deam /. vre]
```

```
{{True}}
```

In this case the substitution worked cleanly. The answer also matches the text's.

12 - 19 Initial value problem  
Solve and graph the solution.

$$13. x^2 y'' + 3 x y' + 0.75 y = 0, y[1] = 1, y'[1] = -1.5$$

```
ClearAll["Global`*"]
```

```
non = {x^2 y''[x] + 3 x y'[x] + 0.75 y[x] == 0, y[1] == 1, y'[1] == -1.5}
```

```
by = DSolve[non, y, x]
```

```
{0.75 y[x] + 3 x y'[x] + x^2 y''[x] == 0, y[1] == 1, y'[1] == -1.5}
```

```
{ {y -> Function[{x}, (1. x^0.5 + 8.88178 × 10^-16 x^1.5) / x^2. ] } }
```

```
rek = Chop[by, 10^-15]
```


```
{ {y -> Function[{x}, (1. x^0.5 + 0 x^1.5) / x^2. ] } }
```

The posed floating point format in the problem created a slight haze which made **Chop** necessary. Allowing this, the answer matches the text's.

```
Simplify[non /. rek]
```

```
{{True, True, True}}
```

```
Plot[y[x] /. rek, {x, 0, 1}, PlotRange -> {0, 100}, ImageSize -> 300]
```



$$15. \quad x^2 y'' + 3xy' + y = 0, \quad y[1] = 3.6, \quad y'[1] = 0.4$$

```
ClearAll["Global`*"]
```

```
hote = {x^2 y''[x] + 3 x y'[x] + y[x] == 0, y[1] == 3.6, y'[1] == 0.4}
```

```
deli = DSolve[hote, y, x]
```


```
{y[x] + 3 x y'[x] + x^2 y''[x] == 0, y[1] == 3.6, y'[1] == 0.4}
```

$$\left\{ \left\{ y \rightarrow \text{Function}\left[ \{x\}, \frac{4 \cdot (0.9 + 1 \cdot \text{Log}[x])}{x} \right] \right\} \right\}$$

```
Simplify[hote /. deli]
```

```
{{True, True, True}}
```

```
Plot[y[x] /. deli, {x, 0, 1}, PlotRange -> {-100, 10}, ImageSize -> 300]
```



The answer matches the text's.

$$17. \quad (x^2 D^2 + x D + I) y = 0, \quad y[1] = 1, \quad y'[1] = 1$$

```
In[185]:= ClearAll["Global`*"]
```



```

In[186]:= zog = {x^2 y''[x] + x y'[x] + y[x] == 0, y[1] == 1, y'[1] == 1}
nilt = DSolve[zog, y, x]

Out[186]= {y[x] + x y'[x] + x^2 y''[x] == 0, y[1] == 1, y'[1] == 1}

Out[187]= {{y -> Function[{x}, Cos[Log[x]] + Sin[Log[x]]]}}

In[188]:= Simplify[zog /. nilt]

Out[188]= {{True, True, True}}


```

The answer matches the text's.

```

In[190]:= Plot[y[x] /. nilt, {x, -1, 1}, PlotRange -> All,
ImageSize -> 300, PlotStyle -> Thickness[0.003]]

```



19.  $(x^2 D^2 - x D - 15 I) y = 0, y[1] = 0.1, y'[1] = -4.5$

```

ClearAll["Global`*"]

wer = {x^2 y''[x] - x y'[x] - 15 y[x] == 0, y[1] == 0.1, y'[1] == -4.5}
jip = DSolve[wer, y, x]

{-15 y[x] - x y'[x] + x^2 y''[x] == 0, y[1] == 0.1, y'[1] == -4.5}

```

```

{{y -> Function[{x}, -  $\frac{0.525 (-1.19048 + 1. x^8)}{x^3}$  ]}}

```

```

Simplify[wer /. jip]
{{x == 0, True, True}}

```

The initial value points are verified. However, the general sol'n is still under a shadow.

```

ard = jip[[1, 1, 2, 2]]
-  $\frac{0.525 (-1.19048 + 1. x^8)}{x^3}$ 

```

```
poi = ExpandNumerator[ard]
```

$$\frac{0.625 - 0.525 x^8}{x^3}$$

```
deke = Expand[poi]
```

$$\frac{0.625}{x^3} - 0.525 x^5$$

The above form matches the text's answer. Below is a series of steps to check the sol'n 'by hand'.

```
mot[x_] :=  $\frac{0.625}{x^3} - 0.5249999999999999 x^5$ 
```

```
mot1 = D[mot[x], x]
```

$$-\frac{1.875}{x^4} - 2.625 x^4$$

```
mot2 = D[mot[x], {x, 2}]
```


$$\frac{7.5}{x^5} - 10.5 x^3$$

```
Simplify[-15 mot[x] - x mot'[x] + x^2 mot''[x]]
```

```
0.
```

The sol'n is checked 'by hand' and found to be correct.

```
Plot[poi, {x, -1, 1}, PlotRange -> {-1000, 1000}, ImageSize -> 300]
```



The vertical line in the plot is an artifact, as the function is undefined at  $x = 0$ .