

BIOS 755: Parametric mean curves

Alexander McLain

January 24, 2023

Parametric Curves

- ▶ An alternative approach for analyzing the parallel-groups repeated measures design is to consider parametric curves for the time trends.
- ▶ We model the means as an explicit function of time
 - ▶ Linear trend
 - ▶ Quadratic Trend
 - ▶ Linear spline

Linear Trend

- ▶ If the means tend to change linearly over time, we can fit the following model:

$$E(Y_{ij}) = \beta_0 + \beta_1 t_j + \beta_2 TRT_i + \beta_3 t_j TRT_i$$

- ▶ Let $Trt=1$ if the subject is in group 1 and 0 otherwise. For subjects in treatment group 1,

$$E(Y_{ij}) = \beta_0 + (\beta_1 + \beta_3)t_j + \beta_2$$

- ▶ For subjects in the control group 0

$$E(Y_{ij}) = \beta_0 + \beta_1 t_j$$

- ▶ Thus, each group's mean is assumed to change linearly over time.

Quadratic Trend

- ▶ If the means tend to change over time in a quadratic manner, we can fit the following model:

$$E(Y_{ij}) = \beta_0 + \beta_1 t_j + \beta_2 t_j^2 + \beta_3 TRT_i + \beta_4 t_j TRT_i + \beta_5 t_j^2 TRT_i$$

- ▶ For subjects in treatment group 1,

$$E(Y_{ij}) = \beta_0 + (\beta_1 + \beta_4)t_j + (\beta_2 + \beta_5)t_j^2 + \beta_3$$

- ▶ For subjects in the control group 0

$$E(Y_{ij}) = \beta_0 + \beta_1 t_j + \beta_2 t_j^2$$

- ▶ Thus, each group's mean is assumed to change quadratically over time.

Quadratic Trend

- ▶ To avoid problems of collinearity in the quadratic (or in any higher-order polynomial) trend model, should always “center” t_j on its mean prior to the analysis (i.e. replace t_j by its deviation from the mean).
- ▶ For example, suppose $t_j = (1, 2, \dots, 10)$. Then the correlation between t_j and t_j^2 is 0.975.
- ▶ However, if we create a “centered” variable, say,

$$t_j^* = t_j - \bar{t}$$

then the correlation between t_j^* and t_j^{*2} is 0.

Log Trend

- ▶ If the means are monotonically increasing (or decreasing), but there appears to be “diminishing returns” in the changes then a log transformation can be used

$$E(Y_{ij}) = \beta_0 + \beta_1 \log(t_j) + \beta_2 TRT_i + \beta_3 \log(t_j) TRT_i$$

- ▶ For subjects in treatment group 1,


$$E(Y_{ij}) = \beta_0 + (\beta_1 + \beta_3) \log(t_j) + \beta_2$$

- ▶ For subjects in the control group 0


$$E(Y_{ij}) = \beta_0 + \beta_1 \log(t_j)$$

- ▶ The square root function \sqrt{t} is another option for “diminishing returns”.

- We can also add a linear trend to a log transformation.




Square-root transformation



Linear Spline

- If the means change over time in a piecewise linear manner, we can fit the following linear spline model with knot at t^* :

$$\begin{aligned} E(Y_{ij}) &= \beta_0 + \beta_1 t_j + \beta_2 TRT_i + \beta_3 t_j TRT_i \quad t_j \leq t^* \\ &= \beta_0 + \beta_1 t^* + \beta_2 TRT_i + \beta_3 t^* TRT_i + \beta_4 (t_j - t^*) + \beta_5 (t_j - t^*) TRT_i \quad t_j > t^* \end{aligned}$$



Linear Spline

- For subjects in group 1


$$\begin{aligned} E(Y_{ij}) &= \beta_0 + \beta_2 + (\beta_1 + \beta_3)t_j \quad t_j \leq t^* \\ &= \beta_0 + \beta_2 + (\beta_1 + \beta_3)t^* + (\beta_4 + \beta_5)(t_j - t^*) \quad t_j > t^* \end{aligned}$$

- For subjects in group 2

$$\begin{aligned} E(Y_{ij}) &= \beta_0 + \beta_1 t_j \quad t_j \leq t^* \\ &= \beta_0 + \beta_1 t^* + \beta_4(t_j - t^*) \quad t_j > t^* \end{aligned}$$

Non-linear Spline

- ▶ Non-linear splines (e.g., cubic B-splines can also be used).



Summary of Features of Parametric Curve Models

- ▶ Allows one to model time trend and treatment effect(s) as a function of a small number of parameters. That is, the treatment effect can be captured in one or two parameters, leading to more powerful tests when these models fit the data.
- ▶ Since $E(Y_{ij})$ is defined as an explicit function of the time of measurement, t_j , there is no reason to require all subjects to have the same set of measurement times, nor even the same number of measurements.
- ▶ May not always be possible to fit data adequately.
- ▶ Finally, note that the aforementioned parametric curve analyses can all be expressed in terms of the general linear model.