



DR. HARSHIL SHAH
ORTHOPAEDIC SURGEON

SHOULDER ARTHROSCOPY

Expert care for rotator cuff tears,
instability and sports injuries.



Minimally Invasive
Small incisions.
Less pain.



Precision Diagnosis
Advanced imaging.
Accurate treatment.



Faster Recovery
Structured rehab.
Stronger outcomes.



Personalised Care
Tailored to you.
Focused on function.



Book a Consultation
+91 98765 43210



www.drharshilshah.com

ROTATOR CUFF TEARS

Symptoms & why timely treatment matters



COMMON SYMPTOMS



Pain with overhead activity

Reaching, lifting or lying on the affected side.



Weakness

Difficulty lifting or rotating the arm.



Night pain

Pain that disturbs sleep.



Stiffness

Loss of motion, especially in the morning.



Clicking or catching

Grinding or popping during movement.


ROTATOR CUFF

Supraspinatus

Infraspinatus

Teres Minor

Subscapularis



WHY TIMELY TREATMENT MATTERS



Prevent tear progression



Restore strength and function



Avoid long-term shoulder arthritis



DR. HARSHIL SHAH
ORTHOPAEDIC SURGEON

HOW A CUFF PROBLEM CAN PROGRESS



Rotator cuff pain & guarding

Pain limits daily activities.
Early rest and therapy can help.



Frozen shoulder

Capsule tightness causes stiffness
and loss of motion.



Cuff tear arthropathy

Chronic tears lead to cartilage wear,
bone changes and weakness.



Advanced stage Reverse shoulder replacement

When damage is severe, a reverse
shoulder replacement can restore
function and reduce pain.



Early diagnosis. Timely treatment. Better outcomes.
Protect your shoulder for the long run.

INSTABILITY AFTER DISLOCATION

Understanding the key structures



Bankart Lesion

Tear of the anterior labrum caused by dislocation.



Hill-Sachs Lesion

Compression injury on the back of the humeral head.



Remplissage

Filling the Hill-Sachs defect with soft tissue to prevent recurrence.



SLAP Tears

Superior labrum injury affecting overhead athletes.



Proper evaluation helps choose the right surgical technique for stability and performance.



DR. HARSHIL SHAH
ORTHOPAEDIC SURGEON

REHABILITATION FOR STRONGER RECOVERY

Guided. Progressive. Goal-oriented.



After Rotator Cuff Repair

Phase 1

Protection
0-4 weeks



Protect the repair.
Passive motion.
Pain control.

Phase 2

Active-Assisted
4-8 weeks



Restore range.
Improve mobility.
Light activities.

Phase 3

Strengthening
8-16 weeks



Build strength.
Endurance and
scapular control.

Phase 4

Return to Activity
4-6 months



Sport-specific.
Overhead strength.
Return to sport.

After Bankart Repair (with or without remplissage)

0-4 weeks

Protection

Sling for passive
movement.



4-8 weeks

Motion

Restore range
of motion.



8-16 weeks

Strengthening

Build shoulder
stability.



4-6+ months

Return to Sport

Gradual return
to play.



Consistent rehab is the key to a pain-free,
stable and strong shoulder.

YOUR QUESTIONS. OUR ANSWERS.



Do I need surgery for every tear?

No. Small tears or early cases can improve with physiotherapy and activity modification.



Can a small cuff tear avoid surgery?

Yes. Many small tears respond well to non-surgical treatment.



How long should I use a sling?

Usually 2–4 weeks depending on the procedure and healing.



When can I return to sport?

Typically 4–6 months, depending on surgery type and your progress.



KEY TAKEAWAYS



Early care
prevents
progression



Expert surgery
restores strength
and stability



Structured rehab
ensures lasting
recovery



Return to life
with confidence
and comfort



DR. HARSHIL SHAH
ORTHOPAEDIC SURGEON



Book a Consultation
+91 98765 43210



www.drharshilshah.com