

DSA 8070 R Session 2: Matrix Algebra

Whitney

August 28, 2021

Contents

| | |
|---|---|
| Motor Trend Car Road Tests Data | 1 |
| Mean Vector and Covariance Matrix | 1 |
| Inverse Matrix | 2 |
| Orthogonal Matrix Example | 2 |
| Eigenvalues and Eigenvectors | 2 |
| Spectral Decomposition | 3 |
| Determinant and Trace | 3 |
| Square-Root Matrices | 4 |
| Partitioning Random vectors | 4 |

Motor Trend Car Road Tests Data

```
data(mtcars)
vars <- which(names(mtcars) %in% c("mpg", "disp", "hp", "drat", "wt"))
cars <- mtcars[, vars]
```

Mean Vector and Covariance Matrix

```
(mean <- apply(cars, 2, mean))
```

```
##      mpg      disp      hp      drat      wt
## 20.090625 230.721875 146.687500  3.596563  3.217250
```

```
n <- dim(cars)[1]; p <- dim(cars)[2]
X <- as.matrix(cars)
ones <- rep(1, n)
(meanCal <- (1 / n) * t(X) %*% ones)
```

```
##      [,1]
## mpg  20.090625
## disp 230.721875
## hp   146.687500
## drat  3.596563
## wt   3.217250
```

```
(S <- cov(cars))
```

```
##      mpg      disp      hp      drat      wt
## mpg  36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
```

```
## disp -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## hp -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## drat 2.195064 -47.06402 -16.45111 0.2858814 -0.3727207
## wt -5.116685 107.68420 44.19266 -0.3727207 0.9573790

(Scal <- (1 / (n - 1)) * t(X) %>% (diag(n) - (1 / n) * ones %>% t(ones)) %>% X)
```

```
##          mpg          disp          hp          drat          wt
## mpg    36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
## disp -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## hp    -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## drat   2.195064 -47.06402 -16.45111  0.2858814 -0.3727207
## wt     -5.116685 107.68420 44.19266 -0.3727207  0.9573790
```

Inverse Matrix

```
S_inv <- solve(S)
(S_inv %>% S)
```

```
##          mpg          disp          hp          drat          wt
## mpg    1.000000e+00 2.842171e-14 7.105427e-15 -1.110223e-16 2.220446e-16
## disp  2.775558e-17 1.000000e+00 -6.661338e-16 -1.734723e-18 -1.387779e-17
## hp    -2.775558e-17 4.440892e-16 1.000000e+00 0.000000e+00 0.000000e+00
## drat -7.549517e-15 1.350031e-13 -1.421085e-14 1.000000e+00 1.165734e-15
## wt     0.000000e+00 1.136868e-13 0.000000e+00 0.000000e+00 1.000000e+00
```

Orthogonal Matrix Example

```
Q <- matrix(c(2, 1, 2, -2, 2, 1, 1, 2, -2), ncol = 3) / 3
(Q %>% t(Q))
```

```
##      [,1] [,2] [,3]
## [1,]  1   0   0
## [2,]  0   1   0
## [3,]  0   0   1
```

Eigenvalues and Eigenvectors

```
eigen <- eigen(S)
(S %>% eigen$vectors[, 1] / eigen$vectors[, 1])
```

```
##      [,1]
## mpg 18636.79
## disp 18636.79
## hp 18636.79
## drat 18636.79
## wt 18636.79
```

```
eigen$values[1]
```

```
## [1] 18636.79
t(eigen$eigenvectors[, 1]) %*% eigen$eigenvectors[, 1]

##      [,1]
## [1,]    1
```

Spectral Decomposition

```
temp <- array(dim = c(5, 5, 5))

for (i in 1:5){
  temp[i,,] <- eigen$values[i] * eigen$eigenvectors[, i] %*% t(eigen$eigenvectors[, i])
}
# Check the spectral decomposition
(out <- apply(temp, 2:3, sum))
```

```
##      [,1]      [,2]      [,3]      [,4]      [,5]
## [1,] 36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
## [2,] -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## [3,] -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## [4,]  2.195064  -47.06402  -16.45111  0.2858814  -0.3727207
## [5,]  -5.116685  107.68420  44.19266  -0.3727207  0.9573790
```

S

```
##      mpg      disp      hp      drat      wt
## mpg    36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
## disp -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## hp    -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## drat   2.195064  -47.06402  -16.45111  0.2858814  -0.3727207
## wt    -5.116685  107.68420  44.19266  -0.3727207  0.9573790
```

Determinant and Trace

```
# Trace
(trace <- sum(diag(S)))
```

```
## [1] 20099.23
```

```
sum(eigen$values)
```

```
## [1] 20099.23
```

```
# Determinant
det(S)
```

```
## [1] 3951786
```

```
prod(eigen$values)
```

```
## [1] 3951786
```

Square-Root Matrices

```
temp1 <- array(dim = c(5, 5, 5))

for (i in 1:5){
  temp1[i,,] <- (1 / eigen$values[i]) * eigen$vectors[, i] %*% t(eigen$vectors[, i])
}
# Check the spectral decomposition
(out1 <- apply(temp1, 2:3, sum))
```

```
##           [,1]      [,2]      [,3]      [,4]      [,5]
## [1,]  0.1695494031 -0.0006468718  0.0058975274 -0.29977161  0.58997555
## [2,] -0.0006468718  0.0005369064 -0.0003801427  0.02257595 -0.03751089
## [3,]  0.0058975274 -0.0003801427  0.0008208474 -0.02678451  0.02595898
## [4,] -0.2997716134  0.0225759526 -0.0267845083  8.50376340  0.40558365
## [5,]  0.5899755523 -0.0375108878  0.0259589804  0.40558365  7.37641228
```

S_inv

```
##           mpg      disp      hp      drat      wt
## mpg  0.1695494031 -0.0006468718  0.0058975274 -0.29977161  0.58997555
## disp -0.0006468718  0.0005369064 -0.0003801427  0.02257595 -0.03751089
## hp    0.0058975274 -0.0003801427  0.0008208474 -0.02678451  0.02595898
## drat -0.2997716134  0.0225759526 -0.0267845083  8.50376340  0.40558365
## wt    0.5899755523 -0.0375108878  0.0259589804  0.40558365  7.37641228
```

```
temp2 <- array(dim = c(5, 5, 5))

for (i in 1:5){
  temp2[i,,] <- sqrt(eigen$values[i]) * eigen$vectors[, i] %*% t(eigen$vectors[, i])
}

out2 <- apply(temp2, 2:3, sum)

(out2 %*% out2)
```

```
##           [,1]      [,2]      [,3]      [,4]      [,5]
## [1,]  36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
## [2,] -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## [3,] -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## [4,]  2.195064 -47.06402 -16.45111  0.2858814 -0.3727207
## [5,] -5.116685 107.68420 44.19266 -0.3727207  0.9573790
```

S

```
##           mpg      disp      hp      drat      wt
## mpg  36.324103 -633.09721 -320.73206  2.1950635 -5.1166847
## disp -633.097208 15360.79983 6721.15867 -47.0640192 107.6842040
## hp   -320.732056 6721.15867 4700.86694 -16.4511089 44.1926613
## drat  2.195064 -47.06402 -16.45111  0.2858814 -0.3727207
## wt   -5.116685 107.68420 44.19266 -0.3727207  0.9573790
```

Partitioning Random vectors

Let's partition the variables into two groups

1. *disp*, *hp*, *wt*
2. *mpg*, *drat*

```
vars1 <- which(names(mtcars) %in% c("disp", "hp", "wt"))
vars2 <- which(names(mtcars) %in% c("mpg", "drat"))
```

```
carPar <- mtcars[, c(vars1, vars2)]
```

```
(Sigma11 <- cov(carPar[1:3, 1:3]))
```

```
##           disp           hp           wt
## disp 901.3333 294.6667 7.410000
## hp   294.6667  96.33333 2.422500
## wt    7.4100   2.42250 0.077175
```

```
(Sigma22 <- cov(carPar[4:5, 4:5]))
```

```
##           mpg           drat
## mpg   3.6450 -0.09450
## drat -0.0945  0.00245
```

```
(Sigma12 <- cov(carPar[1:3, 4:5]))
```

```
##           mpg           drat
## mpg   1.08 -0.0300000000
## drat -0.03  0.0008333333
```