

Lecture 25

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

Text: Chapter 11

STAT 8010 Statistical Methods I
April 21, 2020

Whitney Huang
Clemson University

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class


Analysis of Variance (ANOVA) Approach to Regression

25.1

Notes

Understanding Confidence Intervals

- Suppose $Y = \beta_0 + \beta_1 X + \epsilon$, where $\beta_0 = 3$, $\beta_1 = 1.5$ and $\sigma^2 \sim N(0, 1)$
- We take 100 random sample each with sample size 20
- We then construct the 95% CI for each random sample (\Rightarrow 100 CIs)



Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY


Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.2

Notes

Confidence Intervals vs. Prediction Intervals



Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.3

Notes

Analysis of Variance (ANOVA) Approach to Regression

Partitioning Sums of Squares

- Total sums of squares in response

$$SST = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

- We can rewrite SST as

$$\begin{aligned} \sum_{i=1}^n (Y_i - \bar{Y})^2 &= \sum_{i=1}^n (Y_i - \hat{Y}_i + \hat{Y}_i - \bar{Y})^2 \\ &= \underbrace{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}_{\text{Error}} + \underbrace{\sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2}_{\text{Model}} \end{aligned}$$


Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination



Review of Last Class
Analysis of Variance (ANOVA) Approach to Regression

Notes

Partitioning Total Sums of Squares



Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination



Review of Last Class
Analysis of Variance (ANOVA) Approach to Regression

Notes

Total Sum of Squares: SST

- If we ignored the predictor X , the \bar{Y} would be the best (linear unbiased) predictor

$$Y_i = \beta_0 + \varepsilon_i \tag{1}$$

- SST is the sum of squared deviations for this predictor (i.e., \bar{Y})
- The **total mean square** is $SST/(n - 1)$ and represents an unbiased estimate of σ^2 under the model (1).

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination



Review of Last Class
Analysis of Variance (ANOVA) Approach to Regression

Notes

Regression Sum of Squares: SSR

- SSR: $\sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$
- Degrees of freedom is 1 due to the inclusion of the slope, i.e.,
$$Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i \tag{2}$$
- “Large” MSR = SSR/1 suggests a linear trend, because

$$E[MSE] = \sigma^2 + \beta_1^2 \sum_{i=1}^n (X_i - \bar{X})^2$$

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.7

Notes

Error Sum of Squares: SSE

- SSE is simply the sum of squared residuals
$$SSE = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$$
- Degrees of freedom is $n - 2$ (Why?)
- SSE large when |residuals| are “large” $\Rightarrow Y_i$ ’s vary substantially around fitted regression line
- MSE = SSE/($n - 2$) and represents an unbiased estimate of σ^2 **when taking X into account**

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.8

Notes

ANOVA Table and F test

Source	df	SS	MS
Model	1	$SSR = \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$	$MSR = SSR/1$
Error	$n - 2$	$SSE = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2$	$MSE = SSE/(n-2)$
Total	$n - 1$	$SST = \sum_{i=1}^n (Y_i - \bar{Y})^2$	

- **Goal:** To test $H_0 : \beta_1 = 0$
- Test statistics $F^* = \frac{MSR}{MSE}$
- If $\beta_1 = 0$ then F^* should be near one \Rightarrow reject H_0 when F^* “large”
- We need sampling distribution of F^* under $H_0 \Rightarrow F_{1,n-2}$, where $F(d_1, d_2)$ denotes a F distribution with degrees of freedom d_1 and d_2

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.9

Notes


F Test: $H_0 : \beta_1 = 0$ vs. $H_a : \beta_1 \neq 0$

```
fit <- lm(MaxHeartRate ~ Age)
anova(fit)
```
```

Analysis of Variance Table

Response: MaxHeartRate

|           | Df | Sum Sq    | Mean Sq | F value |
|-----------|----|-----------|---------|---------|
| Age       | 1  | 2724.50   | 2724.50 | 130.01  |
| Residuals | 13 | 272.43    | 20.96   |         |
|           |    | Pr(>F)    |         |         |
| Age       |    | 3.848e-08 | ***     |         |



Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.10

Notes

---

---

---

---

---

---

---

---

SLR: F-Test vs. T-test

ANOVA Table and F-Test

Analysis of Variance Table

Response: MaxHeartRate

|           | Df | Sum Sq  | Mean Sq   |
|-----------|----|---------|-----------|
| Age       | 1  | 2724.50 | 2724.50   |
| Residuals | 13 | 272.43  | 20.96     |
|           |    | F value | Pr(>F)    |
| Age       |    | 130.01  | 3.848e-08 |

Parameter Estimation and T-Test

Coefficients:

|             | Estimate  | Std. Error | t value | Pr(> t ) |
|-------------|-----------|------------|---------|----------|
| (Intercept) | 210.04846 | 2.86694    | 73.27   | < 2e-16  |
| Age         | -0.79773  | 0.06996    | -11.40  | 3.85e-08 |

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.11

Notes

---

---

---

---

---

---

---

---

Correlation and Simple Linear Regression

- **Pearson Correlation:**  $r = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2 \sum_{i=1}^n (Y_i - \bar{Y})^2}}$
- $-1 \leq r \leq 1$  measures the strength of the **linear relationship** between  $Y$  and  $X$
- We can show

$$r = \hat{\beta}_1 \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2}}$$

this implies

$$\beta_1 = 0 \text{ in SLR} \Leftrightarrow \rho = 0$$

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.12

Notes

---

---

---

---

---

---

---

---

## Coefficient of Determination $R^2$

- Defined as the proportion of total variation **explained** by SLR

$$R^2 = \frac{\sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2} = \frac{SSR}{SST} = 1 - \frac{SSE}{SST}$$

- We can show  $r^2 = R^2$ :

$$\begin{aligned} r^2 &= \left( \hat{\beta}_{1,LS} \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2}} \right)^2 \\ &= \frac{\hat{\beta}_{1,LS}^2 \sum_{i=1}^n (X_i - \bar{X})^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2} \\ &= \frac{SSR}{SST} \\ &= R^2 \end{aligned}$$

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.13

## Notes

## Maximum Heart Rate vs. Age: $r$ and $R^2$

```
> summary(fit)$r.squared
[1] 0.9090967
> cor(Age, MaxHeartRate)
[1] -0.9534656
```

### Interpretation:

There is a strong negative linear relationship between MaxHeartRate and Age. Furthermore, ~91% of the variation in MaxHeartRate can be explained by Age.

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY



Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.14

## Notes

## Residual Plot Revisited



⇒ Nonlinear relationship

⇒ Non-constant variance

- Transform  $X$
- Transform  $Y$
- Nonlinear regression
- Weighted least squares

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY


Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.15

## Notes

Extrapolation in SLR



Extrapolation beyond the range of the given data can lead to **seriously biased estimates** if the **assumed relationship does not hold the region of extrapolation**

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.16

Notes

---

---

---

---

---

---

---

---

Summary of SLR

- **Model:**  $Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i$
- **Estimation:** Use the **method of least squares** to estimate the parameters
- **Inference**
  - Hypothesis Testing
  - Confidence/prediction Intervals
  - ANOVA
- **Model Diagnostics and Remedies**

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.17

Notes

---

---

---

---

---

---

---

---

Summary

In this lecture, we learned **ANOVA Approach to Regression** and **Coefficient of Determination**

Next time: **Review**

Simple Linear Regression: ANOVA Approach to Regression and Coefficient of Determination

CLEMSON UNIVERSITY

Review of Last Class

Analysis of Variance (ANOVA) Approach to Regression

25.18

Notes

---

---

---

---

---

---

---

---