

Lecture 13

Continuous Random Variables

Text: Chapter 4

STAT 8010 Statistical Methods I
September 18, 2019

Whitney Huang
Clemson University

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.1

Notes

Agenda

- 1 From Discrete to Continuous Random Variables
- 2 Cumulative Distribution Functions
- 3 Expected Value and Variance
- 4 Normal Distributions

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions


Expected Value and Variance

Normal Distributions

13.2

Notes

Probability Mass Functions vs. Probability Density Functions



Remarks:

- pmf assigns probabilities to each possible values of a discrete distribution
- pdf describes the relative likelihood for this random variable to take on a given interval

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.3

Notes


Probability Mass Functions v.s. Probability Density Functions cont'd

Recall the properties of discrete probability mass functions (Pmfs):

- $0 \leq p_X(x) \leq 1$ for all possible values of x
- $\sum_x p_X(x) = 1$
- $\mathbb{P}(a \leq X \leq b) = \sum_{x=a}^{x=b} p_X(x)$

For continuous distributions, the properties for probability density functions (Pdf's) are similar:

- $f_X(x) \geq 0$ for all possible values of x
- $\int_{-\infty}^{\infty} f_X(x) dx = 1$
- $\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$



Notes


Cumulative Distribution Functions (cdfs) for Continuous Distribution

- The cdf $F_X(x)$ is defined as

$$F_X(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f_X(x) dx$$
- we use cdf to calculate probabilities of a continuous random variable within an interval, i.e.

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx = \int_{-\infty}^b f_X(x) dx - \int_{-\infty}^a f_X(x) dx = \boxed{F_X(b) - F_X(a)}$$

Remark: $\mathbb{P}(X = x) = \int_x^x f_X(x) dx = 0$ for all possible values of x



Notes


Expected Value and Variance

Recall the expected value formula for the discrete random variable: $\mathbb{E}[X] = \sum_x x p_X(x)$

For continuous random variables, we have similar formulas:

Let a , b , and c are constant real numbers

- $\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx$
- $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[cX] = c\mathbb{E}[X]$
- $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$
- $\text{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} x^2 f_X(x) dx - \left(\int_{-\infty}^{\infty} x f_X(x) dx \right)^2$
- $\text{Var}(cX) = c^2 \text{Var}(X)$
- $\text{Var}(X - c) = \text{Var}(X)$



Notes

Example

Let X represent the diameter in inches of a circular disk cut by a machine. Let $f_X(x) = c(4x - x^2)$ for $1 \leq x \leq 4$ and be 0 otherwise. Answer the following questions:

- 1 Find the value of c that makes this a valid pdf
- 2 Find the expected value and variance of X
- 3 What is the probability that X is within .5 inches of the expected diameter?
- 4 Find $F_X(x)$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.7

Notes

Normal Distribution

Characteristics of the Normal random variable:
Let X be a Normal r.v.

- The support for X : $(-\infty, \infty)$
- Its parameter(s) and definition(s): μ : mean and σ^2 : variance
- The probability density function (pdf): $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$ for $-\infty < x < \infty$
- The cumulative distribution function (cdf): No explicit form, look at the value $\Phi(\frac{x-\mu}{\sigma})$ for $-\infty < x < \infty$ from standard normal table
- The expected value: $E[X] = \mu$
- The variance: $Var(X) = \sigma^2$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions


Expected Value and Variance

Normal Distributions

13.8

Notes

Normal Density Curves



- The parameter μ determines the center of the distribution
- The parameter σ^2 determines the spread of the distribution
- Also called bell-shaped distribution

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.9

Notes

Standard Normal $Z \sim N(\mu = 0, \sigma^2 = 1)$

- Normal random variable X with mean μ and standard deviation σ can convert to standard normal Z by the following :

$$Z = \frac{X - \mu}{\sigma}$$

- The cdf of the standard normal, denoted by $\Phi(z)$, can be found from the standard normal table
- The probability $\mathbb{P}(a \leq X \leq b)$ where $X \sim N(\mu, \sigma^2)$ can be compute

$$\begin{aligned} \mathbb{P}(a \leq X \leq b) &= \mathbb{P}\left(\frac{a - \mu}{\sigma} \leq Z \leq \frac{b - \mu}{\sigma}\right) \\ &= \Phi\left(\frac{b - \mu}{\sigma}\right) - \Phi\left(\frac{a - \mu}{\sigma}\right) \end{aligned}$$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.10

Notes

Properties of Φ

- $\Phi(0) = .50 \Rightarrow$ Mean and Median (50_{th} percentile) for standard normal are both 0
- $\Phi(-z) = 1 - \Phi(z)$
- $\mathbb{P}(Z > z) = 1 - \Phi(z) = \Phi(-z)$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.11

Notes

Example

Let us examine Z . Find the following probabilities with respect to Z :

- 1 Z is at most -1.75
- 2 Z is between -2 and 2 inclusive
- 3 Z is less than $.5$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.12

Notes

Example Cont'd

Solution.

- 1 $\mathbb{P}(Z \leq -1.75) = \Phi(-1.75) = .0401$
- 2 $\mathbb{P}(-2 \leq Z \leq 2) = \Phi(2) - \Phi(-2) = .9772 - .0228 = .9544$
- 3 $\mathbb{P}(Z < .5) = \Phi(.5) = .6915$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.13

Notes

Sums of Normal Random Variables

If X_i $1 \leq i \leq n$ are independent normal random variables with mean μ_i and variance σ_i^2 , respectively.

- 1 Let $S_n = \sum_{i=1}^n X_i$ then $S_n \sim N(\sum_{i=1}^n \mu_i, \sum_{i=1}^n \sigma_i^2)$
- 2 This can be applied for any integer n

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.14

Notes

Example

Let X_1 , X_2 , and X_3 be mutually independent, Normal random variables. Let their means and standard deviations be $3k$ and k for $k = 1, 2$, and 3 respectively. Find the following distributions:

- 1 $\sum_{i=1}^3 X_i$
- 2 $X_1 + 2X_2 - 3X_3$
- 3 $X_1 + 5X_3$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.15

Notes

Example Cont'd

Solution.

- 1. $\sum_{i=1}^3 X_i \sim N(\mu = 3 + 6 + 9 = 15, \sigma^2 = 1^2 + 2^2 + 3^2 = 14)$
- 2. $X_1 + 2X_2 - 3X_3 \sim N(\mu = 3 + 12 - 27 = -12, \sigma^2 = 1^2 + 4 \times 2^2 + 9 \times 3^2 = 98)$
- 3. $X_1 + 5X_3 \sim N(\mu = 3 + 45 = 48, \sigma^2 = 1^2 + 25 \times 3^2 = 226)$

Continuous Random Variables

CLEMSON UNIVERSITY

From Discrete to Continuous Random Variables

Cumulative Distribution Functions

Expected Value and Variance

Normal Distributions

13.16

Notes

Notes

Notes
